

AGRICULTURE WATER DEMAND MODEL

Report for Regional District of Nanaimo

May 2013



AGRICULTURE WATER DEMAND MODEL

Report for Regional District of Nanaimo

Authors

Ted van der Gulik, P.Eng.

Senior Engineer
B.C. Ministry of Agriculture
Sustainable Agriculture Management Branch
Abbotsford, BC

Denise Neilsen, Ph.D.

Research Scientist
Agriculture and Agri-Food Canada
Pacific Agri-Food Research Centre
Summerland, BC

Ron Fretwell

Program Developer
RHF Systems Ltd.
Kelowna, BC

Stephanie Tam, P.Eng.

Water Management Engineer
B.C. Ministry of Agriculture
Sustainable Agriculture Management Branch
Abbotsford, BC

Partially Funded By



BRITISH COLUMBIA
CONSERVATION
FOUNDATION



REGIONAL
DISTRICT
OF NANAIMO

Agriculture and Agri-Food Canada 

May 2013

DISCLAIMER

The data that is presented in this report provides the best estimates for agriculture water demand that can be generated at this time. While every effort has been made to ensure the accuracy and completeness of the information, the information should not be considered as final. The Government of Canada, the BC Ministry of Agriculture, and the BC Agriculture Council or its directors, agents, employees, or contractors will not be liable for any claims, damages, or losses of any kind whatsoever arising out of the use of, or reliance upon, this information.

Table of Contents

ACKNOWLEDGEMENTS	5
BACKGROUND.....	6
METHODOLOGY	7
Cadastre	7
Land Use Survey	8
Soil Information.....	10
Climate Information.....	11
MODEL CALCULATIONS.....	12
Crop	12
Irrigation	12
Soil.....	13
Climate.....	13
Agricultural Water Demand Equation.....	13
LIVESTOCK WATER USE.....	18
DEFINITION AND CALCULATION OF INDIVIDUAL TERMS USED IN THE IRRIGATION WATER DEMAND EQUATION	19
Growing Season Boundaries	19
Evapotranspiration (ET _o).....	21
Availability Coefficient (AC).....	21
Rooting Depth (RD)	21
Stress Factor (stressFactor)	22
Available Water Storage Capacity (AWSC).....	22
Maximum Soil Water Deficit (MSWD).....	22
Deep Percolation Factor (soilPercFactor).....	22
Maximum Evaporation Factor (maxEvaporation).....	23
Irrigation Efficiency (I _e).....	23
Soil Water Factor (swFactor)	23
Early Season Evaporation Factor (earlyEvaporationFactor).....	23
Crop Coefficient (K _c)	23
Growing Degree Days (GDD)	24
Frost Indices	24
Corn Heat Units (CHU)	24
Corn Season Start and End.....	25
Tsum Indices.....	25
Wet/Dry Climate Assessment	25
Groundwater Use.....	25

LAND USE RESULTS.....	26
AGRICULTURAL WATER DEMAND MODEL RESULTS.....	29
Annual Crop Water Demand – Tables A and B	29
Annual Water Demand Reported by Irrigation System – Table C	29
Annual Water Demand by Soil Texture – Table D	29
Annual Water Demand by Aquifer – Table E	30
Irrigated Area by Local Government – Table F	30
Irrigation Management Factors – Table G	30
Deep Percolation – Table H.....	31
Improved Irrigation Efficiency and good Management – Table I.....	31
Water Demand for Frost Protection, Harvesting and Other – Table J	31
Livestock Water Use – Table K.....	32
Climate Change Water Demand for 2050 – Table L.....	32
Agricultural Buildout Crop Water Demand Using 2003 Climate Data – Table M	33
Agricultural Buildout Crop Water Demand for 2050 – Table N	35
Irrigation Systems Used for the Buildout Scenario – Table O.....	35
Water Demand by Aquifer for the Buildout Scenario – Table P	35
Water Demand by Local Government for the Buildout Scenario – Table Q	35
LITERATURE	36
APPENDIX TABLES	37

List of Figures

Figure 1	Map of ALR in Regional District of Nanaimo.....	6
Figure 2	Overlaid Survey Map Sheets, Regional District of Nanaimo.....	7
Figure 3	Land Use Survey.....	8
Figure 4	GIS Map Sheet	8
Figure 5	Cadastre with Polygon	9
Figure 6	GIS Model Graphic.....	10
Figure 7	Nanaimo Area Climate Stations.....	11
Figure 8	Higher Productive Groundwater Aquifers in RDN	28
Figure 9	Annual ET and Effective Precipitation in 2050's	32
Figure 10	Future Irrigation Demand for All Outdoor Uses in the Okanagan in Response to Observed Climate Data (Actuals) and Future Climate Data Projected from a Range of Global Climate Models.....	33
Figure 11	RDN Irrigation Expansion Potential.....	35

List of Tables

Table 1	Livestock Water Demand (Litres/day)	18
Table 2	Overview of RDN's Land and Inventoried Area.....	26
Table 3	Summary of Primary Agricultural Activities within the ALR where Primary Land Use is Agriculture in RDN.....	27
Table 4	Irrigation Management Factors.....	30

Acknowledgements

There are many people that have been involved with the preparation and collection of data used in the development of the Water Demand Model for the Regional District of Nanaimo. The authors wish to express appreciation to the following for their efforts and/or in-kind contribution for the tasks noted.

PROFESSIONALS AND CONTRACTORS

Alex Cannon	Environment Canada	Climate data downscaling
Sam Lee	Ministry of Agriculture	GIS Coordination and report preparation
Corrine Roesler	Ministry of Agriculture	GIS Coordination
Linda Hokanson	Ministry of Agriculture	Publication formatting
Julie Mundy	Ministry of Agriculture	Land Use Survey
Kelsey Lang	Ministry of Agriculture	Land Use Inventory
Michael Dykes	Ministry of Agriculture	Land Use Inventory
Wayne Haddow	Ministry of Agriculture	Land Use Inventory
Jill Hatfield	Ministry of Agriculture	Land Use Inventory
Andrea Lawseth	Contractor	Land Use Survey

COVER PHOTO

The scenic photo of an ocean view in the regional district of Nanaimo in British Columbia, Canada was provided by Scott and Laurie Johnson of Toad Hollow Photography located in Duncan, BC.

The authors thank them for their gift of the beautiful photograph for our front cover.

IN-KIND CONTRIBUTION

The authors would like to express their gratitude to the Partnership of Water Sustainability of B.C. for their generous administration of funding for this project.



Background

The Agriculture Water Demand Model (AWDM) was originally developed in the Okanagan Watershed. It was developed in response to rapid population growth, drought conditions from climate change, and the overall increased demand for water. Many of the watersheds in British Columbia (BC) are fully allocated or will be in the next 15 to 20 years. The AWDM helps to understand current agricultural water use and helps to fulfil the Province’s commitment under the “*Living Water Smart – BC Water Plan*” to reserve water for agricultural lands. The Model can be used to establish agricultural water reserves throughout the various watersheds in BC by providing current and future agriculture water use data.

Climate change scenarios developed by the University of British Columbia (UBC) and the Pacific Agri-Food Research Centre (PARC) in Summerland predict an increase in agricultural water demand due to warmer and longer summers and lower precipitation during summer months in the future.

The Agriculture Water Demand Model was developed to provide current and future agricultural water demands. The Model calculates water use on a property-by-property basis, and sums each property to obtain a total water demand for the entire basin or each sub-basin. Crop, irrigation system type, soil texture and climate data are used to calculate the water demand. Climate data from 2003 was used to present information on one of the hottest and driest years on record and 1997 data was used to represent a wet year. Lands within the Agriculture Land Reserve (ALR), depicted in green in Figure 1 were included in the project.




Figure 1 Map of ALR in Regional District of Nanaimo

Methodology

The Model is based on a Geographic Information System (GIS) database that contains information on cropping, irrigation system type, soil texture and climate data. An explanation of how information was compiled for each is given below. The survey area included all properties within the ALR and areas that were zoned for agriculture by the local government. The inventory was undertaken by Ministry of Agriculture (AGRI) staff, hired professional contractors and summer students. Figure 2 provides a schematic of the map sheets that were generated to conduct the survey.




Figure 2 Overlaid Survey Map Sheets, Regional District of Nanaimo

Cadastre

Cadastre information was provided by the Regional District of Nanaimo (RDN). The entire regional district is covered in one dataset which allows the Model to report out on each sub-basin, local government, water purveyor or groundwater aquifer. A GIS technician used aerial photographs to conduct an initial review of cropping information by cadastre, and divided the cadastre into polygons that separated farmstead and driveways from cropping areas. Different crops were also separated into different polygons if the difference could be identified on the aerial photographs. This data was entered into the database that was used by the field teams to conduct and complete the land use survey.

Land Use Survey

The survey maps and database were created by AGRI for the survey crew to enter data about each property. Surveys were done during the summer of 2012. The survey crew drove by each property where the team checked the database for accuracy using visual observation and the aerial photographs on the survey maps. A Professional Agrologist verified what was on the site and a GIS technician altered the codes in the database as necessary (Figure 3). Corrections were handwritten on the maps. The map sheets were then brought back to the office to have the hand-drawn lines digitized into the GIS system and have the additional polygons entered into the database.

Once acquired through the survey, the land use data was brought into the GIS to facilitate analysis and produce maps. Digital data, in the form of a database and GIS shape files (for maps), is available upon request through a data sharing agreement with the Ministry of Agriculture.




Figure 3 Land Use Survey

Figure 4 provides an example of a map sheet from the RDN. The region was divided into 107 map sheets. Each map sheet also had a key map to indicate where it was located in the region.




Figure 4 GIS Map Sheet

The smallest unit for which water use is calculated are the polygons within each cadastre. A polygon is determined by a change in land use or irrigation system within a cadastre. Polygons are designated as blue lines within each cadastre as shown in Figures 4 and 5. The dataset for RDN encompasses 5,632 inventoried land parcels that are in or partially in the ALR. There are a total of 17,060 polygons generated within these land parcels. Figure 5 provides an enhanced view of a cadastre containing three polygons. Each cadastre has a unique identifier as does each polygon. The polygon identifier is acknowledged by PolygonID. This allows the survey team to call up the cadastre in the database, review the number of polygons within the cadastre and ensure the land use is coded accurately for each polygon.



Figure 5 Cadastre with Polygon

Soil Information

Soil information was obtained digitally from the Ministry of Environment’s Terrain and Soils Information System. The Computer Assisted Planning and Map Production application (CAPAMP) provided detailed (1:20,000 scale) soil surveys that were conducted in the Lower Mainland, on Southeast Vancouver Island, and in the Okanagan-Similkameen areas during the early 1980s. Products developed include soil survey reports, maps, agriculture capability and other related themes. Soil information required for this project was the soil texture (loam, etc.), the available water storage capacity and the peak infiltration rate for each texture type.

The intersection of soil boundaries with the cadastre and land use polygons creates additional polygons that the Model uses to calculate water demand. Figure 6 shows how the land use information is divided into additional polygons using the soil boundaries. The Model calculates water demand using every different combination of crop, soil and irrigation system as identified by each polygon.




Figure 6 GIS Model Graphic

Climate Information

The agricultural water demand is calculated using climate, crop, irrigation system and soil information data. To incorporate the climatic diversity, climate layers were developed for the entire region on a 500 m x 500 m grid. Each grid cell contains daily climate data, minimum and maximum temperature (T_{\min} and T_{\max}), and precipitation which allows the Model to calculate a daily reference evapotranspiration rate (ET_0) value. A range of agro-climatic indices such as growing degree days (GDD), corn heat units (CHU), frost free days and temperature sum (Tsum) can also be calculated for each grid cell based on temperature data. These values are used to determine seeding dates and the length of the growing season in the Model.

The climate dataset has been developed by using existing data from climate stations in and around RDN from 1961 to 2003. This climate data set was then interpolated to provide a climate data layer for the entire watershed on the 500 m x 500 m grid. A detailed description of the Model can be obtained by contacting the authors.

Some of the existing climate stations that were used to determine the climate coverage are shown in Figure 7. The attributes attached to each climate grid cell include:

- Latitude
- Longitude
- Elevation
- Aspect
- Slope
- Daily Precipitation
- Daily T_{\max} and T_{\min}




Figure 7 Nanaimo Area Climate Stations

The climate database generated contains T_{\min} , T_{\max} , T_{mean} and Precipitation for each day of the year from 1961 to 2003. The parameters that need to be selected, calculated and stored within the Model are evapotranspiration (ET_0), Tsum of 1,000 (for the Island), effective precipitation (EP), frost free days, GDD with base temperatures of 5 °C and 10 °C, CHU, and first frost date. These climate and crop parameters are used to determine the growing season length as well as the beginning and end of the growing season in Julian day.

Model Calculations

The Model calculates the water demand for each polygon by using crop, irrigation, soil and climate parameters as explained below. Each polygon has been assigned an ID number as mentioned previously.

It should be noted that in coastal regions like the Regional District of Nanaimo, many low-lying areas have high water tables which will reduce the overall irrigation demand. Agricultural water demand results from the Model will therefore be higher than what may actually be used as water tables have not been taken into the equation.

Crop

The CropID is an attribute of the PolygonID as each polygon will contain a single crop. The crop information (observed during the land use survey) has been collected and stored with PolygonID as part of the land use survey. CropID will provide cropping attributes to the Model for calculating water use for each polygon. CropID along with the climate data will also be used to calculate the growing season length and the beginning and end of the growing season. The attributes for CropID include rooting depth, availability coefficient, crop coefficient and a drip factor.

Rooting depth is the rooting depth for a mature crop in a deep soil.

An availability coefficient is assigned to each crop. The availability coefficient is used with the IrrigID to determine the soil moisture available to the crop for each PolygonID.

The crop coefficient adjusts the calculated ET_0 for the stages of crop growth during the growing season. Crop coefficient curves have been developed for every crop. The crop coefficient curve allows the Model to calculate water demand with an adjusted daily ET_0 value throughout the growing season.

The drip factor is used in the water use calculation for polygons where drip irrigation systems are used. Since the Model calculates water use by area, the drip factor adjusts the percentage of area irrigated by the drip system for that crop.

Irrigation

The IrrigID is an attribute of the PolygonID as each polygon will have a single irrigation system type operating. The irrigation information has been collected and stored (as observed during the land use survey) with the land use data. The land use survey determined if a polygon had an irrigation system operating, what the system type was, and if the system was being used. The IrrigID has an irrigation efficiency listed as an attribute.

Two of the IrrigID's, Overtreedrip and Overtreemicro are polygons that have two systems in place. Two irrigation ID's occur when an overhead irrigation system has been retained to provide crop cooling or frost protection. In this case, the efficiencies used in the Model are the drip and microsprinkler efficiencies.

Soil

The soil layer came from CAPAMP at the Ministry of Environment. In addition, soil data provided by Agriculture and Agri-Food Canada (AAFC) was also used to generate multiple soil layers within each polygon. Each parcel was assigned the most predominant soil polygon, and then for each crop field within that soil polygon, the most predominant texture within the crop's rooting depth was determined and assigned to the crop field.

Note that textures could repeat at different depths – the combined total of the thicknesses determined the most predominant texture. For example, a layer of 20 cm sand, followed by 40 cm clay and then 30 cm of sand would have sand be designated at the predominant soil texture.

The attributes attached to the SoilID is the Available Water Storage Capacity (AWSC) which is calculated using the soil texture and crop rooting depth.

The Maximum Soil Water Deficit (MSWD) is calculated to determine the parameters for the algorithm that is used to determine the Irrigation Requirement (IR). The Soil Moisture Deficit at the beginning of the season is calculated using the same terms as the MSWD.

Climate

The climate data in the Model is used to calculate a daily reference evapotranspiration rate (ET_o) for each climate grid cell. The data that is required to calculate this value are:

- Elevation, metres (m)
- Latitude, degrees ($^{\circ}$)
- Minimum Temperature, degree Celsius ($^{\circ}C$)
- Maximum Temperature, degree Celsius ($^{\circ}C$)
- Classification as Coastal or Interior
- Classification as Arid or Humid
- Julian Day

Data that is assumed or are constants in this calculation are:

- Wind speed 2 m/s
- Albedo or canopy reflection coefficient, 0.23
- Solar constant, G_{sc} 0.082 MJ 2 min $^{-1}$
- Interior and Coastal coefficients, K_{Rs} 0.16 for interior locations
0.19 for coastal locations
- Humid and arid region coefficients, K_o 0 $^{\circ}C$ for humid/sub-humid climates
2 $^{\circ}C$ for arid/semi-arid climates

Agricultural Water Demand Equation

The Model calculates the Agriculture Water Demand (AWD) for each polygon, as a unique crop, irrigation system, soil and climate data is recorded on a polygon basis. The polygons are then summed to determine the AWD for each cadastre. The cadastre water demand values are then summed to determine AWD for the basin, sub-basin, water purveyor or local government. The following steps provide the process used by the Model to calculate Agricultural Water Demand. Detailed information is available on request.

1. *Pre-Season Soil Moisture Content*

Prior to the start of each crop's growing season, the soil's stored moisture content is modelled using the soil and crop evaporation and transpiration characteristics and the daily precipitation values. Precipitation increases the soil moisture content and evaporation (modelled using the reference potential evapotranspiration) depletes it. In general, during the pre-season, the soil moisture depth cannot be reduced beyond the maximum evaporation depth; grass crops in wet climates, however, can also remove moisture through crop transpiration.

The process used to model the pre-season soil moisture content is:

1. Determine whether the modelling area is considered to be in a *wet* or *dry* climate (see *Wet/Dry Climate Assessment*), and retrieve the early season evaporation factor in the modelling area
2. For each crop type, determine the start of the growing season (see *Growing Season Boundaries*)
3. For each crop and soil combination, determine the *maximum soil water deficit* (MSWD) and *maximum evaporation factor* (maxEvaporation)
4. Start the initial storedMoisture depth on January 1 at the MSWD level
5. For each day between the beginning of the calendar year and the crop's growing season start, calculate a new stored moisture from:
 - a. the potential evapotranspiration (ET_o)
 - b. the early season evaporation factor (earlyEvaporationFactor)
 - c. the effective precipitation (EP) = actual precipitation \times earlyEvaporationFactor
 - d. daily Climate Moisture Deficit (CMD) = $ET_o - EP$
 - e. storedMoisture = previous day's storedMoisture – CMD

A negative daily CMD (precipitation in excess of the day's potential evapotranspiration) adds to the stored moisture level while a positive climate moisture deficit reduces the amount in the stored moisture reservoir. The stored moisture cannot exceed the maximum soil moisture deficit; any precipitation that would take the stored moisture level above the MSWD gets ignored.

For all crops and conditions except for grass in wet climates, the stored moisture content cannot drop below the maximum soil water deficit minus the maximum evaporation depth; without any crop transpiration in play, only a certain amount of water can be removed from the soil through evaporative processes alone. Grass in wet climates does grow and remove moisture from the soil prior to the start of the irrigation season, however. In those cases, the stored moisture level can drop beyond the maximum evaporation depth, theoretically to 0.

Greenhouses and mushroom barns have no stored soil moisture content.

2. *In-Season Precipitation*

During the growing season, the amount of precipitation considered effective (EP) depends on the overall wetness of the modelling area's climate (see *Wet/Dry Climate Assessment*). In dry climates, the first 5 mm of precipitation is ignored, and the EP is calculated as 75% of remainder:

$$EP = (\text{Precip} - 5) \times 0.75$$

In wet climates, the first 5 mm is included in the EP. The EP is 75% of the actual precipitation:

$$EP = \text{Precip} \times 0.75$$

Greenhouses and mushroom barns automatically have an EP value of 0.

3. ***Crop Cover Coefficient (K_c)***

As the crops grow, the amount of water they lose due to transpiration changes. Each crop has a pair of polynomial equations that provide the crop coefficient for any day during the crop's growing season. It was found that two curves, one for modelling time periods up to the present and one for extending the modelling into the future, provided a better sequence of crop coefficients than using a single curve for all years (currently 1961 to 2100). The application automatically selects the current or future curve as modelling moves across the *crop Curve Changeover Year*.

For alfalfa crops, there are different sets of equations corresponding to different cuttings throughout the growing season.

4. ***Crop Evapotranspiration (ET_c)***

The evapotranspiration for each crop is calculated as the general ET_o multiplied by the crop coefficient (K_c):

$$ET_c = ET_o \times K_c$$

5. ***Climate Moisture Deficit (CMD)***

During the growing season, the daily Climate Moisture Deficit (CMD) is calculated as the crop evapotranspiration (ET_c) less the Effective Precipitation (EP):

$$CMD = ET_c - EP$$

During each crop's growing season, a stored moisture reservoir methodology is used that is similar to the soil moisture content calculation in the pre-season. On a daily basis, the stored moisture level is used towards satisfying the climate moisture deficit to produce an *adjusted Climate Moisture Deficit* (CMD_a):

$$CMD_a = CMD - \text{storedMoisture}$$

If the storedMoisture level exceeds the day's CMD, then the CMD_a is 0 and the stored moisture level is reduced by the CMD amount. If the CMD is greater than the stored moisture, then all of the stored moisture is used (storedMoisture is set to 0) and the adjusted CMD creates an irrigation requirement.

The upper limit for the storedMoisture level during the growing season is the maximum soil water deficit (MSWD) setting.

6. Crop Water Requirement (CWR)

The Crop Water Requirement is calculated as the adjusted Climate Moisture Deficit (CMD_a) multiplied by the soil water factor (swFactor) and any stress factor (used primarily for grass crops):

$$CWR = CMD_a \times swFactor \times stressFactor$$

7. Irrigation Requirement (IR)

The Irrigation Requirement is the Crop Water Requirement (CWR) after taking into account the irrigation efficiency (I_e) and, for drip systems, the drip factor (D_f):

$$IR = CWR \times \frac{D_f}{I_e}$$

For irrigation systems other than drip, the drip factor is 1.

8. Irrigation Water Demand (IWD_{perc} and IWD)

The portion of the Irrigation Water Demand lost to deep percolation is the Irrigation Requirement (IR) multiplied by the percolation factor (soilPercFactor):

$$IWD_{perc} = IR \times soilPercFactor$$

The final Irrigation Water Demand (IWD) is then the Irrigation Requirement (IR) plus the loss to percolation (IWD_{perc}):

$$IWD = IR + IWD_{perc}$$

9. Frost Protection

For some crops (e.g. cranberries), an application of water is often used under certain climatic conditions to provide protection against frost damage. For cranberries, the rule is: when the temperature drops to 0 °C or below between March 16 and May 20 or between October 1 and November 15, a frost event will be calculated. The calculated value is an application of 2.5 mm per hour for 10 hours. In addition, 60% of the water is recirculated and reused, accounting for evaporation and seepage losses.

This amounts to a modelled water demand of 10 mm over the cranberry crop's area for each day that a frost event occurs between the specified dates.

10. Annual Soil Moisture Deficit

Prior to each crop's growing season, the Model calculates the soil's moisture content by starting it at full (maximum soil water deficit level) on January 1, and adjusting it daily according to

precipitation and evaporation. During the growing season, simple evaporation is replaced by the crop's evapotranspiration as it progresses through its growth stages. At the completion of each crop's growing season, an annual soil moisture deficit (SMD) is calculated as the difference between the soil moisture content at that point and the maximum soil water deficit (MSWD):

$$\text{SMD} = \text{MSWD} - \text{storedMoisture}$$

In dry/cold climates, this amount represents water that the farmer would add to the soil in order to prevent it from freezing. Wet climates are assumed to have sufficient precipitation and warm enough temperatures to avoid the risk of freezing without this extra application of water; the SMD demand is therefore recorded only for dry areas.

There is no fixed date associated with irrigation to compensate for the annual soil moisture deficit. The farmer may choose to do it any time after the end of the growing season and before the freeze up. In the Model's summary reports, the water demand associated with the annual soil moisture deficit shows as occurring at time 0 (week 0, month 0, etc.) simply to differentiate it from other demands that do have a date of occurrence during the crop's growing season.

Greenhouses and mushroom barns do not have an annual soil moisture deficit.

11. *Flood Harvesting*

Cranberry crops are generally harvested using flood techniques. The Model calculates the flood harvesting demand as 250 mm of depth for 10% of the cranberry farmed area. For modelling purposes, it is assumed that 250 mm of water gets applied to the total cranberry crop area, 10% at a time. The water is reused for subsequent portions, but by the time the entire crop is harvested, all of the water is assumed to have been used and either depleted through losses or released from the farm.

The water demand is therefore calculated as a fixed 25 mm over the entire cranberry crop area. The harvesting generally takes place between mid-October and mid-November where the Model treats it as occurring on the fixed date of November 16.

Livestock Water Use

The Model calculates an estimated livestock water demand using agricultural census data and an estimate of the water use per animal. Water use for each animal type is calculated a bit differently depending on requirements. For example, for a dairy milking cow, the water demand for each animal includes, drinking, preparation for milking, pen and barn cleaning, milking system washout, bulk tank washout and milking parlor washing. However, for a dry dairy cow, the demand only includes drinking and pen and barn cleaning.

The water use is estimated on a daily basis per animal even though the facility is not cleaned daily. For example, for a broiler operation, the water use for cleaning a barn is calculated as 4 hours of pressure washing per cycle at a 10 gpm flow rate, multiplied by 6 cycles per barn with each barn holding 50,000 birds. On a daily basis, this is quite small with a value of 0.01 litres per day per bird applied.

For all cases, the daily livestock demand is applied to the farm location. However, in the case of beef, the livestock spend quite a bit of the year on the range. Since the actual location of the animals cannot be ascertained, the water demand is applied to the home farm location, even though most of the demand will not be from this location. Therefore, the animal water demand on a watershed scale will work fine but not when the demand is segregated into sub-watersheds or groundwater areas.

The estimates used for each livestock are shown in Table 1.

Table 1 Livestock Water Demand (Litres/day)				
Animal Type	Drinking	Milking Preparation	Barn Component	Total
Milking Dairy Cow	65	5	15	85
Dry Cow	45		5	50
Swine	12		0.5	12.5
Poultry – Broiler	0.16		0.01	0.17
Poultry – Layer	0.08		0.01	0.09
Turkeys	0.35		0.01	0.36
Goats	8			8
Sheep	8			8
Beef – range, steer, bull, heifer	50			50
Horses	50			50

Definition and Calculation of Individual Terms used in the Irrigation Water Demand Equation

Growing Season Boundaries

There are three sets of considerations used in calculating the start and end of the irrigation season for each crop:

- temperature-based growing season derivations, generally using Temperature Sum (Tsum) or Growing Degree Day (GDD) accumulations
- the growing season overrides table
- the irrigation season overrides table

These form an order of precedence with later considerations potentially overriding the dates established for the previous rules. For example, the temperature-based rules might yield a growing season start date of day 90 for a given crop in a mild year. To avoid unrealistic irrigation starts, the season overrides table might enforce a minimum start day of 100 for that crop; at that point, the season start would be set to day 100. At the same time, a Water Purveyor might not turn on the water supply until day 105; specifying that as the minimum start day in the irrigation season overrides table would prevent any irrigation water demands until day 105.

This section describes the rules used to establish growing season boundaries based on the internal calculations of the Model. The GDD and Tsum Day calculations are described in separate sections. The *standard end of season* specified for several crops is the earlier of the end date of Growing Degree Day with base temperature of 5 °C (GDD₅) or the first frost.

1. *Corn (silage corn)*

- uses the corn_start date for the season start
- season end: earlier of the killing frost or the day that the CHU2700 (2700 Corn Heat Units) threshold is reached

2. *Sweetcorn, Potato, Tomato, Pepper, Strawberry, Vegetable, Pea*

- corn_start date for the season start
- corn start plus 110 days for the season end

3. *Cereal*

- GDD5 start for the season start
- GDD5 start plus 130 days for the season end

4. *AppleHD, AppleMD, AppleLD, Asparagus, Berry, Blueberry, Ginseng, Nuts, Raspberry, Sourcherry, Treefruit, Vineberry*

- season start: $(0.8447 \times \text{tsum600_day}) + 18.877$
- standard end of season

5. *Pumpkin*

- corn_start date
- standard end of season

6. ***Apricot***
 - season start: $(0.9153 \times \text{tsum400_day}) + 5.5809$
 - standard end of season
7. ***CherryHD, CherryMD, CherryLD***
 - season start: $(0.7992 \times \text{tsum450_day}) + 24.878$
 - standard end of season
8. ***Grape, Kiwi***
 - season start: $(0.7992 \times \text{tsum450_day}) + 24.878$
 - standard end of season
9. ***Peach, Nectarine***
 - season start: $(0.8438 \times \text{tsum450_day}) + 19.68$
 - standard end of season
10. ***Plum***
 - season start: $(0.7982 \times \text{tsum500_day}) + 25.417$
 - standard end of season
11. ***Pear***
 - season start: $(0.8249 \times \text{tsum600_day}) + 17.14$
 - standard end of season
12. ***Golf, TurfFarm***
 - season start: later of the GDD₅ start and the tsum300_day
 - standard end of season
13. ***Domestic, Yard, TurfPark***
 - season start: later of the GDD₅ start and the tsum400_day
 - standard end of season
14. ***Greenhouse (interior greenhouses)***
 - fixed season of April 1 – October 30
15. ***GH Tomato, GH Pepper, GH Cucumber***
 - fixed season of January 15 – November 30
16. ***GH Flower***
 - fixed season of March 1 – October 30
17. ***GH Nursery***
 - fixed season of April 1 – October 30
18. ***Mushroom***
 - all year: January 1 – December 31

19. Shrubs/Trees, Fstock, NurseryPOT

- season start: tsum500_day
- end: julian day 275

20. Floriculture

- season start: tsum500_day
- end: julian day 225

21. Cranberry

- season start: tsum500_day
- end: julian day 275

22. Grass, Forage, Alfalfa, Pasture

- season start: later of the GDD₅ and the tsum600_day
- standard end of season

23. Nursery

- season start: tsum400_day
- standard end of season

Evapotranspiration (ET_o)

The ET_o calculation follows the FAO Penman-Montieth equation. Two modifications were made to the equation:

- Step 6 – Inverse Relative Distance Earth-Sun (d_r)
Instead of a fixed 365 days as a divisor, the actual number of days for each year (365 or 366) was used.
- Step 19 – Evapotranspiration (ET_o)
For consistency, a temperature conversion factor of 273.16 was used instead of the rounded 273 listed.

Availability Coefficient (AC)

The availability coefficient is a factor representing the percentage of the soil's total water storage that the crop can readily extract. The factor is taken directly from the crop factors table (*crop_factors*) based on the cropId value.

Rooting Depth (RD)

The rooting depth represents the crop's maximum rooting depth and thus the depth of soil over which the plant interacts with the soil in terms of moisture extraction. The value is read directly from the crop factors table.

Stress Factor (stressFactor)

Some crops, such as *grasses*, are often irrigated to a less degree than their full theoretical requirement for optimal growth. The *stress factor (crop_groups_and_factors)* reduces the calculated demand for these crops.

Available Water Storage Capacity (AWSC)

The available water storage capacity is a factor representing the amount of water that a particular soil texture can hold without the water dropping through and being lost to deep percolation. The factor is taken directly from the soil factors table (*soil_factors*).

Maximum Soil Water Deficit (MSWD)

The maximum soil water deficit is the product of the crop's availability coefficient, rooting depth, and the available water storage capacity of the soil:

$$MSWD = RD \times AWSC \times AC$$

Deep Percolation Factor (soilPercFactor)

The soil percolation factor is used to calculate the amount of water lost to deep percolation under different management practices.

For greenhouse crops, the *greenhouse leaching factor* is used as the basic soil percolation factor. This is then multiplied by a greenhouse recirculation factor, if present, to reflect the percentage of water re-captured and re-used in greenhouse operations.

$$soilPercFactor = soilPercFactor \times (1 - recirculationFactor)$$

For Nursery Pot (Nursery POT) and Forestry Stock (Fstock) crops, the soil percolation factor is fixed at 35%. For other crops, the factor depends on the soil texture, the MSWD, the irrigation system, and the Irrigation Management Practices code. The percolation factors table (*soil_percolation_factors*) is read to find the first row with the correct management practices, soil texture and irrigation system, and a MSWD value that matches or exceeds the value calculated for the current land use polygon.

If the calculated MSWD value is greater than the index value for all rows in the percolation factors table, then the highest MSWD factor is used. If there is no match based on the passed parameters, then a default value of 0.25 is applied.

For example, a calculated MSWD value of 82.5 mm, a soil texture of sandy loam (SL) and an irrigation system of solid set overtree (Ssovertree) would retrieve the percolation factor associated with the MSWD index value of 75 mm in the current table (presently, there are rows for MSWD 50 mm and 75 mm for SL and Ssovertree).

Maximum Evaporation Factor (maxEvaporation)

Just as different soil textures can hold different amounts of water, they also have different depths that can be affected by evaporation. The factor is taken directly from the soil factors table.

Irrigation Efficiency (I_e)

Each irrigation system type has an associated efficiency factor (inefficient systems require the application of more water in order to satisfy the same crop water demand). The factor is read directly from the irrigation factors table (*irrigation_factors*).

Soil Water Factor (swFactor)

For the greenhouse “crop”, the soil water factor is set to 1. For other crops, it is interpolated from a table (*soil_water_factors*) based on the MSWD. For Nurseries, the highest soil water factor (lowest MSWD index) in the table is used; otherwise, the two rows whose MSWD values bound the calculated MSWD are located and a soil water factor interpolated according to where the passed MSDW value lies between those bounds.

For example, using the current table with rows giving soil water factors of 0.95 and 0.9 for MSWD index values of 75 mm and 100 mm respectively, a calculated MSWD value of 82.5 mm would return a soil water factor of:

$$0.95 + \left[\frac{82.5 - 75}{100 - 75} \times (0.9 - 0.95) \right]$$
$$= 0.935$$

If the calculated MSWD value is higher or lower than the index values for all of the rows in the table, then the factor associated with the highest or lowest MSWD index is used.

Early Season Evaporation Factor (earlyEvaporationFactor)

The effective precipitation (precipitation that adds to the stored soil moisture content) can be different in the cooler pre-season than in the growing season. The early season evaporation factor is used to determine what percentage of the precipitation is considered effective prior to the growing season.

Crop Coefficient (K_c)

The crop coefficient is calculated from a set of fourth degree polynomial equations representing the crop’s ground coverage throughout its growing season. The coefficients for each term are read from the crop factors table based on the crop type, with the variable equalling the number of days since the start of the crop’s growing season. For example, the crop coefficient for Grape on day 35 of the growing season would be calculated as:

$$K_c = [0.0000000031 \times (35)^4] + [-0.0000013775 \times (35)^3] + (0.0001634536 \times (35)^2) + (-0.0011179845 \times 35) + 0.2399004137$$
$$= 0.346593241$$

Alfalfa crops have an additional consideration. More than one cutting of alfalfa can be harvested over the course of the growing season, and the terms used for the crop coefficient equation changes for the different cuttings. For alfalfa, the alfalfa cuttings table is first used to determine which cutting period the day belongs to (first, intermediate or last), and after that the associated record in the crop factors table is accessed to determine the terms.

There are two sets of polynomial coefficients used to calculate the crop coefficient; the first set is used for modelling time periods up to the year specified as the *crop curve changeover year*; and the second for modelling into the future. The changeover year will be modified as time goes on and new historical climate observations become available.

Growing Degree Days (GDD)

The Growing Degree Day calculations generate the start and end of GDD accumulation.

1. *Start of GDD Accumulation*

For each base temperature (bases 5 and 10 are always calculated, other base temperature can be derived), the start of the accumulation is defined as occurring after 5 consecutive days of T_{mean} matching or exceeding the base temperature (BaseT). The search for the start day gets reset if a killing frost (< -2 °C) occurs, even after the accumulation has started. The search also restarts if there are 2 or more consecutive days of $T_{\text{min}} \leq 0$ °C. The GDD start is limited to Julian days 1 to 210; if the accumulation has not started by that point, then it is unlikely to produce a reasonable starting point for any crop.

2. *End of GDD accumulation*

The search for the end of the GDD accumulation begins 50 days after its start. The accumulation ends on the earlier of 5 consecutive days where T_{mean} fails to reach BaseT (strictly *less than*) or the first killing frost (-2 °C).

During the GDD accumulation period, the daily contribution is the difference between T_{mean} and BaseT, as long as T_{mean} is not less than BaseT:

$$\text{GDD} = T_{\text{mean}} - \text{BaseT}; 0 \text{ if negative}$$

Frost Indices

Three frost indices are tracked for each year:

- the last spring frost is the latest day in the first 180 days of the year with a $T_{\text{min}} \leq 0$ °C
- the first fall frost is the first day between days 240 and the end of the year where $T_{\text{min}} \leq 0$ °C
- the killing frost is the first day on or after the first fall frost where $T_{\text{min}} \leq -2$ °C

Corn Heat Units (CHU)

The Corn Heat Unit is the average of two terms using T_{min} and T_{max} . Prior to averaging, each term is set to 0 individually if it is negative.

$$\begin{aligned} \text{term1} &= [3.33 \times (T_{\max} - 10)] - [0.084 \times (T_{\max} - 10) \times (T_{\max} - 10)]; 0 \text{ if negative} \\ \text{term2} &= 1.8 \times (T_{\min} - 4.44); 0 \text{ if negative} \\ \text{CHU} &= \frac{(\text{term1} + \text{term2})}{2} \end{aligned}$$

Corn Season Start and End

The corn season boundary derivations are similar to the GDD determinations. The start day is established by 3 consecutive days where $T_{\text{mean}} \geq 11.2$ °C. As in the case of the GDD calculations, the search for the corn season start day gets reset if $T_{\min} \leq -2$ °C, or if there are 2 or more consecutive days of -2 °C $\leq T_{\min} \leq 0$ °C.

The search for the silage corn season end begins 50 days after the start. The season ends on the earlier of a mean temperature dropping below 10.1 or a killing frost.

The end of the sweet corn season is defined as 110 days after the season start.

Tsum Indices

The Tsum day for a given number is defined as the day that the sum of the positive daily T_{mean} reaches that number. For example, the Tsum400 day is the day where the sum of the positive T_{mean} starting on January 1 sum to 400 units or greater.

Days where T_{mean} falls below 0 °C are simply not counted; therefore, the Model does not restart the accumulation sequence.

Wet/Dry Climate Assessment

Starting with the Lower Mainland, some of the modelling calculations depend on an assessment of the general climatic environment as *wet* or *dry*. For example, when modelling the soil moisture content prior to the start of the crop's growing season, the reservoir can only be drawn down by evaporation except for *grass* crops in *wet* climates which can pull additional moisture out of the soil.

The assessment of wet or dry uses the total precipitation between May 1 and September 30. If the total is more than 125 mm during that period, the climate is considered to be *wet* and otherwise *dry*.

Groundwater Use

The Model generates water sources for irrigation systems. This is done by first determining which farms are supplied by a water purveyor, and then coding those farms as such. Most water purveyors use surface water but where groundwater is used, the farms are coded as groundwater use. The second step is to check all water licences and assign the water licences to properties in the database. The remaining farms that are irrigating will therefore not have a water licence or be supplied by a water purveyor. The assumption is made that these farms are irrigated by groundwater sources.

Land Use Results

A summary of the land area and the inventoried area of the Regional District of Nanaimo is shown in Table 2. The inventoried area includes parcels that are in and partially in the Agricultural Land Reserve (ALR). The primary agricultural use of the ARL area is shown in Table 3 where only 1,366 parcels currently have active agriculture. Refer to the [Agricultural Land Use Inventory](#) reports for details.

The Model also reports out on groundwater aquifers. Figure 8 provides a schematic of the higher yielding aquifer areas in RDN based on the information from B.C. Ministry of Environment.

Table 2 Overview of RDN's Land and Inventoried Area		
Area Type	Area (ha)	Number of Parcels
RDN		
Total Area	319,881	-
Area of Water Feature	116,310	-
Area of Land (excluding water features)	203,571	-
ALR Area	18,062	4,006
Area of First Nations Reserve	406	262
Inventoried Area		
Total Inventoried Area	38,976	5,632
Area of First Nations Reserve in ALR	124	92

Table 3 Summary of Primary Agricultural Activities within the ALR where Primary Land Use is Agriculture in RDN		
Primary Agriculture Activity	Total Land Cover (ha)	Number of Parcels
Glass and poly greenhouse	5	30
Tree fruits	14	25
Grapes	4	7
Vines and Berries	84	15
Forage and Pasture	4,093	1,205
Vegetables	64	39
Floriculture	<1	1
Turf, Nut Trees, Specialty	23	8
Nursery	91	27
Cultivated land, Fallow land	18	9
Total	4,395	1,366




Figure 8 Higher Productive Groundwater Aquifers in RDN

Agricultural Water Demand Model Results

The Model has a reporting feature that can save and generate reports for many different scenarios that have been pre-developed. This report will provide a summary of the reported data in the Appendices. Climate data from 1997 and 2003 were chosen as they represent a relatively wet year and dry year respectively. Most reports are based on the 2003 data since the maximum current demand can then be presented.

Annual Crop Water Demand – Tables A and B

The Model can use three different irrigation management factors, good, average and poor. Unless otherwise noted, average management were used in the tables. Table A provides the annual irrigation water demand for current crop and irrigation systems for the year 2003 using average irrigation management, and Table B provides the same data for 1997.

The outdoor irrigated acreage in the ALR for RDN is 1,018 hectares (ha). The total annual irrigation demand for this area was 6,917,243 m³ in 2003 (a dry year), and dropped to 3,106,927 m³ in 1997 (a wet year).

Of interest is that during a wet year like 1997, the demand was only 45% of a hot dry year like 2003. Another point to consider is that the actual water demand supplied by an irrigation system may be less than the numbers shown above. The reason is that the Model does not have an adjustment for water supplied to the crop by high water tables. In coastal regions, agriculture is often located in areas which have predominantly high water tables due to the climate. The high water tables will reduce irrigation demand that is not accounted for in the model outputs. The numbers should therefore be considered the highest estimate demand.

In addition, the Model also calculates demand based on relatively good practices. As such, actual use may actually be higher or lower than what is calculated by the Model.

The predominant irrigated agriculture crop in RDN is forage that includes forage corn, grass, legume and pasture.

Annual Water Demand Reported by Irrigation System – Table C

The crop irrigation demand can also be reported by irrigation system type as shown in Tables C. The total area that is currently irrigated by efficient systems such as drip, microsprinkler or microspray is relatively small as forage is the predominant crop type. Sprinkler and travelling gun systems used on forage and pasture crops account for 80% of the irrigation system types.

Annual Water Demand by Soil Texture – Table D

Table D provides the annual water demand by soil texture. Where soil texture data is missing, the soil texture has been defaulted to sandy loam. The defaults are shown in Table D.

Annual Water Demand by Aquifer – Table E

Table E provides information on the irrigation demand taken from the aquifers within RDN boundaries. The table also shows surface water demands for systems that are licensed to take water from surface sources within the aquifer boundaries. Some properties are located outside of known aquifers; therefore, were listed under “others”. The total groundwater extracted is estimated to be 3,441,185 m³.

Irrigated Area by Local Government – Table F

Table F provides a breakdown of the agricultural irrigated areas within the boundaries of each local government within the RDN.

Irrigation Management Factors – Table G

The Model can estimate water demand based on poor, average and good irrigation management factors. This is accomplished by developing an irrigation management factor for each crop, soil and irrigation system combination based on subjective decision and percolation rates. The Maximum Soil Water Deficit (MSWD) is the maximum amount of water that can be stored in the soil within the crop rooting zone. An irrigation system applying more water than what can be stored will result in percolation beyond the crop’s rooting depth. Irrigation systems with high application rates will have a probability of higher percolation rates, a stationary gun for instance.

For each soil class, a range of four MSWD are provided, which reflect a range of crop rooting depths. An irrigation management factor, which determines the amount of leaching, is established for each of the MSWD values for the soil types (Table 4). The management factor is based on irrigation expertise as to how the various irrigation systems are able to operate. For example, Table 4 indicates that for a loam soil and a MSWD of 38 mm, a solid set overtree system has a management factor of 0.1 for good management while the drip system has a management factor of 0.05. This indicates that it is easier to prevent percolation with a drip system than it is with a solid set sprinkler system. For poor management, the factors are higher.

There are a total of 1,344 irrigation management factors established for the 16 different soil textures, MSWD and 21 different irrigation system combinations used in the Model.

Table 4 Irrigation Management Factors							
Soil Texture	MSWD	Solid Set Overtree			Drip		
		Good	Average	Poor	Good	Average	Poor
Loam	38	0.10	0.15	0.20	0.05	0.10	0.15
	50	0.05	0.10	0.15	0.05	0.075	0.10
	75	0.05	0.10	0.15	0.05	0.075	0.10
	100	0.05	0.075	0.10	0.05	0.075	0.10
Sandy loam	25	0.20	0.225	0.25	0.10	0.15	0.20
	38	0.10	0.15	0.20	0.10	0.125	0.15
	50	0.05	0.10	0.15	0.05	0.10	0.10
	75	0.05	0.10	0.15	0.05	0.075	0.10

The management factors increase as the MSWD decreases because there is less soil storage potential in the crop rooting depth. For irrigation systems such as guns, operating on a pasture which has a shallow rooting depth, on a sandy soil which cannot store much water, the poor irrigation management factor may be as high as 0.5.

The management factor used in the Model assumes all losses are deep percolation while it is likely that some losses will occur as runoff as well.

Table G provides an overview of the impacts on the management factor and irrigation systems used. An improvement of 4% in total water use reduction could be achieved by improved management to reduce percolation. The soils located in the region have a higher water holding capacity so improved management shows limited water use reduction.

Table G also provides percolation rates based on good, average and poor management using 2003 climate data. In summary, there is 628,243 m³ of water lost to percolation on good management, 759,281 m³ on average management, and 890,320 m³ on poor management. Percolation rates for poor management are 41% higher than for good management.

Deep Percolation – Table H

The percolation rates vary by crop, irrigation system type, soil and the management factor used. Table H shows the deep percolation amounts by irrigation system type for average management. The last column provides a good indication of the average percolation per hectare for the various irrigation system types. Landscape systems have a high percolation rate predominantly because application rates are high and the crop rooting depth is quite shallow. Microspray and microsprinklers are also shown to have high percolation rates but these systems are likely inside greenhouse nursery systems and the water may be recirculated

Improved Irrigation Efficiency and good Management – Table I

There is an opportunity to reduce water use by converting irrigation systems to a higher efficiency for some crops. For example, drip systems could be used for all berry crops, vegetable crops and some of the other horticultural crops, but not forage crops. In addition, using better management such as irrigation scheduling techniques will also reduce water use, especially for forage where drip conversion is not possible. Table I provides a scenario of water demand if all sprinkler systems are converted to drip systems for horticultural crops in RDN, using good irrigation management. The water demand for 2003 would reduce from 6,917,243 m³ to 6,458,897 m³ if sprinkler systems were converted to drip and good management practices were implemented. Since forage is such a predominant crop in the region, the amount of reduction achieved is only around 500,000 m³.

Water Demand for Frost Protection, Harvesting and Other – Table J

The algorithms to calculate water demand for frost protection, crop harvesting, greenhouse and potted nursery are different from the conventional irrigation system calculation. These uses are therefore reported separately from field irrigation use. For RDN, the total use is calculated to be 256,525 m³.

Livestock Water Use – Table K

The Model provides an estimate of water use for livestock. The estimate is based on the number of animals in RDN as determined by the latest census, the drinking water required for each animal per day and the barn or milking parlour wash water. Values used are shown in Table K. For RDN, the amount of livestock water is estimated at 86,356 m³.

Climate Change Water Demand for 2050 – Table L

The Model also has access to climate change information until the year 2100. While data can be run for each year, three driest years in the 2050’s were selected to give a representation of climate change. Figure 9 shows the climate data results which indicate that 2053, 2056, and 2059 generate the highest annual ET_o and lowest annual precipitation. These three years were used in this report.




Figure 9 Annual ET and Effective Precipitation in 2050's

Table L provides the results of climate change on irrigation demand for the three years selected using current crops and irrigation systems. Current crops and irrigation systems are used to show the increase due to climate change only, with no other changes taking place.

Figure 10 shows all of the climate change scenario runs for the Okanagan using 12 climate models from 1960 to 2100. This work was compiled by Denise Neilsen at the Agriculture and Agri-Food Canada – Summerland Research Station. There is a lot of scatter in this figure, but it is obvious that there is a trend of increasing water demand.

The three climate change models used in this report are RCP26, RCP45 and RCP85. Running only three climate change models on three selected future years in RDN is not sufficient to provide a trend like in Figure 10. What the results do show is that in an extreme climate scenario, it is possible to have an annual water demand that is 30% higher than what was experienced in 2003. Averaging the data between the three climate change models shows that if the data for just the year 2053 is examined, the increase in demand is 10% higher than 2003. More runs of the climate change models will be required to better estimate a climate change trend for RDN.




Figure 10 Future Irrigation Demand for All Outdoor Uses in the Okanagan in Response to Observed Climate Data (Actuals) and Future Climate Data Projected from a Range of Global Climate Models

Agricultural Buildout Crop Water Demand Using 2003 Climate Data – Table M

An agricultural buildout scenario was developed that looked at potential agricultural lands that could be irrigated in the future. The rules used to establish where potential additional agricultural lands were located in RDN are as follows:

- within 1,000 m of water supply (lake)
- within 1,000 m of water supply (water course)
- within 1,000 m of water supply (wetland)

- within 1,000 m of high productivity aquifer
- within 1,000 m of water purveyor
- with Ag Capability class 1-4 only where available
- must be within the ALR
- below 250 m average elevation

For the areas that are determined to be eligible for future buildout, a crop and irrigation system need to be applied. Where a crop already existed in the land use inventory, that crop would remain and an irrigation system assigned. If no crop existed, then a crop and irrigation system are assigned as per the criteria below.

- **Forage crops:** 60% of buildout area with sprinkler irrigation
- **Pasture:** 20% of buildout area with sprinkler irrigation
- **Blueberries:** 10% of buildout area with drip irrigation
- **Vegetables:** 10% of buildout area with drip irrigation

Figure 11 indicates the location of agricultural land that is currently irrigated (dark green) and the land that can be potentially irrigated (red). Based on the scenario provided for RDN, the additional agricultural land that could be irrigated is 3,111 ha, bringing the total irrigated area to 4,129 ha. The water demand for a year like 2003 would be 26,082,504 m³ assuming efficient irrigation systems and good management.




Figure 11 RDN Irrigation Expansion Potential

Agricultural Buildout Crop Water Demand for 2050 – Table N

The same irrigation expansion and cropping scenario used to generate the values in Table M were used to generate the climate change water demand shown in Table N. Three climate models were used and the results averaged. When climate change is added to the buildout scenario the water demand increases from 26 million m³ in 2003 to 29.5 million m³ if averaging the three climate change models for the 2053 scenario. Again, more runs are required to develop a good trend with the climate change data. See discussion under Table L.

Irrigation Systems Used for the Buildout Scenario – Table O

Table O provides an account of the irrigation systems used by area for the buildout scenario in the previous two examples. Note that sprinkler irrigation is still most predominant as forage is projected to be the major crop.

Water Demand by Aquifer for the Buildout Scenario – Table P

Table P provides the water demand based on aquifers for the buildout scenario in Table M. It can be compared with the values in Table E without buildout.

Water Demand by Local Government for the Buildout Scenario – Table Q

Table Q provides the water demand based on local governments for the buildout scenario in Table M. It can be compared with the values in Table F without buildout.

Literature

Cannon, A.J., and Whitfield, P.H. (2002), Synoptic map classification using recursive partitioning and principle component analysis. *Monthly Weather Rev.* 130:1187-1206.

Cannon, A.J. (2008), Probabilistic multi-site precipitation downscaling by an expanded Bernoulli-gamma density network. *Journal of Hydrometeorology*. <http://dx.doi.org/10.1175%2F2008JHM960.1>
Intergovernmental Panel on Climate Change (IPCC) (2008), Fourth Assessment Report –AR4. <http://www.ipcc.ch/ipccreports/ar4-syr.htm>

Neilsen, D., Duke, G., Taylor, W., Byrne, J.M., and Van der Gulik T.W. (2010). Development and Verification of Daily Gridded Climate Surfaces in the Okanagan Basin of British Columbia. *Canadian Water Resources Journal* 35(2), pp. 131-154. <http://www4.agr.gc.ca/abstract-resume/abstract-resume.htm?lang=eng&id=21183000000448>

Allen, R. G., Pereira, L. S., Raes, D. and Smith, M. (1998). Crop evapotranspiration Guidelines for computing crop water requirements. FAO Irrigation and Drainage Paper 56. United Nations Food and Agriculture Organization. Rome. 100pp

Appendix Tables

Appendix Table A	2003 Water Demand by Crop with Average Management
Appendix Table B	1997 Water Demand by Crop with Average Management
Appendix Table C	2003 Water Demand by Irrigation System with Average Management
Appendix Table D	2003 Water Demand by Soil Texture with Average Management
Appendix Table E	2003 Water Demand by Aquifer with Average Management
Appendix Table F	2003 Water Demand by Local Government with Average Management
Appendix Table G	2003 Management Comparison on Irrigation Demand and Percolation Volumes
Appendix Table H	2003 Percolation Volumes by Irrigation System with Average Management
Appendix Table I	2003 Crop Water Demand for Improved Irrigation System Efficiency and Good Management
Appendix Table J	2003 Water Demand for Frost Protection, Harvesting and Other Use with Average Management
Appendix Table K	2003 Water Demand by Animal Type with Average Management
Appendix Table L	Climate Change Water Demand Circa 2050 for a High Demand Year with Good Management using Current Crops and Irrigation Systems
Appendix Table M	Buildout Crop Water Demand for 2003 Climate Data and Good Management
Appendix Table N	Buildout Crop Water Demand for Climate Change Data Circa 2050 and Good Management
Appendix Table O	Buildout Irrigation System Demand for 2003 Climate Data and Good Management
Appendix Table P	Buildout Water Demand by Aquifer for 2003 Climate Data and Good Management
Appendix Table Q	Buildout Water Demand by Local Government for 2003 Climate Data and Good Management

Appendix Table A 2003 Water Demand by Crop with Average Management

Water Source	Surface Water			Reclaimed Water			Groundwater			Total		
	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)
Apple	0.2	1,427	779	-	-	-	5.6	32,767	590	5.7	34,194	596
Berry	-	-	-	-	-	-	9.3	48,870	526	9.3	48,870	526
Blueberry	1.5	7,222	467	-	-	-	59.6	242,556	407	61.1	249,778	409
Cranberry	-	-	-	-	-	-	20.2	300,423	1,490	20.2	300,423	1,490
Forage	462.2	3,019,008	653	-	-	-	242.3	1,753,321	724	704.4	4,772,329	677
Golf	16.1	131,788	817	-	-	-	37.5	298,572	795	53.7	430,360	802
Grape	-	-	-	-	-	-	4.8	10,571	222	4.8	10,571	222
Greenhouse	0.7	13,071	1,801	-	-	-	11.3	207,336	1,842	12.0	220,407	1,839
Nursery Floriculture	-	-	-	-	-	-	0.2	646	342	0.2	646	342
Nursery Shrubs/Trees	0.6	3,682	298	-	-	-	16.9	64,601	286	17.5	68,283	287
Pasture/Grass	11.7	82,788	705	-	-	-	7.5	50,531	675	19.2	133,319	693
Raspberry	0.2	1,127	579	-	-	-	7.7	37,043	478	7.9	38,170	481
Recreational Turf	-	-	-	-	-	-	6.8	47,479	698	6.8	47,479	698
Strawberry	-	-	-	-	-	-	3.3	13,918	422	3.3	13,918	422
Sweetcorn	-	-	-	-	-	-	29.0	113,340	391	29.0	113,340	391
Turf Farm	20.9	178,124	853	-	-	-	3.9	30,959	795	24.8	209,083	844
Vegetable	5.9	37,821	636	-	-	-	32.6	188,254	578	38.5	226,075	587
TOTALS	520.2	3,476,058	668	-	-	-	498.3	3,441,185	691	1,018.4	6,917,243	679

Appendix Table B 1997 Water Demand by Crop with Average Management

Water Source	Surface Water			Reclaimed Water			Groundwater			Total		
	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)
Apple	0.2	592	323	-	-	-	5.6	13,396	241	5.7	13,988	244
Berry	-	-	-	-	-	-	9.3	17,272	186	9.3	17,272	186
Blueberry	1.5	2,595	168	-	-	-	59.6	66,296	111	61.1	68,891	113
Cranberry	-	-	-	-	-	-	20.2	125,027	620	20.2	125,027	620
Forage	462.2	1,295,008	280	-	-	-	242.3	763,317	315	704.4	2,058,325	292
Golf	16.1	73,917	458	-	-	-	37.5	152,932	407	53.7	226,849	423
Grape	-	-	-	-	-	-	4.8	2,355	49	4.8	2,355	49
Greenhouse	0.7	12,260	1,690	-	-	-	11.3	192,960	1,714	12.0	205,221	1,712
Nursery Floriculture	-	-	-	-	-	-	0.2	288	152	0.2	288	152
Nursery Shrubs/Trees	0.6	1,827	116	-	-	-	16.9	28,312	99	17.5	30,139	100
Pasture/Grass	11.7	36,934	315	-	-	-	7.5	24,137	322	19.2	61,070	318
Raspberry	0.2	364	187	-	-	-	7.7	8,979	116	7.9	9,343	118
Recreational Turf	-	-	-	-	-	-	6.8	25,640	377	6.8	25,640	377
Strawberry	-	-	-	-	-	-	3.3	5,717	173	3.3	5,717	173
Sweetcorn	-	-	-	-	-	-	29.0	34,061	118	29.0	34,061	118
Turf Farm	20.9	92,139	441	-	-	-	3.9	15,574	400	24.8	107,713	435
Vegetable	5.9	20,626	347	-	-	-	32.6	94,401	290	38.5	115,027	298
TOTALS	520.2	1,536,262	295	-	-	-	498.3	1,570,664	315	1,018.4	3,106,927	305

Appendix Table C 2003 Water Demand by Irrigation System with Average Management

Water Source	Surface Water			Reclaimed Water			Groundwater			Total		
	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)
Drip	1.8	7,850	432	-	-	-	80.3	304,797	380	82.1	312,647	381
Flood	-	-	-	-	-	-	20.2	300,423	1,490	20.2	300,423	1,490
Golfsprinkler	4.1	27,800	676	-	-	-	34.6	274,234	792	38.7	302,035	780
Handline	35.3	278,731	790	-	-	-	20.2	119,516	592	55.5	398,247	718
Landscapesprinkler	12.0	103,988	865	-	-	-	9.7	71,816	738	21.8	175,804	808
Microsprinkler	0.7	13,071	1,801	-	-	-	11.3	207,336	1,842	12.0	220,407	1,839
Overtreedrip	-	-	-	-	-	-	7.3	31,721	434	7.3	31,721	434
SDI	-	-	-	-	-	-	4.0	17,605	440	4.0	17,605	440
Sprinkler	407.6	2,528,947	620	-	-	-	130.9	745,651	570	538.4	3,274,598	608
Ssovertime	-	-	-	-	-	-	12.9	80,578	627	12.9	80,578	627
Ssundertree	1.8	14,558	815	-	-	-	-	-	-	1.8	14,558	815
Travgun	56.8	501,113	882	-	-	-	116.2	865,943	745	173.0	1,367,056	790
Wheeline	-	-	-	-	-	-	50.8	421,566	830	50.8	421,566	830
TOTALS	520.2	3,476,058	668	-	-	-	498.3	3,441,185	691	1,018.4	6,917,243	679

Appendix Table D 2003 Water Demand by Soil Texture with Average Management

Water Source	Surface Water			Reclaimed Water			Groundwater			Total		
	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)
Clayey Sand	-	-	-	-	-	-	0.3	1,142	361	0.3	1,142	361
Cultured Medium	1.1	15,830	1,490	-	-	-	14.8	233,636	1,584	15.8	249,466	1,578
Fine Sandy Loam	26.7	212,385	795	-	-	-	3.9	30,874	794	30.6	243,260	795
Loam	0.1	955	805	-	-	-	11.6	71,109	614	11.7	72,063	616
Loamy Sand	37.4	317,893	850	-	-	-	79.3	631,932	797	116.7	949,825	814
Organic	38.4	159,535	415	-	-	-	70.8	320,250	452	109.2	479,785	439
Peat	-	-	-	-	-	-	20.2	300,423	1,490	20.2	300,423	1,490
Sand	20.2	192,928	953	-	-	-	3.3	29,583	910	23.5	222,511	947
Sandy Loam	87.2	650,882	747	-	-	-	80.8	547,985	678	168.0	1,198,867	714
Sandy Loam (defaulted)	0.9	5,453	639	-	-	-	5.4	30,050	552	6.3	35,504	564
Silt Loam	308.1	1,920,197	623	-	-	-	208.0	1,244,200	598	516.2	3,164,397	613
TOTALS	520.2	3,476,058	668	-	-	-	498.3	3,441,185	691	1,018.4	6,917,243	679

Appendix Table E 2003 Water Demand by Aquifer with Average Management

Water Source	Surface Water			Reclaimed Water			Groundwater			Total		
Agriculture Soil Texture	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)
Others	1.7	8,912	519	-	-	-	60.2	337,094	560	61.9	346,006	559
Between Big & Little Qual	-	141	662	-	-	-	49.1	491,408	1,001	49.1	491,549	1,000
Cassidy	11.2	87,692	780	-	-	-	21.3	173,034	811	32.6	260,726	800
Cedar, North Holden Lake	-	-	-	-	-	-	1.2	7,480	644	1.2	7,480	644
Cedar, Yellow Point, N.O	19.6	153,752	784	-	-	-	39.8	345,878	868	59.5	499,630	840
Errington	115.7	619,958	536	-	-	-	51.4	282,038	549	167.1	901,996	540
Errington, Morison Creek	60.0	283,150	472	-	-	-	21.4	124,190	579	81.5	407,340	500
Extension (Nanaimo)	11.1	78,186	702	-	-	-	1.1	6,903	610	12.3	85,088	694
Gabriola excluding North	0.7	4,735	669	-	-	-	3.6	25,734	724	4.3	30,470	715
Gabriola Northern Area	-	-	-	-	-	-	0.5	3,716	803	0.5	3,716	803
Lantzville	17.5	106,631	681	-	-	-	10.3	65,186	637	27.8	171,817	617
Little Qualicum R. Valley	29.1	183,740	632	-	-	-	-	-	-	29.1	183,740	632
Nanaimo	1.1	5,232	487	-	-	-	5.4	22,956	422	6.5	28,188	432
Nanoose Creek	138.2	1,092,091	790	-	-	-	12.9	104,369	806	151.2	1,196,460	791
Nanoose Hill	0.5	2,775	599	-	-	-	3.0	17,301	580	3.4	20,076	583
Parksville	22.7	183,302	833	-	-	-	101.6	824,610	837	124.3	1,007,914	837
Qualicum	61.2	432,900	707	-	-	-	81.7	407,986	499	142.9	840,886	588
South Wellington	21.2	188,872	892	-	-	-	7.3	47,776	658	28.4	236,648	832
Spider Lk nr Home Lk	-	-	-	-	-	-	0.2	2,493	1,014	0.2	2,493	1,014
Thames River to Mplegaur	-	-	-	-	-	-	1.1	8,948	787	1.1	8,948	787
Upper reaches of Whisky C	-	-	-	-	-	-	0.2	1,717	934	0.2	1,717	934
Westwood Lake, Nanaimo	8.7	43,987	507	-	-	-	24.7	140,367	569	33.4	184,354	553
TOTALS	520.2	3,476,058	668	-	-	-	498.3	3,441,185	691	1,018.4	6,917,243	679

Appendix Table F 2003 Water Demand by Local Government with Average Management

Water Source	Surface Water			Reclaimed Water			Groundwater			Total		
Agriculture Local Government	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)
Lantzville	4.8	39,973	835	-	-	-	8.8	57,711	652	13.6	97,685	716
Nanaimo	506.5	3,365,962	665	-	-	-	468.4	3,229,302	689	975.0	6,595,265	676
Parksville	8.8	70,123	793	-	-	-	0.6	4,330	765	9.4	74,452	791
Qualicum Beach	-	-	-	-	-	-	20.4	149,841	734	20.4	149,841	734
TOTALS	520.2	3,476,058	668	-	-	-	498.3	3,441,185	691	1,018.4	6,917,243	679

Appendix Table G 2003 Management Comparison on Irrigation Demand and Percolation Volumes

Water Source	Surface Water				Reclaimed Water				Groundwater				Total				
Agriculture Management	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)	Deep Percolation (m ³)	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)	Deep Percolation (m ³)	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)	Deep Percolation (m ³)	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)	Deep Percolation (m ³)	Percolation (m ³ /ha)
Poor	520.2	3,546,161	682	403,820	-	-	-	-	498.3	3,509,179	704	486,500	1,018.4	7,055,340	693	890,320	874
Avg	520.2	3,476,058	668	333,717	-	-	-	-	498.3	3,448,244	692	425,564	1,018.4	6,924,302	680	759,281	746
Good	520.2	3,405,955	655	263,614	-	-	-	-	498.3	3,387,308	680	364,629	1,018.4	6,793,263	667	628,243	617

Appendix Table H 2003 Percolation Volumes by Irrigation System with Average Management

Water Source	Surface Water			Reclaimed Water			Groundwater			Total			
	Irrigated Area (ha)	Irrigation Demand (m ³)	Deep Percolation (m ³)	Irrigated Area (ha)	Irrigation Demand (m ³)	Deep Percolation (m ³)	Irrigated Area (ha)	Irrigation Demand (m ³)	Deep Percolation (m ³)	Irrigated Area (ha)	Irrigation Demand (m ³)	Deep Percolation (m ³)	Percolation (m ³ /ha)
Drip	1.8	7,850	809	-	-	-	80.3	304,797	24,458	82.1	312,647	25,268	308
Flood	-	-	-	-	-	-	20.2	300,423	68,456	20.2	300,423	68,456	3,389
Golfsprinkler	4.1	27,800	5,630	-	-	-	34.6	274,234	41,387	38.7	302,035	47,017	1,215
Handline	35.3	278,731	33,247	-	-	-	20.2	119,516	11,425	55.5	398,247	44,672	805
Landscapesprinkler	12.0	103,988	21,052	-	-	-	9.7	71,816	10,475	21.8	175,804	31,528	1,446
Microsprinkler	0.7	13,071	3,735	-	-	-	11.3	207,336	59,239	12.0	220,407	62,973	5,248
Overtreedrip	-	-	-	-	-	-	7.3	31,721	3,077	7.3	31,721	3,077	422
SDI	-	-	-	-	-	-	4.0	17,605	1,100	4.0	17,605	1,100	275
Sprinkler	407.6	2,528,947	214,082	-	-	-	130.9	745,651	66,622	538.4	3,274,598	280,704	521
Ssovertree	-	-	-	-	-	-	12.9	80,578	12,943	12.9	80,578	12,943	1,003
Ssundertree	1.8	14,558	1,175	-	-	-	-	-	-	1.8	14,558	1,175	653
Travgun	56.8	501,113	53,987	-	-	-	116.2	865,943	84,707	173.0	1,367,056	138,694	802
Wheelline	-	-	-	-	-	-	50.8	421,566	41,676	50.8	421,566	41,676	820
TOTALS	520.2	3,476,058	333,717	-	-	-	498.3	3,441,185	425,564	1,018.4	6,917,243	759,281	746

Appendix Table I 2003 Crop Water Demand for Improved Irrigation System Efficiency and Good Management

Water Source	Surface Water			Reclaimed Water			Groundwater			Total		
	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)
Apple	0.2	864	472	-	-	-	5.6	26,793	483	5.7	27,657	482
Berry	-	-	-	-	-	-	9.3	34,717	374	9.3	34,717	374
Blueberry	1.5	6,964	450	-	-	-	59.6	231,016	388	61.1	237,980	389
Cranberry	-	-	-	-	-	-	20.2	107,953	535	20.2	107,953	535
Forage	462.2	2,954,296	639	-	-	-	242.3	1,719,607	710	704.4	4,673,903	664
Golf	16.1	130,744	810	-	-	-	37.5	293,451	782	53.7	424,194	790
Grape	-	-	-	-	-	-	4.8	6,551	137	4.8	6,551	137
Greenhouse	0.7	13,071	1,801	-	-	-	11.3	207,336	1,842	12.0	220,407	1,839
Nursery Floriculture	-	-	-	-	-	-	0.2	619	328	0.2	619	328
Nursery Shrubs/Trees	0.6	3,662	292	-	-	-	16.9	63,691	280	17.5	67,352	280
Pasture/Grass	11.7	81,169	691	-	-	-	7.5	49,228	657	19.2	130,396	678
Raspberry	0.2	648	333	-	-	-	7.7	25,612	331	7.9	26,260	331
Recreational Turf	-	-	-	-	-	-	6.8	46,711	687	6.8	46,711	687
Strawberry	-	-	-	-	-	-	3.3	10,412	316	3.3	10,412	316
Sweetcorn	-	-	-	-	-	-	29.0	111,366	384	29.0	111,366	384
Turf Farm	20.9	176,479	845	-	-	-	3.9	30,359	779	24.8	206,838	835
Vegetable	5.9	20,303	341	-	-	-	32.6	105,277	323	38.5	125,581	326
TOTALS	520.2	3,388,199	651	-	-	-	498.3	3,070,699	616	1,018.4	6,458,897	634

Appendix Table J 2003 Water Demand for Frost Protection, Harvesting and Other Use with Average Management

Water Source	Surface Water			Reclaimed Water			Groundwater			Total		
Agriculture Crop Group	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)
Cranberry Frost Protection	-	-	-	-	-	-	20.2	2,017	10	20.2	2,017	10
Cranberry Harvesting	-	-	-	-	-	-	20.2	5,042	25	20.2	5,042	25
Greenhouse	0.7	13,071	1,801	-	-	-	11.3	207,336	1,842	12.0	220,407	1,839
Nursery Pot	0.3	2,759	819	-	-	-	3.5	26,300	753	3.8	29,059	759
TOTALS	1.1	15,830	1,490	-	-	-	55.1	240,694	437	56.1	256,525	457

Appendix Table K 2003 Water Demand by Animal Type

Animal Type	Demand (m ³)
Beef	29,090
Dairy - dry	10,330
Dairy - milking	17,560
Goats	712
Horses	13,596
Poultry - broiler	6,250
Poultry - laying	3,309
Sheep	4,126
Swine	1,382
TOTALS	86,356

Appendix Table L Climate Change Water Demand Circa 2050 for High Demand Year with Good Management Using Current Crops and Irrigation Systems

Climate Change	rcp26			rcp45			rcp85			Average		
	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)
Year												
2053	1,018.4	8,344,478	819	1,018.4	5,398,171	530	1,018.4	9,075,637	891	1,018.4	7,606,095	747
2056	1,018.4	5,085,250	603	1,018.4	7,385,931	725	1,018.4	5,337,802	524	1,018.4	5,936,328	617
2059	1,018.4	4,320,083	424	1,018.4	7,843,401	770	1,018.4	8,699,672	854	1,018.4	6,954,385	683
Average	1,018.4	5,916,604	615	1,018.4	6,875,834	675	1,018.4	7,704,370	756	1,018.4	6,832,269	682

Appendix Table M Buildout Crop Water Demand for 2003 Climate Data with Good Management

Water Source	Surface Water			Reclaimed Water			Groundwater			Total		
Agriculture Crop Group	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)
Apple	0.2	1,398	763	-	-	-	5.6	31,884	574	5.7	33,282	580
Berry	-	-	-	-	-	-	9.3	47,942	516	9.3	47,942	516
Blueberry	426.5	1,574,807	369	-	-	-	63.2	250,166	396	489.6	1,824,973	373
Cranberry	-	-	-	-	-	-	20.2	300,423	1,490	20.2	300,423	1,490
Forage	2,304.5	16,087,534	698	-	-	-	311.3	2,248,286	722	2,615.8	18,335,820	701
Golf	16.1	130,744	810	-	-	-	37.5	293,451	782	53.7	424,194	790
Grape	-	-	-	-	-	-	4.8	10,343	217	4.8	10,343	217
Greenhouse	0.7	13,071	1,801	-	-	-	11.3	207,336	1,842	12.0	220,407	1,839
Nursery Floriculture	-	-	-	-	-	-	0.2	619	328	0.2	619	328
Nursery Shrubs/Trees	0.6	3,662	292	-	-	-	16.9	63,691	280	17.5	67,352	280
Pasture/Grass	518.4	3,261,259	629	-	-	-	25.6	163,875	641	544.0	3,425,134	630
Raspberry	0.2	1,105	568	-	-	-	7.7	36,431	470	7.9	37,536	473
Recreational Turf	-	-	-	-	-	-	6.8	46,711	687	6.8	46,711	687
Strawberry	-	-	-	-	-	-	3.3	13,564	411	3.3	13,564	411
Sweetcorn	-	-	-	-	-	-	29.0	111,366	384	29.0	111,366	384
Turf Farm	20.9	176,479	845	-	-	-	3.9	30,359	779	24.8	206,838	835
Vegetable	250.7	785,540	313	-	-	-	34.6	190,458	550	285.4	975,998	342
TOTALS	3,538.9	22,035,599	623	-	-	-	591.1	4,046,905	685	4,129.9	26,082,504	632

Appendix Table N Buildout Crop Water Demand for Climate Change Data Circa 2050 and Good Management

Climate Change	rcp26			rcp45			rcp85			Average		
Year	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)
2053	4,129.9	33,388,424	808	4,129.9	20,952,630	507	4,129.9	34,589,240	838	4,129.9	29,643,431	718
2056	4,129.9	20,986,528	570	4,129.9	28,502,084	690	4,129.9	20,866,430	505	4,129.9	23,451,681	588
2059	4,129.9	17,064,005	413	4,129.9	31,272,821	757	4,129.9	33,347,438	807	4,129.9	27,228,088	659
Average	4,129.9	23,812,986	597	4,129.9	26,909,178	651	4,129.9	29,601,036	717	4,129.9	26,774,400	655

Appendix Table O Buildout Irrigation System Demand for 2003 Climate Data and Good Management

Water Source	Surface Water			Reclaimed Water			Groundwater			Total		
Agriculture Irrigation System	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)
Drip	671.5	2,324,003	346	-	-	-	85.9	318,651	371	757.4	2,642,654	349
Flood	-	-	-	-	-	-	20.2	300,423	1,490	20.2	300,423	1,490
Golfsprinkler	4.1	27,713	674	-	-	-	34.6	269,402	779	38.7	297,116	767
Handline	35.3	275,061	780	-	-	-	20.2	117,166	580	55.5	392,227	707
Landscapesprinkler	12.0	103,030	857	-	-	-	9.7	70,760	727	21.8	173,790	799
Microsprinkler	0.7	13,071	1,801	-	-	-	11.3	207,336	1,842	12.0	220,407	1,839
Overtreedrip	-	-	-	-	-	-	7.3	31,223	428	7.3	31,223	428
SDI	-	-	-	-	-	-	4.0	17,267	432	4.0	17,267	432
Sprinkler	2,756.6	18,789,707	682	-	-	-	218.0	1,372,965	630	2,974.6	20,162,672	678
Ssovertime	-	-	-	-	-	-	12.9	78,247	609	12.9	78,247	609
Ssundertree	1.8	13,992	784	-	-	-	-	-	-	1.8	13,992	784
Travgun	56.8	489,021	860	-	-	-	116.2	847,820	730	173.0	1,336,840	773
Wheelline	-	-	-	-	-	-	50.8	415,646	818	50.8	415,646	818
TOTALS	3,538.9	22,035,599	623	-	-	-	591.1	4,046,905	685	4,129.9	26,082,504	632

Appendix Table P Buildout Water Demand by Aquifer for 2003 Climate Data and Good Management

Water Source	Surface Water			Reclaimed Water			Groundwater			Total		
Aquifer	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)
Others	160.1	1,003,204	627	-	-	-	84.2	517,420	615	244.3	1,520,624	623
Between Big & Little Qual	864.6	5,055,148	585	-	-	-	49.1	488,072	994	913.7	5,543,220	607
Between Big Qualicum R. &	13.4	97,291	728	-	-	-	-	-	-	13.4	97,291	728
Cassidy	301.9	1,967,140	652	-	-	-	60.3	448,667	744	362.2	2,415,807	667
Cedar, North Holden Lake	-	-	-	-	-	-	1.2	7,174	617	1.2	7,174	617
Cedar, Yellow Point, N.O	60.5	440,559	728	-	-	-	44.5	373,481	839	105.0	814,040	775
Errington	506.1	3,096,658	612	-	-	-	51.4	277,193	539	557.5	3,373,851	605
Errington, Morison Creek	313.4	1,535,279	490	-	-	-	21.4	121,439	566	334.9	1,656,718	495
Extension (Nanaimo)	38.6	258,380	669	-	-	-	1.1	6,717	594	39.8	265,096	666
Gabriola excluding North	0.7	4,536	641	-	-	-	3.6	24,867	700	4.3	29,403	690
Gabriola Northern Area	-	-	-	-	-	-	0.5	3,688	796	0.5	3,688	796
Lantzville	29.9	162,786	662	-	-	-	10.3	64,233	626	40.2	227,020	594
Little Qualicum R. Valley	74.0	506,479	685	-	-	-	-	-	-	74.0	506,479	685
Madrona Point / Parksville	84.8	683,502	806	-	-	-	-	-	-	84.8	683,502	806
Nanaimo	8.0	38,908	485	-	-	-	5.4	22,431	412	13.5	61,340	455
Nanoose Creek	301.9	2,065,529	684	-	-	-	24.0	180,824	752	325.9	2,246,354	689
Nanoose Hill	0.5	3,327	617	-	-	-	3.0	16,713	561	3.5	20,040	569
Parksville	82.3	516,019	737	-	-	-	101.6	811,140	828	184.0	1,327,159	788
Qualicum	578.2	3,802,653	658	-	-	-	81.7	396,973	486	659.9	4,199,626	636
South Wellington	67.9	543,633	801	-	-	-	21.4	135,480	633	89.3	679,113	760
Spider Lk nr Home Lk	0.1	691	670	-	-	-	0.2	2,440	992	0.3	3,131	897
Thames River to Maplegaur	-	-	-	-	-	-	1.1	8,928	786	1.1	8,928	786
Upper reaches of Whisky C	20.3	102,015	502	-	-	-	0.2	1,717	934	20.5	103,732	505
Westwood Lake, Nanaimo	31.6	151,861	481	-	-	-	24.7	137,307	557	56.3	289,168	514
TOTALS	3,538.9	22,035,599	623	-	-	-	591.1	4,046,905	685	4,129.9	26,082,504	632

Appendix Table Q Buildout Water Demand by Local Government for 2003 Climate Data and Good Management

Water Source	Surface Water			Reclaimed Water			Groundwater			Total		
Agriculture Local Government	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m ³)	Avg. Req. (mm)
Lantzville	17.2	97,655	568	-	-	-	8.8	56,934	643	26.0	154,589	594
Nanaimo	3,377.0	21,078,161	624	-	-	-	561.2	3,838,987	684	3,938.2	24,917,148	633
Parksville	32.2	229,950	713	-	-	-	0.6	4,236	748	32.8	234,185	714
Qualicum Beach	112.4	629,833	560	-	-	-	20.4	146,749	719	132.9	776,582	585
TOTALS	3,538.9	22,035,599	623	-	-	-	591.1	4,046,905	685	4,129.9	26,082,504	632