

# AGRICULTURE WATER DEMAND MODEL

## Report for the Similkameen Watershed

October 2013





# AGRICULTURE WATER DEMAND MODEL

## Report for the Similkameen Watershed

### Authors

**Ted van der Gulik, P.Eng.**

Senior Engineer  
B.C. Ministry of Agriculture  
Sustainable Agriculture Management Branch  
Abbotsford, BC

**Denise Neilsen, Ph.D.**

Research Scientist  
Agriculture and Agri-Food Canada  
Pacific Agri-Food Research Centre  
Summerland, BC

**Ron Fretwell**

Program Developer  
RHF Systems Ltd.  
Kelowna, BC

**Andrew Petersen, P.Ag.**

Regional Resource Specialist  
B.C. Ministry of Agriculture  
Sustainable Agriculture Management Branch  
Kamloops, BC

**Stephanie Tam, P.Eng.**

Water Management Engineer  
B.C. Ministry of Agriculture  
Sustainable Agriculture Management Branch  
Abbotsford, BC

### Funded By

**Canada-British Columbia Water Supply Expansion Program**



**October 2013**

## **DISCLAIMER**

The data that is presented in this report provides the best estimates for agriculture water demand that can be generated at this time. While every effort has been made to ensure the accuracy and completeness of the information, the information should not be considered as final. The Government of Canada, the BC Ministry of Agriculture, and the BC Agriculture Council or its directors, agents, employees, or contractors will not be liable for any claims, damages, or losses of any kind whatsoever arising out of the use of, or reliance upon, this information.

# Table of Contents

---

<b>ACKNOWLEDGEMENTS .....</b>	<b>5</b>
<b>BACKGROUND.....</b>	<b>6</b>
<b>METHODOLOGY .....</b>	<b>7</b>
Cadastre.....	7
Land Use Survey.....	8
Soil Information .....	10
Climate Information .....	11
<b>MODEL CALCULATIONS.....</b>	<b>12</b>
Crop .....	12
Irrigation .....	12
Soil .....	12
Climate .....	13
Agricultural Water Demand Equation .....	13
<b>LIVESTOCK WATER USE.....</b>	<b>18</b>
<b>DEFINITION AND CALCULATION OF INDIVIDUAL TERMS USED IN THE IRRIGATION WATER DEMAND EQUATION .....</b>	<b>19</b>
Growing Season Boundaries.....	19
Evapotranspiration ( $ET_o$ ) .....	21
Availability Coefficient (AC) .....	21
Rooting Depth (RD).....	21
Stress Factor (stressFactor) .....	22
Available Water Storage Capacity (AWSC).....	22
Maximum Soil Water Deficit (MSWD).....	22
Deep Percolation Factor (SoilpercFactor) .....	22
Maximum Evaporation Factor (maxEvaporation) .....	23
Irrigation Efficiency ( $I_e$ ) .....	23
Soil Water Factor (swFactor) .....	23
Early Season Evaporation Factor (earlyEvaporationFactor) .....	23
Crop Coefficient ( $K_c$ ).....	23
Growing Degree Days (GDD).....	24
Frost Indices.....	24
Corn Heat Units (CHU).....	24
Corn Season Start and End.....	25
Tsum Indices .....	25
Wet/Dry Climate Assessment.....	25
Groundwater Use .....	25
<b>LAND USE RESULTS.....</b>	<b>26</b>

<b>AGRICULTURAL WATER DEMAND RESULTS.....</b>	<b>30</b>
Annual Crop Water Demand – Tables A and B .....	30
Annual Water Demand by Irrigation System – Table C .....	30
Annual Water Demand by Soil Texture – Table D .....	30
Annual Water Demand by Water Purveyor – Table E .....	31
Irrigated Area within Local Governments – Table F .....	31
Irrigation Management Factors – Table G .....	31
Deep Percolation – Table H .....	32
Improved Irrigation Efficiency and Good Management – Table I .....	32
Livestock Water Use – Table J .....	32
Climate Change Water Demand for 2050 – Table K .....	33
Agricultural Buildout Crop Water Demand Using 2003 Climate Data – Table L .....	34
Agricultural Buildout Crop Water Demand for 2050 – Table M .....	35
Irrigation Systems Used for the Buildout Scenario for 2003 Climate Data– Table N .....	35
Water Demand for the Buildout Area by Water Purveyor 2003 Climate Data – Table O .....	35
Water Demand for the Buildout Area by Local Government 2003 Climate Data – Table P .....	36
<b>LITERATURE .....</b>	<b>37</b>
<b>APPENDIX TABLES .....</b>	<b>38</b>

## List of Figures

Figure 1	Map of ALR in the Similkameen Watershed .....	6
Figure 2	Map of the Similkameen Watershed Overlaid with Map Sheets .....	7
Figure 3	Land Use Survey .....	8
Figure 4	GIS Map Sheet .....	8
Figure 5	Cadastre with Polygons .....	9
Figure 6	GIS Model Graphics .....	10
Figure 7	Similkameen Climate Stations .....	11
Figure 8	Water Areas in the Similkameen Watershed .....	27
Figure 9	ALR Areas in the Similkameen Watershed .....	28
Figure 10	Land Parcels in the Similkameen Watershed .....	28
Figure 11	Higher Productive Groundwater Aquifers in the Similkameen Watershed .....	29
Figure 12	Annual ET and Effective Precipitation in 2050's .....	33
Figure 13	Future Irrigation Demand for All Outdoor Uses in the Okanagan in Response to Observed Climate Data (Actuals) and Future Climate Data Projected from a Range of Global Climate Models .....	34
Figure 14	Similkameen Watershed Irrigation Expansion Potential .....	36

## List of Tables

Table 1	Livestock Water Demand (Litres/day) .....	18
Table 2	Overview of Similkameen Basin's Land and Inventoried Area .....	26
Table 3	Summary of Primary Agricultural Activities of Inventoried Parcels where Primary Land Use is agriculture in Similkameen .....	27
Table 4	Irrigation Management Factors .....	31

## Acknowledgements

---

There are over twenty people that have been involved with the preparation and collection of data for the development of the Water Demand Model in the Similkameen. The authors wish to express appreciation to the following for their in-kind contribution for the tasks noted.


Rae Anne Bell	Agriculture and Agri-Food Canada	Land Use Inventory
Linda Decker	Contractor	GIS data preparation
Beckie Rozander	Contractor	Land Use Inventory
Alex Cannon	Environment Canada	Climate data downscaling
Bill Taylor	Environment Canada	Climate data layer
Andrew Petersen	Ministry of Agriculture	Land Use Inventory
Linda Hokanson	Ministry of Agriculture	Cover and Publication Design
Sam Lee	Ministry of Agriculture	GIS data preparation
Stephanie Tam	Ministry of Agriculture	Model Development

## Background

The Agriculture Water Demand Model (AWDM) was developed in the Okanagan Watershed. It was developed in response to rapid population growth, drought conditions from climate change, and the overall increased demand for water. Many of the watersheds in British Columbia (BC) are fully allocated or will be in the next 15 to 20 years. The AWDM helps to understand current agricultural water use and helps to fulfil the Province's commitment under the "*Living Water Smart – BC Water Plan*" to reserve water for agricultural lands. The Model can be used to establish agricultural water reserves throughout the various watersheds in BC by providing current and future agricultural water use data.

Climate change scenarios developed by the University of British Columbia (UBC) and the Pacific Agri-Food Research Centre (PARC) in Summerland predict an increase in agricultural water demand due to warmer and longer summers and lower precipitation during summer months in the future.

The Agriculture Water Demand Model was developed to provide current and future agricultural water demands. The Model calculates water use on a property-by-property basis, and sums each property to obtain a total water demand for the entire basin or each sub-basin. Crop, irrigation system type, soil texture and climate data are used to calculate the water demand. Climate data from 2003 was used to present information on one of the hottest and driest years on record, and 1997 data was used to represent a wet year. Lands within the Agriculture Land Reserve (ALR), depicted in green in Figure 1, were included in the project.




**Figure 1 Map of ALR in the Similkameen Watershed**



## Methodology

The Model is based on a Geographic Information System (GIS) database that contains information on cropping, irrigation system type, soil texture and climate. An explanation of how the information was compiled for each is given below. The survey area included all properties within the ALR and areas that were zoned for agriculture by the local governments. The inventory was undertaken by Ministry of Agriculture (AGRI) staff, hired professional contractors and summer students. Figure 2 provides a schematic of the map sheets that were generated to conduct the survey.



**Figure 2 Map of the Similkameen Watershed Overlaid with Map Sheets**

### Cadastre

Cadastre information was provided by the Okanagan-Similkameen Regional District and local governments in the Similkameen Watershed. The entire Similkameen Watershed is covered in one dataset which allows the Model to report out on each sub-basin, local government, water purveyor or groundwater aquifer. A GIS technician used aerial photographs to conduct an initial review of cropping information by cadastre, and divided the cadastre into polygons that separate farmstead and driveways from cropping areas. Different crops were also separated into different polygons if the difference could

be identified on the aerial photographs. This data was entered into a database that was used by the field teams to conduct and complete the land use survey.

## Land Use Survey


The survey maps and database were created by AGRI for the survey crew to enter data about each property. Surveys were done during the summer of 2008. The survey crew drove by each property where the team checked the database for accuracy using visual observation and the aerial photographs on the survey maps. A Professional Agrologist verified what was on the site, and a GIS technician altered the codes in the database as necessary (Figure 3). Corrections were handwritten on the maps. The maps were then brought back to the office to have the hand-drawn lines digitized into the GIS system and have the additional polygons entered into the database.

Once acquired through the survey, the land use data was brought into the GIS to facilitate analysis and produce maps. Digital data, in the form of a database and GIS shape files (for maps), is available upon request through a data sharing agreement with the Ministry of Agriculture.



**Figure 3 Land Use Survey**

Figure 4 provides an example of a map sheet. The Similkameen Watershed was divided into 250 map sheets. Each map sheet also had a key map to indicate where it was located in the region.



**Figure 4 GIS Map Sheet**

The smallest unit for which water use is calculated are the polygons within each cadastre. A polygon is determined by a change in land use or irrigation system within a cadastre. Polygons are designated as blue lines within each cadastre as shown in Figures 4 and 5. The Similkameen watershed encompasses 2,599 inventoried land parcels that are in or partially in the ALR. There are a total of 6,322 polygons (land covers) generated for the Similkameen Watershed for this project. Figure 5 provides an enhanced view of a cadastre containing three polygons. Each cadastre has a unique identifier as does each polygon. The polygon identifier is acknowledged by PolygonID. This allows the survey team to call up the cadastre in the database, review the number of polygons within the cadastre and ensure the land use is coded accurately for each polygon.




**Figure 5 Cadastre with Polygons**

## Soil Information

Soil information was obtained digitally from the Ministry of Environment's Terrain and Soils Information System. The Computer Assisted Planning and Map Production application (CAPAMP) provided detailed (1:20,000 scale) soil surveys that were conducted in the Lower Mainland, on Southeast Vancouver Island, and in the Okanagan-Similkameen areas during the early 1980s. Products developed include soil survey reports, maps, agriculture capability and other related themes. Soil information required for this project was the soil texture (loam, etc.), the available water storage capacity and the peak infiltration rate for each texture type.

The intersection of soil boundaries with the cadastre and land use polygons creates additional polygons that the Model uses to calculate water demand. Figure 6 shows how the land use information is divided into additional polygons using the soil boundaries. The Model calculates water demand using every different combination of crop, soil and irrigation system as identified by each polygon.



**Figure 6 GIS Model Graphics**

The next section will discuss about climate information where the climate grid does not develop additional polygons. Each polygon has the climate grid cell which is prominent for that polygon assigned to it.




## Climate Information

The agricultural water demand is calculated using climate, crop, irrigation system and soil information data. The climate in the interior region is quite diverse. The climate generally gets cooler and wetter from south to north and as elevation increases. To incorporate the climatic diversity, climate layers were developed for the entire region on a 500 m x 500 m grid. Each grid cell contains daily climate data, minimum and maximum temperature ( $T_{\min}$  and  $T_{\max}$ ), and precipitation which allows the Model to calculate a daily reference evapotranspiration rate ( $ET_o$ ) value. A range of agro-climatic indices such as growing degree days (GDD), corn heat units (CHU), frost free days and temperature sum (Tsum) can also be calculated for each grid cell based on temperature data. These values are used to determine seeding dates and the length of the growing season in the Model.

The climate dataset is generated by using existing data from climate stations in and around the Similkameen Basin from 1961 to 2006, and other station data close to the region. This climate dataset was then downscaled to provide a climate data layer for the entire watershed on the 500 m x 500 m grid. Since the Similkameen Watershed is a little over 7,571 square km, there are a total of 30,284 grid cells populated with daily data. A detailed description of the Model can be obtained by contacting the authors.

Existing climate stations that were used to determine the climate coverage are shown in Figure 7. The attributes attached to each climate grid cell include:

- Latitude
- Longitude
- Elevation
- Aspect
- Slope
- Daily Precipitation
- Daily  $T_{\min}$  and  $T_{\max}$



**Figure 7 Similkameen Climate Stations**

A climate database generated contains  $T_{\min}$ ,  $T_{\max}$ ,  $T_{\text{mean}}$  and Precipitation for each day of the year from 1961 until 2006. The parameters that need to be selected, calculated and stored within the Model are evapotranspiration ( $ET_o$ ), Tsum of 600 (for the Similkameen region), effective precipitation (EP), frost free days, GDD with base temperatures of 5 °C and 10 °C, CHU, and first frost date. These climate and crop parameters are used to determine the growing season length as well as the beginning and end of the growing season in Julian day.

## **Model Calculations**

---

The model calculates the water demand for each polygon by using crop, irrigation, soil and climate parameters as explained below. Each polygon has been assigned an ID number as mentioned previously.

### **Crop**

The CropID is an attribute of the PolygonID as each polygon will contain a single crop. The crop information (observed during the land use survey) has been collected and stored with PolygonID as part of the land use survey. CropID will provide cropping attributes to the Model for calculating water use for each polygon. CropID along with the climate data will also be used to calculate the growing season length and the beginning and end of the growing season. The attributes for CropID include rooting depth, availability coefficient, crop coefficient and a drip factor.

Rooting depth is the rooting depth for a mature crop in a deep soil.

An availability coefficient is assigned to each crop. The availability coefficient is used with the IrrigID to determine the soil moisture available to the crop for each PolygonID.

The crop coefficient adjusts the calculated  $ET_o$  for the stages of crop growth during the growing season. Crop coefficient curves have been developed for every crop. The crop coefficient curve allows the Model to calculate water demand with an adjusted daily  $ET_o$  value throughout the growing season.

The drip factor is used in the water use calculation for polygons where drip irrigation systems are used. Since the Model calculates water use by area, the drip factor adjusts the percentage of area irrigated by the drip system for that crop.

### **Irrigation**

The IrrigID is an attribute of the PolygonID as each polygon will have a single irrigation system type operating. The irrigation information has been collected and stored (as observed during the land use survey) with the land use data. The land use survey determined if a polygon had an irrigation system operating, what the system type was, and if the system was being used. The IrrigID has an irrigation efficiency listed as an attribute.

Two of the IrrigID, Overtreedrip and Overtreemicro are polygons that have two systems in place. Two irrigation ID's occur when an overhead irrigation system has been retained to provide crop cooling or frost protection. In this case, the efficiencies used in the Model are the drip and microsprinkler efficiencies.

### **Soil**

The soil layer came from CAPAMP at the Ministry of Environment. In addition, soil data provided by Agriculture and Agri-Food Canada (AAFC) was also used to generate multiple soil layers within each polygon. Each parcel was assigned the most predominant soil polygon, and then for each crop field

within that soil polygon, the most predominant texture within the crop's rooting depth was determined and assigned to the crop field.

Note that textures could repeat at different depths – the combined total of the thicknesses determined the most predominant texture. For example, a layer of 20 cm sand, followed by 40 cm clay and then 30 cm of sand would have sand be designated as the predominant soil texture.

The attributes attached to the SoilID is the Available Water Storage Capacity (AWSC) which is calculated using the soil texture and crop rooting depth.

The Maximum Soil Water Deficit (MSWD) is calculated to determine the parameters for the algorithm that is used to determine the Irrigation Requirement (IR). The Soil Moisture Deficit at the beginning of the season is calculated using the same terms as the MSWD.

## Climate

The climate data in the Model is used to calculate a daily reference evapotranspiration rate ( $ET_o$ ) for each climate grid cell. The data that is required to calculate this value are:

- Elevation, metres (m)
- Latitude, degrees ( $^{\circ}$ )
- Minimum Temperature, degree Celsius ( $^{\circ}C$ )
- Maximum Temperature, degree Celsius ( $^{\circ}C$ )
- Classification as Coastal or Interior
- Classification as Arid or Humid
- Julian Day

Data that is assumed or are constants in this calculation are:

- |   |   |
|---|---|
| • Wind speed                                  | 2 m/s   |
| • Albedo or canopy reflection coefficient,    | 0.23  |
| • Solar constant, $G_{sc}$                    | $0.082 \text{ MJ}^{-2}\text{min}^{-1}$  |
| • Interior and Coastal coefficients, $K_{Rs}$ | 0.16 for interior locations<br>0.19 for coastal locations                               |
| • Humid and arid region coefficients, $K_o$   | 0 $^{\circ}C$ for humid/sub-humid climates<br>2 $^{\circ}C$ for arid/semi-arid climates |

## Agricultural Water Demand Equation

The Model calculates the Agriculture Water Demand (AWD) for each polygon, as a unique crop, irrigation system, soil and climate data is recorded on a polygon basis. The polygons are then summed to determine the AWD for each cadastre. The cadastre water demand values are then summed to determine AWD for the basin, sub-basin, water purveyor or local government. The following steps provide the process used by the Model to calculate Agricultural Water Demand. The entire process is outlined although not all of the steps may be used for the Similkameen, e.g., flood harvesting. Detailed information is available on request.

## 1. *Pre-Season Soil Moisture Content*

Prior to the start of each crop's growing season, the soil's stored moisture content is modelled using the soil and crop evaporation and transpiration characteristics and the daily precipitation values. Precipitation increases the soil moisture content and evaporation (modelled using the reference potential evapotranspiration) depletes it. In general, during the pre-season, the soil moisture depth cannot be reduced beyond the maximum evaporation depth; grass crops in wet climates, however, can also remove moisture through crop transpiration.

The process used to model the pre-season soil moisture content is:

1. Determine whether the modelling area is considered to be in a *wet* or *dry* climate (see *Wet/Dry Climate Assessment*), and retrieve the early season evaporation factor in the modelling area
2. For each crop type, determine the start of the growing season (see *Growing Season Boundaries*)
3. For each crop and soil combination, determine the *maximum soil water deficit* (MSWD) and *maximum evaporation factor* (maxEvaporation)
4. Start the initial storedMoisture depth on January 1 at the MSWD level
5. For each day between the beginning of the calendar year and the crop's growing season start, calculate a new storedMoisture from:
  - a. the potential evapotranspiration ( $ET_o$ )
  - b. the early season evaporation factor (earlyEvaporationFactor)
  - c. the effective precipitation (EP) = actual precipitation x earlyEvaporationFactor
  - d. daily Climate Moisture Deficit (CMD) =  $ET_o - EP$
  - e. storedMoisture = previous day's storedMoisture – CMD

A negative daily CMD (precipitation in excess of the day's potential evapotranspiration) adds to the stored moisture level while a positive climate moisture deficit reduces the amount in the stored moisture reservoir. The stored moisture cannot exceed the maximum soil moisture deficit; any precipitation that would take the stored moisture level above the MSWD gets ignored.

For all crops and conditions except for grass in wet climates, the stored moisture content cannot drop below the maximum soil water deficit minus the maximum evaporation depth; without any crop transpiration in play, only a certain amount of water can be removed from the soil through evaporative processes alone. Grass in wet climates does grow and remove moisture from the soil prior to the start of the irrigation season however. In those cases, the stored moisture level can drop beyond the maximum evaporation depth, theoretically to 0.

Greenhouses and mushroom barns have no stored soil moisture content.

## 2. *In-Season Precipitation*

During the growing season, the amount of precipitation considered effective (EP) depends on the overall wetness of the modelling area's climate (see *Wet/Dry Climate Assessment*). In dry climates, the first 5 mm of precipitation is ignored, and the EP is calculated as 75% of the remainder:

$$EP = (\text{Precip} - 5) \times 0.75$$



In wet climates, the first 5 mm is included in the EP. The EP is 75% of the actual precipitation:

$$EP = \text{Precip} \times 0.75$$

Greenhouses and mushroom barns automatically have an EP value of 0.

### 3. ***Crop Cover Coefficient ( $K_c$ )***

As the crops grow, the amount of water they lose due to transpiration changes. Each crop has a pair of polynomial equations that provide the crop coefficient for any day during the crop's growing season. It was found that two curves, one for modelling time periods up to the present and one for extending the modelling into the future, provided a better sequence of crop coefficients than using a single curve for all years (currently 1961 to 2100). The application automatically selects the current or future curve as modelling moves across the *crop Curve Changeover Year*.

For alfalfa crops, there are different sets of equations corresponding to different cuttings throughout the growing season.

### 4. ***Crop Evapotranspiration ( $ET_c$ )***

The evapotranspiration for each crop is calculated as the general  $ET_o$  multiplied by the crop coefficient ( $K_c$ ):

$$ET_c = ET_o \times K_c$$

### 5. ***Climate Moisture Deficit (CMD)***

During the growing season, the daily Climate Moisture Deficit (CMD) is calculated as the crop evapotranspiration ( $ET_c$ ) less the Effective Precipitation (EP):

$$CMD = ET_c - EP$$

During each crop's growing season, a stored moisture reservoir methodology is used that is similar to the soil moisture content calculation in the pre-season. On a daily basis, the stored moisture level is used towards satisfying the climate moisture deficit to produce an *adjusted Climate Moisture Deficit* ( $CMD_a$ ):

$$CMD_a = CMD - \text{storedMoisture}$$

If the storedMoisture level exceeds the day's CMD, then the  $CMD_a$  is 0 and the stored moisture level is reduced by the CMD amount. If the CMD is greater than the stored moisture, then all of the stored moisture is used (storedMoisture is set to 0) and the adjusted CMD creates an irrigation requirement.

The upper limit for the storedMoisture level during the growing season is the maximum soil water deficit (MSWD) setting.

#### 6. ***Crop Water Requirement (CWR)***

The Crop Water Requirement is calculated as the adjusted Climate Moisture Deficit (CMD<sub>a</sub>) multiplied by the soil water factor (swFactor) and any stress factor (used primarily for grass crops):

$$CWR = CMD_a \times swFactor \times stressFactor$$

#### 7. ***Irrigation Requirement (IR)***

The Irrigation Requirement is the Crop Water Requirement (CWR) after taking into account the irrigation efficiency (I<sub>e</sub>) and, for drip systems, the drip factor (D<sub>f</sub>):

$$IR = CWR \times \frac{D_f}{I_e}$$

For irrigation systems other than drip, the drip factor is 1.

#### 8. ***The Irrigation Water Demand (IWD<sub>perc</sub> and IWD)***

The portion of the Irrigation Water Demand lost to deep percolation is the Irrigation Requirement (IR) multiplied by the percolation factor (soilPercFactor):

$$IWD_{perc} = IR \times soilPercFactor$$

The final Irrigation Water Demand (IWD) is then the Irrigation Requirement (IR) plus the loss to percolation (IWD<sub>perc</sub>):

$$IWD = IR + IWD_{perc}$$

#### 9. ***Frost Protection***

For some crops (e.g. cranberries), an application of water is often used under certain climatic conditions to provide protection against frost damage. For cranberries, the rule is: when the temperature drops to 0 °C or below between March 16 and May 20 or between October 1 and November 15, a frost event will be calculated. The calculated value is an application of 2.5 mm per hour for 10 hours. In addition, 60% of the water is recirculated and reused, accounting for evaporation and seepage losses.

This amounts to a modelled water demand of 10 mm over the cranberry crop's area for each day that a frost event occurs between the specified dates.

#### 10. ***Annual Soil Moisture Deficit***

Prior to each crop's growing season, the Model calculates the soil's moisture content by starting it at full (maximum soil water deficit level) on January 1, and adjusting it daily according to

precipitation and evaporation. During the growing season, simple evaporation is replaced by the crop's evapotranspiration as it progresses through its growth stages. At the completion of each crop's growing season, an annual soil moisture deficit (SMD) is calculated as the difference between the soil moisture content at that point and the maximum soil water deficit (MSWD):

$$\text{SMD} = \text{MSWD} - \text{storedMoisture}$$

In dry/cold climates, this amount represents water that the farmer would add to the soil in order to prevent it from freezing. Wet climates are assumed to have sufficient precipitation and warm enough temperatures to avoid the risk of freezing without this extra application of water; the SMD demand is therefore recorded only for dry areas.

There is no fixed date associated with irrigation to compensate for the annual soil moisture deficit. The farmer may choose to do it any time after the end of the growing season and before the freeze up. In the Model's summary reports, the water demand associated with the annual soil moisture deficit shows as occurring at time 0 (week 0, month 0, etc.) simply to differentiate it from other demands that do have a date of occurrence during the crop's growing season.

Greenhouses and mushroom barns do not have an annual soil moisture deficit.

## **11. *Flood Harvesting***

Cranberry crops are generally harvested using flood techniques. The Model calculates the flood harvesting demand as 250 mm of depth for 10% of the cranberry farmed area. For modelling purposes, it is assumed that 250 mm of water gets applied to the total cranberry crop area, 10% at a time. The water is reused for subsequent portions, but by the time the entire crop is harvested, all of the water is assumed to have been used and either depleted through losses or released from the farm.

The water demand is therefore calculated as a fixed 25 mm over the entire cranberry crop area. The harvesting generally takes place between mid-October and mid-November where the Model treats it as occurring on the fixed date of November 16.

## Livestock Water Use

The Model calculates an estimated livestock water demand using agricultural census data and an estimate of the water use per animal. Water use for each animal type is calculated a bit differently depending on requirements. For example, for a dairy milking cow, the water demand for each animal includes, drinking, preparation for milking, pen and barn cleaning, milking system washout, bulk tank washout and milking parlor washing. However, for a dry dairy cow, the demand only includes drinking and pen and barn cleaning.

The water use is estimated on a daily basis per animal even though the facility is not cleaned daily. For example, for a broiler operation, the water use for cleaning a barn is calculated as 4 hours of pressure washing per cycle at a 10 gpm flow rate, multiplied by 6 cycles per barn with each barn holding 50,000 birds. On a daily basis, this is quite small with a value of 0.01 litres per day per bird applied.

For all cases, the daily livestock demand is applied to the farm location. However, in the case of beef, the livestock spend quite a bit of the year on the range. Since the actual location of the animals cannot be ascertained, the water demand is applied to the home farm location, even though most of the demand will not be from this location. Therefore, the animal water demand on a watershed scale will work fine but not when the demand is segregated into sub-watersheds or groundwater areas.

The estimates used for each livestock are shown in Table 1.

<b>Table 1      Livestock Water Demand (Litres/day)</b>				
<b>Animal Type</b>	<b>Drinking</b>	<b>Milking Preparation</b>	<b>Barn Component</b>	<b>Total</b>
Milking Dairy Cow	65	5	15	85
Dry Cow	45		5	50
Swine	12		0.5	12.5
Poultry – Broiler	0.16		0.01	0.17
Poultry – Layer	0.08		0.01	0.09
Turkeys	0.35		0.01	0.36
Goats	8			8
Sheep	8			8
Beef – range, steer, bull, heifer	50			50
Horses	50			50



# Definition and Calculation of Individual Terms used in the Irrigation Water Demand Equation

---

## Growing Season Boundaries

There are three sets of considerations used in calculating the start and end of the irrigation season for each crop:

- temperature-based growing season derivations, generally using Temperature Sum (Tsum) or Growing Degree Day (GDD) accumulations
- the growing season overrides table
- the irrigation season overrides table

These form an order of precedence with later considerations potentially overriding the dates established for the previous rules. For example, the temperature-based rules might yield a growing season start date of day 90 for a given crop in a mild year. To avoid unrealistic irrigation starts, the season overrides table might enforce a minimum start day of 100 for that crop; at that point, the season start would be set to day 100. At the same time, a Water Purveyor might not turn on the water supply until day 105; specifying that as the minimum start day in the irrigation season overrides table would prevent any irrigation water demands until day 105.

This section describes the rules used to establish growing season boundaries based on the internal calculations of the Model. The GDD and Tsum Day calculations are described in separate sections. The *standard end of season* specified for several crops is the earlier of the end date of Growing Degree Day with base temperature of 5 °C (GDD<sub>5</sub>) or the first frost.

### 1. *Corn (silage corn)*

- uses the corn\_start date for the season start
- season end: earlier of the killing frost or the day that the CHU2700 (2700 Corn Heat Units) threshold is reached

### 2. *Sweetcorn, Potato, Tomato, Pepper, Strawberry, Vegetable, Pea*

- corn\_start date for the season start
- corn start plus 110 days for the season end

### 3. *Cereal*

- GDD<sub>5</sub> start for the season start
- GDD<sub>5</sub> start plus 130 days for the season end

### 4. *AppleHD, AppleMD, AppleLD, Asparagus, Berry, Blueberry, Ginseng, Nuts, Raspberry, Sourcherry, Treefruit, Vineberry*

- season start:  $(0.8447 \times \text{tsum600\_day}) + 18.877$
- standard end of season

### 5. *Pumpkin*

- corn\_start date
- standard end of season

6. ***Apricot***
  - season start:  $(0.9153 \times \text{tsum400\_day}) + 5.5809$
  - standard end of season
7. ***CherryHD, CherryMD, CherryLD***
  - season start:  $(0.7992 \times \text{tsum450\_day}) + 24.878$
  - standard end of season
8. ***Grape, Kiwi***
  - season start:  $(0.7992 \times \text{tsum450\_day}) + 24.878$
  - standard end of season
9. ***Peach, Nectarine***
  - season start:  $(0.8438 \times \text{tsum450\_day}) + 19.68$
  - standard end of season
10. ***Plum***
  - season start:  $(0.7982 \times \text{tsum500\_day}) + 25.417$
  - standard end of season
11. ***Pear***
  - season start:  $(0.8249 \times \text{tsum600\_day}) + 17.14$
  - standard end of season
12. ***Golf, TurfFarm***
  - season start: later of the GDD<sub>5</sub> start and the tsum300\_day
  - standard end of season
13. ***Domestic, Yard, TurfPark***
  - season start: later of the GDD<sub>5</sub> start and the tsum400\_day
  - standard end of season
14. ***Greenhouse (interior greenhouses)***
  - fixed season of April 1 – October 30
15. ***GH Tomato, GH Pepper, GH Cucumber***
  - fixed season of January 15 – November 30
16. ***GH Flower***
  - fixed season of March 1 – October 30
17. ***GH Nursery***
  - fixed season of April 1 – October 30
18. ***Mushroom***
  - all year: January 1 – December 31

**19. *Shrubs/Trees, Fstock, NurseryPOT***

- season start: tsum500\_day
- end: Julian day 275

**20. *Floriculture***

- season start: tsum500\_day
- end: Julian day 225

**21. *Cranberry***

- season start: tsum500\_day
- end: Julian day 275

**22. *Grass, Forage, Alfalfa, Pasture***

- season start: later of the GDD<sub>5</sub> and the tsum600\_day
- standard end of season

**23. *Nursery***

- season start: tsum400\_day
- standard end of season

## **Evapotranspiration (ET<sub>o</sub>)**

The ET<sub>o</sub> calculation follows the FAO Penman-Montieth equation. Two modifications were made to the equation:

- Step 6 – Inverse Relative Distance Earth-Sun (d<sub>r</sub>)  
Instead of a fixed 365 days as a divisor, the actual number of days for each year (365 or 366) was used.
- Step 19 – Evapotranspiration (ET<sub>o</sub>)  
For consistency, a temperature conversion factor of 273.16 was used instead of the rounded 273 listed.

## **Availability Coefficient (AC)**

The availability coefficient is a factor representing the percentage of the soil's total water storage that the crop can readily extract. The factor is taken directly from the crop factors table (crop\_factors) based on the cropId value.

## **Rooting Depth (RD)**

The rooting depth represents the crop's maximum rooting depth and thus the depth of soil over which the plant interacts with the soil in terms of moisture extraction. The value is read directly from the crop factors table.

### Stress Factor (stressFactor)

Some crops, such as *grasses*, are often irrigated to a less degree than their full theoretical requirement for optimal growth. The *stress factor (crop\_groups\_and\_factors)* reduces the calculated demand for these crops.

### Available Water Storage Capacity (AWSC)

The available water storage capacity is a factor representing the amount of water that a particular soil texture can hold without the water dropping through and being lost to deep percolation. The factor is taken directly from the soil factors table (*soil\_factors*).

### Maximum Soil Water Deficit (MSWD)

The maximum soil water deficit is the product of the crop's availability coefficient, rooting depth, and the available water storage capacity of the soil:

$$MSWD = RD \times AWSC \times AC$$

### Deep Percolation Factor (Soilpercfactor)

The soil percolation factor is used to calculate the amount of water lost to deep percolation under different management practices.

For greenhouse crops, the *greenhouse leaching factor* is used as the basic soil percolation factor. This is then multiplied by a greenhouse recirculation factor, if present, to reflect the percentage of water re-captured and re-used in greenhouse operations.

$$soilPercFactor = soilPercFactor \times (1 - recirculationFactor)$$

For Nursery Pot (Nursery POT) and Forestry Stock (Fstock) crops, the soil percolation factor is fixed at 35%. For other crops, the factor depends on the soil texture, the MSWD, the irrigation system, and the Irrigation Management Practices code. The percolation factors table (*soil\_percolation\_factors*) is read to find the first row with the correct management practices, soil texture and irrigation system, and a MSWD value that matches or exceeds the value calculated for the current land use polygon.

If the calculated MSWD value is greater than the index value for all rows in the percolation factors table, then the highest MSWD factor is used. If there is no match based on the passed parameters, then a default value of 0.25 is applied.

For example, a calculated MSWD value of 82.5 mm, a soil texture of sandy loam (SL) and an irrigation system of solid set overtree (Ssovertime) would retrieve the percolation factor associated with the MSWD index value of 75 mm in the current table (presently, there are rows for MSWD 50 mm and 75 mm for SL and Ssovertime).



### Maximum Evaporation Factor (maxEvaporation)

Just as different soil textures can hold different amounts of water, they also have different depths that can be affected by evaporation. The factor is taken directly from the soil factors table.

### Irrigation Efficiency ( $I_e$ )

Each irrigation system type has an associated efficiency factor (inefficient systems require the application of more water in order to satisfy the same crop water demand). The factor is read directly from the irrigation factors table (*irrigation\_factors*).

### Soil Water Factor (swFactor)

For the greenhouse “crop”, the soil water factor is set to 1. For other crops, it is interpolated from a table (*soil\_water\_factors*) based on the MSWD. For Nurseries, the highest soil water factor (lowest MSWD index) in the table is used; otherwise, the two rows whose MSWD values bound the calculated MSWD are located and a soil water factor interpolated according to where the passed MSDW value lies between those bounds.

For example, using the current table with rows giving soil water factors of 0.95 and 0.9 for MSWD index values of 75 mm and 100 mm respectively, a calculated MSWD value of 82.5 mm would return a soil water factor of:

$$0.95 + \left[ \frac{82.5 - 75}{100 - 75} \times (0.9 - 0.95) \right] \\ = 0.935$$

If the calculated MSWD value is higher or lower than the index values for all of the rows in the table, then the factor associated with the highest or lowest MSWD index is used.

### Early Season Evaporation Factor (earlyEvaporationFactor)

The effective precipitation (precipitation that adds to the stored soil moisture content) can be different in the cooler pre-season than in the growing season. The early season evaporation factor is used to determine what percentage of the precipitation is considered effective prior to the growing season.

### Crop Coefficient ( $K_c$ )

The crop coefficient is calculated from a set of fourth degree polynomial equations representing the crop’s ground coverage throughout its growing season. The coefficients for each term are read from the crop factors table based on the crop type, with the variable equalling the number of days since the start of the crop’s growing season. For example, the crop coefficient for Grape on day 35 of the growing season would be calculated as:

$$K_c = [0.0000000031 \times (35)^4] + [-0.0000013775 \times (35)^3] + (0.0001634536 \times (35)^2) + (-0.0011179845 \times 35) + 0.2399004137 \\ = 0.346593241$$

Alfalfa crops have an additional consideration. More than one cutting of alfalfa can be harvested over the course of the growing season, and the terms used for the crop coefficient equation changes for the different cuttings. For alfalfa, the alfalfa cuttings table is first used to determine which cutting period the day belongs to (first, intermediate or last), and after that the associated record in the crop factors table is accessed to determine the terms.

There are two sets of polynomial coefficients used to calculate the crop coefficient; the first set is used for modelling time periods up to the year specified as the *crop curve changeover year*; and the second for modelling into the future. The changeover year will be modified as time goes on and new historical climate observations become available.

## Growing Degree Days (GDD)

The Growing Degree Day calculations generate the start and end of GDD accumulation.

### 1. *Start of GDD Accumulation*

For each base temperature (bases 5 and 10 are always calculated, other base temperature can be derived), the start of the accumulation is defined as occurring after 5 consecutive days of  $T_{\text{mean}}$  matching or exceeding the base temperature (BaseT). The search for the start day gets reset if a killing frost ( $< -2^{\circ}\text{C}$ ) occurs, even after the accumulation has started. The search also restarts if there are 2 or more consecutive days of  $T_{\text{min}} \leq 0^{\circ}\text{C}$ . The GDD start is limited to Julian days 1 to 210; if the accumulation has not started by that point, then it is unlikely to produce a reasonable starting point for any crop.

### 2. *End of GDD accumulation*

The search for the end of the GDD accumulation begins 50 days after its start. The accumulation ends on the earlier of 5 consecutive days where  $T_{\text{mean}}$  fails to reach BaseT (strictly *less than*) or the first killing frost ( $-2^{\circ}\text{C}$ ).

During the GDD accumulation period, the daily contribution is the difference between  $T_{\text{mean}}$  and BaseT, as long as  $T_{\text{mean}}$  is not less than BaseT:

$$\text{GDD} = T_{\text{mean}} - \text{BaseT}; 0 \text{ if negative}$$

## Frost Indices

Three frost indices are tracked for each year:

- the last spring frost is the latest day in the first 180 days of the year with a  $T_{\text{min}} \leq 0^{\circ}\text{C}$
- the first fall frost is the first day between days 240 and the end of the year where  $T_{\text{min}} \leq 0^{\circ}\text{C}$
- the killing frost is the first day on or after the first fall frost where  $T_{\text{min}} \leq -2^{\circ}\text{C}$

## Corn Heat Units (CHU)

The Corn Heat Unit is the average of two terms using  $T_{\text{min}}$  and  $T_{\text{max}}$ . Prior to averaging, each term is set to 0 individually if it is negative.

$$\begin{aligned} \text{term1} &= [3.33 \times (T_{\text{max}} - 10)] - [0.084 \times (T_{\text{max}} - 10) \times (T_{\text{max}} - 10)]; 0 \text{ if negative} \\ \text{term2} &= 1.8 \times (T_{\text{min}} - 4.44); 0 \text{ if negative} \\ \text{CHU} &= \frac{(\text{term1} + \text{term2})}{2} \end{aligned}$$

## Corn Season Start and End

The corn season boundary derivations are similar to the GDD determinations. The start day is established by 3 consecutive days where  $T_{\text{mean}} \geq 11.2$  °C. As in the case of the GDD calculations, the search for the corn season start day gets reset if  $T_{\text{min}} \leq -2$  °C, or if there are 2 or more consecutive days of  $-2$  °C  $\leq T_{\text{min}} \leq 0$  °C.

The search for the silage corn season end begins 50 days after the start. The season ends on the earlier of a mean temperature dropping below 10.1 or a killing frost.

The end of the sweet corn season is defined as 110 days after the season start.

## Tsum Indices

The Tsum day for a given number is defined as the day that the sum of the positive daily  $T_{\text{mean}}$  reaches that number. For example, the Tsum400 day is the day where the sum of the positive  $T_{\text{mean}}$  starting on January 1 sum to 400 units or greater.

Days where  $T_{\text{mean}}$  falls below 0 °C are simply not counted; therefore, the Model does not restart the accumulation sequence.

## Wet/Dry Climate Assessment

Starting with the Lower Mainland, some of the modelling calculations depend on an assessment of the general climatic environment as *wet* or *dry*. For example, when modelling the soil moisture content prior to the start of the crop's growing season, the reservoir can only be drawn down by evaporation except for *grass* crops in *wet* climates which can pull additional moisture out of the soil.

The assessment of wet or dry uses the total precipitation between May 1 and September 30. If the total is more than 125 mm during that period, the climate is considered to be *wet* and otherwise *dry*.

## Groundwater Use


The Model generates water sources for irrigation systems. This is done by first determining which farms are supplied by a water purveyor, and then coding those farms as such. Most water purveyors use surface water but where groundwater is used, the farms are coded as groundwater use. The second step is to check all water licences and assign the water licences to properties in the database. The remaining farms that are irrigating will therefore not have a water licence or be supplied by a water purveyor. The assumption is made that these farms are irrigated by groundwater sources.

## Land Use Results


A summary of the land area and the inventoried area of the Similkameen Watershed are shown in Table 2. The primary agricultural use of the ARL area is shown in Table 3. Figures 8, 9 and 10 show the areas of water, ALR land and land parcels in the basin graphically. Figure 11 provides a schematic of the higher yielding aquifer areas in the Similkameen Watershed.

<b>Table 2 Overview of Similkameen Basin's Land and Inventoried Area</b>		
<b>Area Type</b>	<b>Area (ha)</b>	<b>Number of Parcels</b>
<b>Similkameen Watershed</b>		
Total Area	757,157	-
Area of Water Feature	4,374	-
Area of Land (excluding water features)	752,783	-
ALR Area	62,108	2,585
Area of First Nations Reserve	17,344	71
<b>Inventoried Area</b>		
Total Inventoried Area	62,649	2,599
Area of First Nations Reserve in ALR	4,903	63


<b>Table 3 Summary of Primary Agricultural Activities of Inventoried Parcels where Primary Land Use is agriculture in Similkameen</b>		
<b>Primary Agriculture Activity</b>	<b>Total ALR Area (ha)</b>	<b>Number of Parcels</b>
Polyhouse	2	5
Floriculture	1	2
Forage and pasture	4,637	473
Grains, cereals and oilseeds	14	3
Nursery/trees	6	4
Specialty crops	4	6
Tree fruit	673	305
Vegetables	72	55
Vine and berry	231	60
Other	324	95
<b>Total</b>	<b>5,966</b>	<b>1,012</b>



**Figure 8 Water Areas in the Similkameen Watershed**




**Figure 9 ALR Areas in the Similkameen Watershed**



**Figure 10 Land Parcels in the Similkameen Watershed**





**Figure 11 Higher Productive Groundwater Aquifers in the Similkameen Watershed**

## ***Agricultural Water Demand Results***

---

The Model has a reporting feature that can save and generate reports for many different scenarios that have been pre-developed. This report will provide a summary of the reported data in the Appendices. Climate data from 1997 and 2003 were chosen as they represent a relatively wet year and dry year respectively. Most reports are based on the 2003 data since the maximum current demand can then be presented.

### **Annual Crop Water Demand – Tables A and B**

The Model can use three different irrigation management factors, good, average and poor. Unless otherwise noted, average management were used in the tables. Appendix Table A provides the annual irrigation water demand for current crop and irrigation systems used for the year 2003 using average irrigation management, and Table B provides the same data for 1997.

Where a crop was not established, the acreage was assigned a forage crop so that the Model could determine a water demand. The total irrigated acreage in the Similkameen Watershed is 4,533 hectares (ha), including 3,064 ha (70%) for forage crops (alfalfa, grass and corn), and 1,148 ha (26%) for fruit crops. In the Similkameen, 2,499 ha (55%) is supplied by licensed surface water sources, and 2,033 ha (45%) is irrigated with groundwater.

The total annual irrigation demand was 39,178,239 m<sup>3</sup> in 2003, and dropped to 21,898,136 m<sup>3</sup> in 1997. During a wet year like 1997, the demand was only 55% of a hot dry year like 2003. There were 352.7 ha of land surveyed that had an irrigation system but were deemed to be not in use at the time of the survey.

In addition, the Model also calculates demand based on relatively good practices. As such, actual use may actually be higher or lower than what is calculated by the Model.

### **Annual Water Demand by Irrigation System – Table C**

The crop irrigation demand can also be reported by irrigation system type as shown in Table C. The total area that is currently irrigated by efficient drip systems is 248 ha (6%). Low-pressure pivot irrigation systems irrigate 275 ha (6%). Sprinkler systems (including solid set overtree and undertree), guns, wheelline and handline, irrigate 3,076 ha (70%) of the agricultural crops in Similkameen.

### **Annual Water Demand by Soil Texture – Table D**

Table D provides the annual water demand by soil texture. Where soil texture data is missing, the soil texture has been defaulted to sandy loam. All outdoor domestic irrigated areas were also assumed to use sandy loam. The defaults are shown in Table D.

## Annual Water Demand by Water Purveyor – Table E

The Model calculates water demand on a property-by-property basis, and can summarize the data for each water purveyor in the Similkameen Watershed. Table E provides the water demands for the area supplied by a water purveyor, First Nations lands that are irrigated (457 ha or 10%), and privately irrigated areas (3,038 ha or 66%).

## Irrigated Area within Local Governments – Table F

Table F provides a breakdown of the agriculture irrigated areas within the boundaries of each local government within the Similkameen watershed.

## Irrigation Management Factors – Table G

The Model can estimate water demand based on poor, average and good irrigation management factors. This is accomplished by developing an irrigation management factor for each crop, soil and irrigation system combination. The Maximum Soil Water Deficit (MSWD) is the maximum amount of water that can be stored in the soil within the crop rooting zone. An irrigation system applying more water than what can be stored will result in percolation beyond the crop's rooting depth. Irrigation systems with high application rates will have a probability of higher percolation rates, a stationary gun for instance.

For each soil class, ranges of four MSWD are provided, which reflect a range of crop rooting depths. An irrigation management factor, which determines the amount of leaching, is established for each of the MSWD values for the soil types (Table 4). The management factor is based on irrigation expertise as to how the various irrigation systems are able to operate. For example, Table 5 indicates that for a loam soil and a MSWD of 38 mm, a solid set overtree system has a management factor of 0.10 for good management while the drip system has a management factor of 0.05. This indicates that it is easier to prevent percolation with a drip system than it is with a solid set sprinkler system. For poor management, the factors are higher.

There are a total of 1,344 irrigation management factors established for the 16 different soil textures, MSWD and 21 different irrigation system combinations used in the Model.

Table 4 Irrigation Management Factors							
Soil Texture	MSWD	Solid Set Overtree			Drip		
		Good	Average	Poor	Good	Average	Poor
Loam	38	0.10	0.15	0.20	0.05	0.10	0.15
	50	0.05	0.10	0.15	0.05	0.075	0.10
	75	0.05	0.10	0.15	0.05	0.075	0.10
	100	0.05	0.075	0.10	0.05	0.075	0.10
Sandy loam	25	0.20	0.225	0.25	0.10	0.15	0.20
	38	0.10	0.15	0.20	0.10	0.125	0.15
	50	0.05	0.10	0.15	0.05	0.10	0.10
	75	0.05	0.10	0.15	0.05	0.075	0.10

The management factors increase as the MSWD decreases because there is less soil storage potential in the crop rooting depth. For irrigation systems such as guns, operating on a pasture which has a shallow rooting depth, and on a sandy soil which cannot store much water, the poor irrigation management factor may be as high as 0.50.

The management factor used in the Model assumes all losses are deep percolation while it is likely that some losses will occur as runoff as well. It should be recognized that the values are a bit subjective but they do indicate the potential for improvements.

Table G provides an overview of the impacts on the management factor and irrigation systems used. Management improvements could be more significant if irrigation systems were converted to more efficient systems.

Table G provides percolation rates based on good, average and poor management using 2003 climate data. In summary, there is 3,479,837 m<sup>3</sup> of water lost to percolation on good management, 4,237,588 m<sup>3</sup> on average management, and 4,995,339 m<sup>3</sup> on poor management. Percolation rates for poor management are 43% higher than for good management.

### **Deep Percolation – Table H**

The percolation rates vary by crop, irrigation system type, soil and the management factor used. Table H shows the deep percolation amounts by irrigation system type for average management. The last column provides a good indication of the average percolation per hectare for the various irrigation system types. For example, drip irrigation systems have only 13% of the percolation rates of gun systems. Landscape systems have a high percolation rate predominantly because application rates are high and the crop rooting depth is quite shallow.

### **Improved Irrigation Efficiency and Good Management – Table I**

There is an opportunity to reduce water use by converting irrigation systems to a higher efficiency for some crops. For example, drip systems could be used for all berry crops, vegetable crops and some of the other horticultural crops. Forage crops could use low pressure center pivot systems for all field sizes larger than 10 ha. In addition, using better management such as irrigation scheduling techniques will also reduce water use. Table I provides a scenario of water demand if all sprinkler systems are converted to drip systems for horticultural crops and forage fields larger than 10 ha are converted to pivot systems, using good irrigation management. The water demand for 2003 would then reduce from 39,178,239 m<sup>3</sup> to 34,153,841 m<sup>3</sup>. This is about an 13% reduction in water demand.

### **Livestock Water Use – Table J**

The Model provides an estimate of water use for livestock. The estimate is based on the number of animals in the Similkameen Watershed as determined by the latest census, the drinking water required for each animal per day and the barn or milking parlour wash water. Values used are shown in Table J. For Similkameen, the amount of livestock water is estimated at 303,068 m<sup>3</sup>.

Climate Change Water Demand for 2050 – Table K

The Model also has access to climate change information until the year 2100. While data can be run for each year, three driest years in the 2050’s were selected to give a representation of climate change. Figure 12 shows the climate data results which indicate that 2053, 2056, and 2059 generate the highest annual ET<sub>o</sub> and lowest annual precipitation. These three years were used in this report.





Figure 12 Annual ET and Effective Precipitation in 2050's

Table K provides the results of climate change on irrigation demand for the three years selected using current crops and irrigation systems. Current crops and irrigation systems are used to show the increase due to climate change only, with no other changes taking place.

Figure 13 shows all of the climate change scenario runs for the Okanagan using 12 climate models from 1960 to 2100. This work was compiled by Denise Neilsen at the Agriculture and Agri-Food Canada – Summerland Research Station. There is a lot of scatter in this figure, but it is obvious that there is a trend of increasing water demand.

The three climate change models used in this report are RCP26, RCP45 and RCP85. Running only three climate change models on three selected future years in the Similkameen Watershed is not sufficient to provide a trend like in Figure 13. What the results do show is that in an extreme climate scenario, it is possible to have an annual water demand that is 15% higher than what was experienced in 2003 based on RCP85 climate model in 2053. More runs of the climate change models will be required to better estimate a climate change trend for the region.



**Figure 13 Future Irrigation Demand for All Outdoor Uses in the Okanagan in Response to Observed Climate Data (Actuals) and Future Climate Data Projected from a Range of Global Climate Models**

### **Agricultural Buildout Crop Water Demand Using 2003 Climate Data – Table L**

An agricultural buildout scenario was developed that looked at potential agricultural lands that could be irrigated in the future. The rules used to establish where potential additional agricultural lands were located are as follows:

- within 1,000 m of water supply (lake)
- within 1,000 m of water supply (water course)
- within 1,000 m of water supply (wetland)
- within 1,000 m of high productivity aquifer
- within 1,000 m of water purveyor
- with Ag Capability class 1-4 only where available
- must be within the ALR
- below 800 m average elevation
- remove parcels to the East of Richter Mountain (due to a lack of access to water)
- must be private ownership

For the areas that are determined to be eligible for future buildout, a crop and irrigation system need to be applied. Where a crop already existed in the land use inventory, that crop would remain and an



irrigation system assigned. If no crop existed, then a crop and irrigation system is assigned as per the criteria below:

- **West of Hedley:** 35% grass with sprinkler irrigation, and 35% alfalfa with sprinkler irrigation or low pressure pivot
- **East of Hedley:** 10% grapes with drip irrigation, 10% vegetables with drip irrigation, and 10% apples with drip irrigation

The sum of the percentages is the total of the entire Similkameen Watershed. West of Hedley, the properties where land covers are available would be split 50%-50% for grass-sprinkler and alfalfa-sprinkler. East of Hedley, properties where land covers are available would be split 33%-33%-33% for grape-drip/vegetable-drip/apple-drip.

For irrigated forage crops that are equal to or over 10 ha in size, the irrigation system type will be changed from sprinkler to low-pressure pivot (if not already using a low-pressure pivot). It is anticipated that current irrigation systems will be replaced by more efficient systems like low-pressure pivots in the future to reduce water demand when water resources are more stretched.

Figure 14 indicates the location of agricultural land that is currently irrigated (dark green) and the land that can be potentially irrigated (red). Based on the scenario provided for the Similkameen Watershed, the additional agricultural land that could be irrigated is 1,827 ha. The water demand for a climate year like 2003 would be 46,224,424 m<sup>3</sup> assuming efficient irrigation systems and good management.

### **Agricultural Buildout Crop Water Demand for 2050 – Table M**


The same irrigation expansion and cropping scenario used to generate the values in Table L were used to generate the climate change water demand shown in Table M. See discussion under Table K section. When climate change is added to the buildout scenario, the water demand increases from 45 million m<sup>3</sup> to 59 million m<sup>3</sup> based on climate change model RCP85 in 2059. This demonstrates the extremes that could occur in the future.

### **Irrigation Systems Used for the Buildout Scenario for 2003 Climate Data– Table N**

Table N provides an account of the irrigation systems used by area for the buildout scenario in the previous two examples. Note that low pressure pivot and drip irrigation are the two most predominant system types. This is because the model is applying the most efficient irrigation systems to the future irrigated area.

### **Water Demand for the Buildout Area by Water Purveyor 2003 Climate Data – Table O**

Table O provides the water demand within the current water purveyed areas of the Similkameen Watershed for the buildout scenario used in this report. Comparing these values with the result in Table E will provide information on the possible increased water demand for the areas supplied by water purveyors.



**Figure 14 Similkameen Watershed Irrigation Expansion Potential**

### **Water Demand for the Buildout Area by Local Government 2003 Climate Data – Table P**

Table P provides the future water demand within local government boundaries using previous scenarios. Comparing these values with the result in Table F will provide information on the possible increased water demand within local governments if the buildout scenarios actually occurred in the future.

## Literature

---

Cannon, A.J., and Whitfield, P.H. (2002), Synoptic map classification using recursive partitioning and principle component analysis. *Monthly Weather Rev.* 130:1187-1206.

Cannon, A.J. (2008), Probabilistic multi-site precipitation downscaling by an expanded Bernoulli-gamma density network. *Journal of Hydrometeorology*. <http://dx.doi.org/10.1175%2F2008JHM960.1>  
Intergovernmental Panel on Climate Change (IPCC) (2008), Fourth Assessment Report –AR4. <http://www.ipcc.ch/ipccreports/ar4-syr.htm>

Merritt, W, Alila, Y., Barton, M., Taylor, B., Neilsen, D., and Cohen, S. 2006. Hydrologic response to scenarios of climate change in the Okanagan Basin, British Columbia. *J. Hydrology*. 326: 79-108.

Neilsen, D., Smith, S., Frank, G., Koch, W., Alila, Y., Merritt, W., Taylor, B., Barton, M, Hall, J. and Cohen, S. 2006. Potential impacts of climate change on water availability for crops in the Okanagan Basin, British Columbia. *Can. J. Soil Sci.* 86: 909-924.

Neilsen, D., Duke, G., Taylor, W., Byrne, J.M., and Van der Gulik T.W. (2010). Development and Verification of Daily Gridded Climate Surfaces in the Okanagan Basin of British Columbia. *Canadian Water Resources Journal* 35(2), pp. 131-154. <http://www4.agr.gc.ca/abstract-resume/abstract-resume.htm?lang=eng&id=21183000000448>

Allen, R. G., Pereira, L. S., Raes, D. and Smith, M. 1998. Crop evapotranspiration Guidelines for computing crop water requirements. FAO Irrigation and Drainage Paper 56. United Nations Food and Agriculture Organization. Rome. 100pp

## ***Appendix Tables***

---

<b>Appendix Table A</b>	<b>2003 Water Demand by Crop with Average Management</b>
<b>Appendix Table B</b>	<b>1997 Water Demand by Crop with Average Management</b>
<b>Appendix Table C</b>	<b>2003 Water Demand by Irrigation System with Average Management</b>
<b>Appendix Table D</b>	<b>2003 Water Demand by Soil Texture with Average Management</b>
<b>Appendix Table E</b>	<b>2003 Water Demand by Purveyor with Average Management</b>
<b>Appendix Table F</b>	<b>2003 Water Demand by Local Government with Average Management</b>
<b>Appendix Table G</b>	<b>2003 Management Comparison on Irrigation Demand and Percolation Volumes</b>
<b>Appendix Table H</b>	<b>2003 Percolation Volumes by Irrigation System with Average Management</b>
<b>Appendix Table I</b>	<b>2003 Crop Water Demand for Improved Irrigation System Efficiency and Good Management</b>
<b>Appendix Table J</b>	<b>2003 Water Demand by Animal Type with Average Management</b>
<b>Appendix Table K</b>	<b>Climate Change Water Demand Circa 2050 for a High Demand Year with Good Management using Current Crops and Irrigation Systems</b>
<b>Appendix Table L</b>	<b>Buildout Crop Water Demand for 2003 Climate Data and Good Management</b>
<b>Appendix Table M</b>	<b>Buildout Crop Water Demand for Climate Change Data Circa 2050 and Good Management</b>
<b>Appendix Table N</b>	<b>Buildout Irrigation System Demand for 2003 Climate Data and Good Management</b>
<b>Appendix Table O</b>	<b>Buildout Water Demand by Purveyor for 2003 Climate Data and Good Management</b>
<b>Appendix Table P</b>	<b>Buildout Water Demand by Local Government for 2003 Climate Data and Good Management</b>

**Appendix Table A 2003 Water Demand by Crop with Average Management**

Water Source	Surface Water			Reclaimed Water			Groundwater			Total		
Agriculture Crop Group	Irrigated Area (ha)	Irrigation Demand (m <sup>3</sup> )	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m <sup>3</sup> )	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m <sup>3</sup> )	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m <sup>3</sup> )	Avg. Req. (mm)
Alfalfa	1,526.9	13,692,147	897	-	-	-	645.9	5,660,493	876	2,172.8	19,352,641	891
Apple	140.0	1,131,373	808	-	-	-	479.6	3,952,356	824	619.7	5,083,729	820
Berry	4.2	38,945	931	-	-	-	1.9	11,617	618	6.1	50,562	834
Cherry	9.9	86,143	866	-	-	-	93.5	836,813	895	103.4	922,956	892
Corn	3.9	26,554	677	-	-	-	16.7	142,441	853	20.6	168,994	819
Forage	541.4	5,167,246	954	-	-	-	329.7	3,023,966	917	871.0	8,191,212	940
Fruit	12.7	110,170	865	-	-	-	141.0	1,248,725	886	153.7	1,358,895	884
Golf	-	-	-	-	-	-	29.3	252,078	861	29.3	252,078	861
Grape	124.4	549,717	442	-	-	-	140.7	530,655	377	265.1	1,080,372	407
Greenhouse/Nursery	3.7	39,482	1,061	-	-	-	4.0	104,058	1,046	7.7	143,540	1,059
Recreational Turf	122.0	1,161,655	953	-	-	-	80.6	818,846	1,015	202.6	1,980,501	978
Vegetable	10.4	80,942	781	-	-	-	70.9	511,816	722	81.3	592,758	729
<b>TOTALS</b>	2,499.5	22,084,374	884	-	-	-	2,033.8	17,093,865	840	4,533.3	39,178,239	864

**Appendix Table B 1997 Water Demand by Crop with Average Management**

Water Source	Surface Water			Reclaimed Water			Groundwater			Total		
Agriculture Crop Group	Irrigated Area (ha)	Irrigation Demand (m <sup>3</sup> )	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m <sup>3</sup> )	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m <sup>3</sup> )	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m <sup>3</sup> )	Avg. Req. (mm)
Alfalfa	1,526.9	7,104,331	465	-	-	-	645.9	2,978,894	461	2,172.8	10,083,226	464
Apple	140.0	630,752	450	-	-	-	479.6	2,261,695	472	619.7	2,892,447	467
Berry	4.2	20,989	502	-	-	-	1.9	4,959	264	6.1	25,948	428
Cherry	9.9	49,216	495	-	-	-	93.5	497,927	533	103.4	547,143	529
Corn	3.9	13,639	348	-	-	-	16.7	79,837	478	20.6	93,476	453
Forage	541.4	3,088,608	571	-	-	-	329.7	1,896,849	575	871.0	4,985,457	572
Fruit	12.7	61,808	485	-	-	-	141.0	720,350	511	153.7	782,158	509
Golf	-	-	-	-	-	-	29.3	185,641	634	29.3	185,641	634
Grape	124.4	220,982	178	-	-	-	140.7	207,985	148	265.1	428,967	162
Greenhouse/Nursery	3.7	20,092	794	-	-	-	4.0	91,422	801	7.7	111,515	795
Recreational Turf	122.0	777,679	638	-	-	-	80.6	562,269	697	202.6	1,339,948	661
Vegetable	10.4	56,651	547	-	-	-	70.9	356,560	503	81.3	413,211	508
<b>TOTALS</b>	<b>2,499.5</b>	<b>12,044,747</b>	<b>482</b>	<b>-</b>	<b>-</b>	<b>-</b>	<b>2,033.8</b>	<b>9,844,389</b>	<b>484</b>	<b>4,533.3</b>	<b>21,889,136</b>	<b>483</b>



**Appendix Table C 2003 Water Demand by Irrigation System with Average Management**

Water Source	Surface Water			Reclaimed Water			Groundwater			Total		
Agriculture Irrigation System	Irrigated Area (ha)	Irrigation Demand (m <sup>3</sup> )	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m <sup>3</sup> )	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m <sup>3</sup> )	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m <sup>3</sup> )	Avg. Req. (mm)
Drip	71.9	286,648	399	-	-	-	176.2	798,450	453	248.1	1,085,098	437
Flood	106.5	1,573,349	1,478	-	-	-	10.0	150,374	1,510	116.4	1,723,722	1,480
GolfSprinkler	-	-	-	-	-	-	19.7	168,038	852	19.7	168,038	852
Gun	85.2	1,048,470	1,230	-	-	-	34.8	360,486	1,035	120.1	1,408,956	1,173
Handline	421.0	3,784,115	899	-	-	-	197.3	1,859,005	942	618.3	5,643,120	913
Landscapesprinkler	16.7	145,664	871	-	-	-	49.8	491,380	986	66.5	637,044	957
Microspray	13.5	96,497	714	-	-	-	180.7	1,377,851	762	194.2	1,474,347	759
Microsprinkler	89.8	740,850	825	-	-	-	228.5	1,910,303	836	318.3	2,651,153	833
Pivot	26.9	218,345	812	-	-	-	27.5	264,268	962	54.4	482,613	888
PivotLP	242.8	1,881,409	775	-	-	-	32.3	253,013	784	275.1	2,134,422	776
SDI	-	-	-	-	-	-	35.1	203,454	579	35.1	203,454	579
Sprinkler	433.3	3,729,168	861	-	-	-	350.4	3,095,842	884	783.6	6,825,011	871
Ssovertree	78.3	430,412	550	-	-	-	35.8	245,896	686	114.1	676,308	593
Sssprinkler	0.4	4,246	1,048	-	-	-	5.0	55,471	1,114	5.4	59,717	1,109
Ssundertree	17.1	162,258	952	-	-	-	130.3	1,236,689	949	147.4	1,398,947	949
Travgun	40.2	336,836	838	-	-	-	0.4	3,389	851	40.6	340,225	838
Wheeline	856.0	7,646,109	893	-	-	-	520.0	4,619,955	888	1,376.0	12,266,064	891
<b>TOTALS</b>	<b>2,499.5</b>	<b>22,084,374</b>	<b>884</b>	<b>-</b>	<b>-</b>	<b>-</b>	<b>2,033.8</b>	<b>17,093,865</b>	<b>840</b>	<b>4,533.3</b>	<b>39,178,239</b>	<b>864</b>

**Appendix Table D 2003 Water Demand by Soil Texture with Average Management**

Water Source	Surface Water			Reclaimed Water			Groundwater			Total		
Agriculture Soil Texture	Irrigated Area (ha)	Irrigation Demand (m <sup>3</sup> )	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m <sup>3</sup> )	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m <sup>3</sup> )	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m <sup>3</sup> )	Avg. Req. (mm)
Clayey Sand	73.3	561,079	766	-	-	-	42.5	331,602	781	115.7	892,681	771
Clayey Sandy Loam	78.2	620,977	795	-	-	-	30.4	243,365	800	108.6	864,342	796
Fine Sandy Loam	57.4	505,135	880	-	-	-	0.2	1,429	816	57.6	506,564	880
Loam	553.5	4,439,167	802	-	-	-	423.9	3,143,617	742	977.5	7,582,783	776
Loamy Sand	414.4	4,534,557	1,094	-	-	-	159.4	1,637,924	1,028	573.7	6,172,481	1,076
Organic	73.4	693,615	945	-	-	-	8.9	74,553	835	82.3	768,168	933
Sand	3.0	35,072	1,182	-	-	-	0.1	507	945	3.0	35,579	1,178
Sandy Clay Loam	189.7	1,783,512	940	-	-	-	61.4	443,469	722	251.1	2,226,981	887
Sandy Loam	602.3	5,425,974	901	-	-	-	615.2	5,639,191	917	1,217.5	11,065,166	909
Sandy Loam (defaulted)	66.2	594,398	898	-	-	-	42.0	361,931	861	108.2	956,330	884
Silt Loam	387.0	2,881,962	745	-	-	-	581.7	4,733,900	814	968.7	7,615,862	786
Silty Clay Loam	1.2	8,924	720	-	-	-	68.2	482,375	707	69.5	491,300	707
<b>TOTALS</b>	<b>2,499.5</b>	<b>22,084,374</b>	<b>884</b>	<b>-</b>	<b>-</b>	<b>-</b>	<b>2,033.8</b>	<b>17,093,865</b>	<b>840</b>	<b>4,533.3</b>	<b>39,178,239</b>	<b>864</b>

**Appendix Table E 2003 Water Demand by Purveyor with Average Management**

Water Source	Surface Water			Reclaimed Water			Groundwater			Total		
Agriculture Purveyor	Irrigated Area (ha)	Irrigation Demand (m <sup>3</sup> )	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m <sup>3</sup> )	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m <sup>3</sup> )	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m <sup>3</sup> )	Avg. Req. (mm)
Cawston	-	-	-	-	-	-	193.6	1,644,655	850	193.6	1,644,655	850
Fairview Heights	-	-	-	-	-	-	263.2	2,169,257	824	263.2	2,169,257	824
Keremeos	-	5	932	-	-	-	581.6	4,948,610	851	581.6	4,948,615	851
Lower Similkameen First Nation	378.0	3,198,316	846	-	-	-	20.9	214,183	1,023	398.9	3,412,499	855
Upper Similkameen First Nation	48.9	424,814	870	-	-	-	8.9	79,651	899	57.7	504,465	874
Private	2,072.7	18,461,239	891	-	-	-	965.6	8,037,509	832	3,038.3	26,498,748	872
<b>TOTALS</b>	<b>2,499.5</b>	<b>22,084,374</b>	<b>884</b>	<b>-</b>	<b>-</b>	<b>-</b>	<b>2,033.8</b>	<b>17,093,865</b>	<b>840</b>	<b>4,533.3</b>	<b>39,178,239</b>	<b>864</b>

**Appendix Table F 2003 Water Demand by Local Government with Average Management**

Water Source	Surface Water			Reclaimed Water			Groundwater			Total		
Agriculture Local Government	Irrigated Area (ha)	Irrigation Demand (m <sup>3</sup> )	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m <sup>3</sup> )	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m <sup>3</sup> )	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m <sup>3</sup> )	Avg. Req. (mm)
Keremeos	-	-	-	-	-	-	28.5	243,276	852	28.5	243,276	852
Lower Similkameen First Nation	378.0	3,198,316	846	-	-	-	20.9	214,183	1,023	398.9	3,412,499	855
Okanagan-Similkameen	2,072.7	18,461,244	891	-	-	-	1,892.0	15,817,026	836	3,964.6	34,278,270	865
Thompson-Nicola	-	-	-	-	-	-	3.1	25,858	840	3.1	25,858	840
Upper Similkameen First Nation	-	-	-	-	-	-	23.7	186,984	789	23.7	186,984	789
Others	48.9	424,814	870	-	-	-	8.9	79,651	899	57.7	504,465	874
<b>TOTALS</b>	<b>2,499.5</b>	<b>22,084,374</b>	<b>884</b>	<b>-</b>	<b>-</b>	<b>-</b>	<b>2,033.8</b>	<b>17,093,865</b>	<b>840</b>	<b>4,533.3</b>	<b>39,178,239</b>	<b>864</b>

**Appendix Table G 2003 Management Comparison on Irrigation Demand and Percolation Volumes**

Water Source	Surface Water				Reclaimed Water				Groundwater				Total				
Agriculture Management	Irrigated Area (ha)	Irrigation Demand (m <sup>3</sup> )	Avg. Req. (mm)	Deep Percolation (m <sup>3</sup> )	Irrigated Area (ha)	Irrigation Demand (m <sup>3</sup> )	Avg. Req. (mm)	Deep Percolation (m <sup>3</sup> )	Irrigated Area (ha)	Irrigation Demand (m <sup>3</sup> )	Avg. Req. (mm)	Deep Percolation (m <sup>3</sup> )	Irrigated Area (ha)	Irrigation Demand (m <sup>3</sup> )	Avg. Req. (mm)	Deep Percolation (m <sup>3</sup> )	Percolation (m <sup>3</sup> /ha)
Poor	2,499.5	22,459,861	899	2,897,430	-	-	-	-	2,033.8	17,476,129	859	2,097,910	4,533.3	39,935,990	881	4,995,339	1,102
Avg	2,499.5	22,084,374	884	2,521,942	-	-	-	-	2,033.8	17,093,865	840	1,715,646	4,533.3	39,178,239	864	4,237,588	935
Good	2,499.5	21,708,887	869	2,146,455	-	-	-	-	2,033.8	16,711,601	822	1,333,382	4,533.3	38,420,488	848	3,479,837	768

**Appendix Table H 2003 Percolation Volumes by Irrigation System with Average Management**

Water Source	Surface Water			Reclaimed Water			Groundwater			Total			
Agriculture Irrigation System	Irrigated Area (ha)	Irrigation Demand (m <sup>3</sup> )	Deep Percolation (m <sup>3</sup> )	Irrigated Area (ha)	Irrigation Demand (m <sup>3</sup> )	Deep Percolation (m <sup>3</sup> )	Irrigated Area (ha)	Irrigation Demand (m <sup>3</sup> )	Deep Percolation (m <sup>3</sup> )	Irrigated Area (ha)	Irrigation Demand (m <sup>3</sup> )	Deep Percolation (m <sup>3</sup> )	Percolation (m <sup>3</sup> /ha)
Drip	71.9	286,648	17,145	-	-	-	176.2	798,450	51,388	248.1	1,085,098	68,533	276
Flood	106.5	1,573,349	398,699	-	-	-	10.0	150,374	38,623	116.4	1,723,722	437,321	3,757
GolfSprinkler	-	-	-	-	-	-	19.7	168,038	26,012	19.7	168,038	26,012	1,320
Gun	85.2	1,048,470	197,150	-	-	-	34.8	360,486	62,162	120.1	1,408,956	259,312	2,159
Handline	421.0	3,784,115	402,772	-	-	-	197.3	1,859,005	236,244	618.3	5,643,120	639,016	1,034
Landscapesprinkler	16.7	145,664	24,599	-	-	-	49.8	491,380	83,021	66.5	637,044	107,619	1,618
Microspray	13.5	96,497	8,077	-	-	-	180.7	1,377,851	105,728	194.2	1,474,347	113,805	586
Microsprinkler	89.8	740,850	55,101	-	-	-	228.5	1,910,303	136,521	318.3	2,651,153	191,622	602
Pivot	26.9	218,345	9,791	-	-	-	27.5	264,268	20,645	54.4	482,613	30,436	559
PivotLP	242.8	1,881,409	133,732	-	-	-	32.3	253,013	15,396	275.1	2,134,422	149,128	542
SDI	-	-	-	-	-	-	35.1	203,454	14,423	35.1	203,454	14,423	411
Sprinkler	433.3	3,729,168	448,164	-	-	-	350.4	3,095,842	382,726	783.6	6,825,011	830,889	1,060
Ssovertime	78.3	430,412	27,118	-	-	-	35.8	245,896	21,036	114.1	676,308	48,155	422
Sssprinkler	0.4	4,246	368	-	-	-	5.0	55,471	7,388	5.4	59,717	7,756	1,436
Ssundertree	17.1	162,258	15,124	-	-	-	130.3	1,236,689	105,249	147.4	1,398,947	120,373	817
Travgun	40.2	336,836	31,034	-	-	-	0.4	3,389	312	40.6	340,225	31,347	772
Wheelline	856.0	7,646,109	753,070	-	-	-	520.0	4,619,955	408,772	1,376.0	12,266,064	1,161,842	844
<b>TOTALS</b>	<b>2,499.5</b>	<b>22,084,374</b>	<b>2,521,942</b>	<b>-</b>	<b>-</b>	<b>-</b>	<b>2,033.8</b>	<b>17,093,865</b>	<b>1,715,646</b>	<b>4,533.3</b>	<b>39,178,239</b>	<b>4,237,588</b>	<b>935</b>

**Appendix Table I 2003 Crop Water Demand for Improved Irrigation System Efficiency and Good Management**

Water Source	Surface Water			Reclaimed Water			Groundwater			Total		
Agriculture Crop Group	Irrigated Area (ha)	Irrigation Demand (m <sup>3</sup> )	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m <sup>3</sup> )	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m <sup>3</sup> )	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m <sup>3</sup> )	Avg. Req. (mm)
Alfalfa	1,526.9	12,449,787	815	-	-	-	645.9	5,463,588	846	2,172.8	17,913,375	824
Apple	140.0	834,420	596	-	-	-	479.6	2,872,457	599	619.7	3,706,877	598
Berry	4.2	23,432	560	-	-	-	1.9	7,472	398	6.1	30,904	510
Cherry	9.9	59,223	595	-	-	-	93.5	582,164	623	103.4	641,386	620
Corn	3.9	26,151	667	-	-	-	16.7	138,615	830	20.6	164,766	799
Forage	541.4	4,410,774	815	-	-	-	329.7	2,785,921	845	871.0	7,196,695	826
Fruit	12.7	78,293	615	-	-	-	141.0	871,848	618	153.7	950,141	618
Golf	-	-	-	-	-	-	29.3	248,480	849	29.3	248,480	849
Grape	124.4	374,827	301	-	-	-	140.7	433,595	308	265.1	808,422	305
Greenhouse/Nursery	3.7	38,714	1,050	-	-	-	4.0	103,968	1,036	7.7	142,682	1,049
Recreational Turf	122.0	1,140,215	935	-	-	-	80.6	809,343	1,004	202.6	1,949,558	962
Vegetable	10.4	50,970	492	-	-	-	70.9	349,585	493	81.3	400,556	493
<b>TOTALS</b>	<b>2,499.5</b>	<b>19,486,807</b>	<b>780</b>	<b>-</b>	<b>-</b>	<b>-</b>	<b>2,033.8</b>	<b>14,667,035</b>	<b>721</b>	<b>4,533.3</b>	<b>34,153,841</b>	<b>753</b>

Appendix Table J 2003 Water Demand by Animal Type	
Animal Type	Demand (m <sup>3</sup> )
Beef	281,079
Dairy - dry	260
Dairy - milking	442
Goats	574
Horses	16,321
Poultry - broiler	878
Poultry - laying	465
Sheep	2,587
Swine	460
<b>TOTALS</b>	<b>303,066</b>

Appendix Table K Climate Change Water Demand Circa 2050 for High Demand Year with Good Management Using Current Crops and Irrigation Systems												
Climate Change	rcp26			rcp45			rcp85			Total		
Year	Irrigated Area (ha)	Irrigation Demand (m <sup>3</sup> )	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m <sup>3</sup> )	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m <sup>3</sup> )	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m <sup>3</sup> )	Avg. Req. (mm)
2053	4,533.3	38,957,339	859	4,533.3	35,576,098	785	4,533.3	45,126,622	995	4,533.3	39,886,686	880
2056	4,533.3	36,114,922	797	4,533.3	37,688,940	831	4,533.3	29,069,328	641	4,533.3	34,291,063	756
2059	4,533.3	21,829,606	482	4,533.3	45,384,375	1,001	4,533.3	45,777,296	1,010	4,533.3	37,663,759	831



**Appendix Table L Buildout Crop Water Demand for 2003 Climate Data with Good Management**

Water Source	Surface Water			Reclaimed Water			Groundwater			Total		
Agriculture Crop Group	Irrigated Area (ha)	Irrigation Demand (m <sup>3</sup> )	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m <sup>3</sup> )	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m <sup>3</sup> )	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m <sup>3</sup> )	Avg. Req. (mm)
Alfalfa	1,941.9	15,623,145	805	-	-	-	747.5	6,387,266	855	2,689.3	22,010,410	818
Apple	220.4	1,276,401	579	-	-	-	594.1	3,483,014	586	814.5	4,759,415	584
Berry	4.2	23,432	560	-	-	-	2.5	9,874	388	6.7	33,306	495
Cherry	9.9	59,223	595	-	-	-	93.8	584,633	623	103.8	643,856	620
Corn	3.9	26,151	667	-	-	-	16.7	138,615	830	20.6	164,766	799
Forage	1,051.3	8,295,085	789	-	-	-	478.6	4,001,959	836	1,529.9	12,297,044	804
Fruit	12.7	78,293	615	-	-	-	141.7	876,291	618	154.5	954,584	618
Golf	-	-	-	-	-	-	29.3	248,480	849	29.3	248,480	849
Grape	197.6	591,024	299	-	-	-	223.6	665,965	298	421.2	1,256,989	298
Greenhouse	0.1	2,969	2,165	-	-	-	3.6	99,824	2,798	3.7	102,793	2,774
Greenhouse	0.1	2,969	1,100	-	-	-	3.6	99,824	1,100	3.7	102,793	1,100
Nursery	3.6	35,745	1,000	-	-	-	0.4	4,144	971	4.0	39,889	997
Greenhouse/Nursery	3.7	38,714	1,050	-	-	-	4.0	103,968	1,036	7.7	142,682	1,049
Recreational Turf	123.0	1,150,637	936	-	-	-	97.6	952,924	977	220.5	2,103,562	954
Vegetable	122.8	548,470	447	-	-	-	239.5	1,060,860	443	362.3	1,609,330	444
<b>TOTALS</b>	<b>3,691.3</b>	<b>27,710,576</b>	<b>751</b>	<b>-</b>	<b>-</b>	<b>-</b>	<b>2,669.0</b>	<b>18,513,848</b>	<b>694</b>	<b>6,360.3</b>	<b>46,224,424</b>	<b>727</b>

**Appendix Table M Buildout Crop Water Demand for Climate Change Data Circa 2050 and Good Management**

Climate Change	rcp26			rcp45			rcp85			Total		
Year	Irrigated Area (ha)	Irrigation Demand (m <sup>3</sup> )	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m <sup>3</sup> )	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m <sup>3</sup> )	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m <sup>3</sup> )	Avg. Req. (mm)
2053	6,360.3	49,621,769	780	6,360.3	45,921,885	722	6,360.3	58,254,082	916	6,360.3	51,265,912	806
2056	6,360.3	46,637,980	733	6,360.3	48,552,814	763	6,360.3	37,271,189	586	6,360.3	44,153,994	694
2059	6,360.3	27,944,806	439	6,360.3	58,546,261	920	6,360.3	58,916,566	926	6,360.3	48,469,211	762

**Appendix Table N Buildout Irrigation System Demand for 2003 Climate Data and Good Management**

Water Source	Surface Water			Reclaimed Water			Groundwater			Total		
Agriculture Irrigation System	Irrigated Area (ha)	Irrigation Demand (m <sup>3</sup> )	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m <sup>3</sup> )	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m <sup>3</sup> )	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m <sup>3</sup> )	Avg. Req. (mm)
Drip	567.2	2,576,717	454	-	-	-	1,292.7	6,664,970	516	1,859.8	9,241,687	497
Flood	55.2	830,303	1,503	-	-	-	4.1	60,874	1,500	59.3	891,177	1,503
GolfSprinkler	-	-	-	-	-	-	19.7	165,180	837	19.7	165,180	837
Gun	25.9	301,614	1,166	-	-	-	17.2	203,183	1,184	43.0	504,796	1,173
Handline	216.0	1,919,342	889	-	-	-	207.2	1,938,721	936	423.2	3,858,063	912
Landscapesprinkler	17.2	148,855	865	-	-	-	50.2	490,198	976	67.4	639,052	948
Microspray	-	-	-	-	-	-	0.2	1,606	750	0.2	1,606	750
Microsprinkler	-	-	-	-	-	-	0.4	4,144	971	0.4	4,144	971
Pivot	-	-	-	-	-	-	14.9	155,812	1,047	14.9	155,812	1,047
PivotLP	2,027.0	15,154,319	748	-	-	-	453.6	3,279,918	723	2,480.6	18,434,236	743
SDI	-	-	-	-	-	-	0.1	366	378	0.1	366	378
Sprinkler	456.3	3,914,918	858	-	-	-	172.1	1,655,942	962	628.4	5,570,860	887
Ssovertree	-	-	-	-	-	-	3.9	35,164	905	3.9	35,164	905
Sssprinkler	-	-	-	-	-	-	7.0	72,742	1,035	7.0	72,742	1,035
Ssundertree	0.8	7,814	941	-	-	-	2.9	25,721	880	3.8	33,535	893
Travgun	-	-	-	-	-	-	0.4	3,711	932	0.4	3,711	932
Wheeline	325.7	2,856,694	877	-	-	-	422.4	3,755,599	889	748.1	6,612,292	884
<b>TOTALS</b>	<b>3,691.3</b>	<b>27,710,576</b>	<b>751</b>	<b>-</b>	<b>-</b>	<b>-</b>	<b>2,669.0</b>	<b>18,513,848</b>	<b>694</b>	<b>6,360.3</b>	<b>46,224,424</b>	<b>727</b>

**Appendix Table O Buildout Water Demand by Purveyor for 2003 Climate Data and Good Management**

Water Source	Surface Water			Reclaimed Water			Groundwater			Total		
Agriculture Purveyor	Irrigated Area (ha)	Irrigation Demand (m³)	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m³)	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m³)	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m³)	Avg. Req. (mm)
Cawston	-	-	-	-	-	-	219.0	1,521,411	695	219.0	1,521,411	695
Fairview Heights	-	-	-	-	-	-	272.5	1,682,176	617	272.5	1,682,176	617
Keremeos	39.9	185,711	465	-	-	-	690.3	4,688,281	679	730.2	4,873,992	667
Lower Similkameen First Nations	499.7	3,811,137	763	-	-	-	20.9	189,579	905	520.6	4,000,716	768
Upper Similkameen First Nations	126.7	984,108	777	-	-	-	8.9	78,042	881	135.6	1,062,150	783
Private	3,025.0	22,729,620	751	-	-	-	1,457.4	10,354,359	710	4,482.3	33,083,979	738
<b>TOTALS</b>	<b>3,691.3</b>	<b>27,710,576</b>	<b>751</b>	<b>-</b>	<b>-</b>	<b>-</b>	<b>2,669.0</b>	<b>18,513,848</b>	<b>694</b>	<b>6,360.3</b>	<b>46,224,424</b>	<b>727</b>

**Appendix Table P Buildout Water Demand by Local Government for 2003 Climate Data and Good Management**

Water Source	Surface Water			Reclaimed Water			Groundwater			Total		
Agriculture Local Government	Irrigated Area (ha)	Irrigation Demand (m³)	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m³)	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m³)	Avg. Req. (mm)	Irrigated Area (ha)	Irrigation Demand (m³)	Avg. Req. (mm)
Keremeos	-	-	-	-	-	-	29.7	199,439	671	29.7	199,439	671
Okanagan-Similkameen	3,057.5	22,855,479	748	-	-	-	2,515.3	17,331,168	689	5,572.8	40,186,647	721
Princeton	7.4	59,851	809	-	-	-	3.1	25,793	838	10.5	85,645	817
Thompson-Nicola	-	-	-	-	-	-	34.3	185,371	540	34.3	185,371	540
Lower Similkameen First Nations	499.7	3,811,137	763	-	-	-	20.9	189,579	905	520.6	4,000,716	768
Upper Similkameen First Nations	126.7	984,108	777	-	-	-	8.9	78,042	881	135.6	1,062,150	783
Private	-	-	-	-	-	-	56.7	504,456	889	56.7	504,456	889
<b>TOTALS</b>	<b>3,691.3</b>	<b>27,710,576</b>	<b>751</b>	<b>-</b>	<b>-</b>	<b>-</b>	<b>2,669.0</b>	<b>18,513,848</b>	<b>694</b>	<b>6,360.3</b>	<b>46,224,424</b>	<b>727</b>