

Hairy Water-clover (*Marsilea vestita*) Marsileaceae (Water Clover Family)

<u>RANGE</u>

- In western North America from British Columbia to Saskatchewan, south to California through Arkansas
- In B.C., known from four sites near Kamloops and three in the Okanagan Valley (two north of Kelowna, one along Osoyoos Lake); populations at two of the Okanagan sites may have been extirpated




Figure 1 B.C. distribution of Marsilea vestita (BC CDC 2014)

<u>HABITAT</u>

- Seasonally submerged or wet, usually sand-rich soils of open depressions or floodplains alongside lakes and rivers in the Bunchgrass, Ponderosa Pine and Interior Douglas-fir Biogeoclimatic Zones
- Associates include spike-rushes, Engelmann's spike-rush (*Eleocharis engelmannii*), awned cyperus (*Cyperus squarrosus*), bentgrasses (*Agrostis* spp.), common witchgrass (*Panicum capillare*) and toothcup meadow-foam (*Rotala ramosior*)



Figure 2 Seasonally submerged open depression habitat Ministry of Forests, Lands and Natural Resource Operations | Thompson Okanagan Region Resource Management | Ecosystems Section | Penticton, B.C.



Figure 3 Marsilea vestita growing in open sandy substrate

LIFE HISTORY

- Perennial fern that annually grows vegetatively via rhizomes, forming small patches in open soil
- Sporocarps produced from summer into October
- When water conditions are favourable, sporocarp absorbs water and expels male spores (microspores) and female spores (megaspores)
- Spores germinate into tiny male and female plants, which produce sperm and eggs, respectively
- Female plant will form a gelatinous layer that traps sperm found in water surrounding plant
- When fertilization takes place, a young plant is formed that eventually produce fronds




Figure 4 Illustration of *Marsilea vestita* by Jeanne R. Janish (Hitchcock et al. 1969)

Status: Red / Not Assessed Best Survey Time: Jul to Sep General Habitat: Wetland

Marsilea vestita (continued)

DESCRIPTION

<u>General</u>

- Semi-terrestrial, perennial fern
- Each plant composed of numerous, thread-like stems topped by 'four leaf clover'-like leaflets (fronds) that grow from buds along thin, creeping underground rhizomes

Leaves

- Individual leaflets wedge-shaped and rounded, from 0.5 to 3 cm long and wide
- When growing under water, leaves are either submerged or floating on water surface
- When terrestrial, following water drawdown, leaflets held above ground on often arching stems

Sporocarps

- Spores produced in distinctive pod-like sporocarps, which are clustered on short stalks at base of leaves
- Sporocarps 4 to 8 mm long and slightly hairy with two distinctive bumps at one end near stalk




Figure 5 Patch showing thread-like stems, creeping rhizomes and sporocarp (arrow); growing with red-listed *Eleocharis engelmannii*

IDENTIFICATION TIPS

- Readily identified by its 'four leaf clover'-like leaves
- Flowerless clovers (*Trifolium* spp.) are distinguished from *Marsilea vestita* by usually having three leaflets and growing from fibrous roots
- Except for clovers, which occasionally grow nearby, no other plant in the habitats where *Marsilea vestita* is found can be mistaken for it



Figure 6 Plants showing characteristic 'four leaf clover'-like leaflets

GENERAL THREATS AND GUIDANCE

- Avoid development in areas with known occurrences of *Marsilea vestita* through project relocation or redesign
- Protect seasonally submerged and wet habitat alongside lakes and rivers from disturbance and development, including exclusion of livestock and ATVs through fencing and/or signage, and consider restoration including invasive plant removal following professional advice
- Follow provincial methods for when and how to conduct plant species at risk surveys
- Follow provincial policy and guidance on how to avoid, minimize, restore and offset impacts to plant species at risk and their habitats
- Report any sightings to the B.C. Conservation Data Centre (<u>cdcdata@gov.bc.ca</u>) and FLNR Ecosystems Section (josie.symonds@gov.bc.ca)

REFERENCES

B.C. Conservation Data Centre. 2014. http://a100.gov.bc.ca/pub/eswp/

Douglas et al. (editors). 2000. *Illustrated Flora of British Columbia*. Vol. 5. B.C. Min. Environ., Lands and Parks, and Min. For., Victoria, B.C.

Hitchcock et al.. 1969. Vascular Plants of the Pacific Northwest. Univ. Washington Press, Seattle, W.A.

Klinkenberg, Brian (editor). 2013. E-Flora BC. <u>http://linnet.geog.ubc.ca/Atlas/Atlas.aspx?sciname=M</u> <u>arsilea%20vestita</u>

NatureServe. 2012. <u>http://www.NatureServe.org/explorer</u>

<u>ACKNOWLEDGEMENTS</u>

Marsilea vestita Plant Species at Risk Fact Sheet developed by Josie Symonds, based on content produced under contract by Terry McIntosh and review comments by Brenda Costanzo and Orville Dyer