

Introduction

This is the fourth annual air quality report for the Southern Interior Air Zone. Annual air zone reporting is a commitment under the national Air Quality Management System (AQMS). This report describes achievement of the Canadian Ambient Air Quality Standards (CAAQS) for ground-level ozone (O₃) and fine particulates (PM_{2.5}), the associated management levels and recent actions to improve air quality. A province-wide summary can be found at: <http://www.env.gov.bc.ca/soe/indicators/air/>.

Background

The AQMS is the national approach to managing air quality in Canada. Under the AQMS, the CAAQS are developed to drive action to protect human health and the environment. Air zones are areas that exhibit similar air quality characteristics, issues and trends, and that form the basis for monitoring, reporting and taking action on air quality. The Southern Interior Air Zone (see Figure 1) is one of seven broad air zones across the province. Under the AQMS progressively more rigorous actions are expected as air quality approaches or exceeds the CAAQS. The level of action is guided by the Air Zone Management Framework outlined in Table 1.




Figure 1. Southern Interior Air Zone.

Table 1. Air zone management framework for ground-level ozone and PM_{2.5}. The CAAQS define the upper threshold, separating the “red” and “orange” management levels.

Management Level	O ₃ (ppb)		PM _{2.5} – Annual (µg/m ³)		PM _{2.5} - 24h (µg/m ³)	
	2015	2020	2015	2020	2015	2020
Red	Actions for Achieving Air Zone CAAQS					
Threshold (CAAQS)	63	62	10	8.8	28	27
Orange	Actions for Preventing CAAQS Exceedance					
Threshold	56		6.4		19	
Yellow	Actions for Preventing Air Quality Deterioration					
Threshold	50		4		10	
Green	Actions for Keeping Clean Areas Clean					

Ozone Levels

Ozone measurements in the Southern Interior Air Zone are summarized in Figure 2.

Concentrations ranged from 42 ppb in Castlegar to 53 ppb in Kelowna.¹ All sites were below the national standard of 63 ppb.

Trends in ozone levels are shown in Figure 3.² Concentrations have remained below the level of the national standard throughout this period, with the exception of 2012 in Kelowna, when wildfire smoke led to higher ozone concentrations.³




Figure 2. Ozone concentrations in the Georgia Strait Air Zone, based on annual 4th highest daily 8-hour maxima, averaged over 2014-2016. Red dashed line identifies the CAAQS of 63 ppb.




Figure 3. Trends in ozone concentrations (2007-2016), based on annual 4th highest daily 8-hour maxima for a single year. Red dashed line identifies CAAQS of 63 ppb. Asterisk (*) flags combined dataset from multiple sites in Kamloops.

¹ Concentrations based on 4th highest daily 8-hour maximum, averaged over three years (2014-2016).

² Concentrations based on 4th highest daily 8-hour maximum, averaged over a single year.

³ Teakles, A.D., So, Rita, Ainslie, B. et al. (2017) Impacts of the July 2012 Siberian fire plume on air quality in the Pacific Northwest. Atmos. Chem. Phys. 17, pp. 2593-2611.

PM_{2.5} Levels

PM_{2.5} refers to inhalable particles up to 2.5 micrometres in diameter. PM_{2.5} measurements are summarized in Figure 4. A distinction is made between data collected using the new Federal Equivalent Method (FEM) technology and the older TEOM instruments that are being phased out. The FEMs are the preferred instrument as they provide a more complete measure of PM_{2.5} than the TEOMs.

Daily concentrations (upper plot) ranged from 18 µg/m³ in Grand Forks to 52 µg/m³ in Castlegar.⁴ Castlegar was the only site to exceed the national standard of 28 µg/m³, as a result of wildfire influences. Annual concentrations (lower plot) ranged from 5.8 µg/m³ in Grand Forks to 8.5 µg/m³ in Golden.⁵ All monitoring sites achieved the national standard of 10 µg/m³.

Trends in annual mean concentrations between 2007 and 2016 are shown in Figure 5 for a subset of these sites.⁶ A shift to higher reported concentrations is seen with the change from TEOM to FEM instruments from about 2010 onward. However, all monitoring sites remained below the CAAQS level.




Figure 4. PM_{2.5} concentrations in Southern Interior Air Zone. Upper plot based on 24-hour concentration (annual 98th percentile, averaged over 2014-2016). Lower plot based on annual mean concentration (averaged over 2014-2016). Red dashed lines identify CAAQS of 28 µg/m³ (upper plot) and 10 µg/m³ (lower plot). Asterisk (*) flags combined dataset from multiple sites in Golden.

⁴ Concentrations based on the annual 98th percentile of 24-hour values, averaged over three years (2014-2016).

⁵ Concentrations based on the annual average of 24-hour values, averaged over three years (2014-2016).

⁶ Concentrations based on the annual average of 24-hour values over a single year.




Figure 5. Trends in PM_{2.5} concentrations (2007-2016), based on annual mean concentrations from a single year. The CAAQS value of 10 µg/m³ is shown by the dashed line. PM_{2.5} measurements prior to 2011 are reported at 25°C and 1 atm. From 2011 onward, measurements are reported at local conditions. Asterisk (*) flags combined dataset from more than one site in community.

Air Zone Management Levels

Air zone management levels are assigned on the basis of the highest concentrations within an air zone, excluding contributions from transboundary flows (TF) and exceptional events (EE) such as wildfires. This is done so that long-term management strategies are not developed on the basis of events that are beyond local or provincial control.

In the Southern Interior Air Zone, wildfires are the primary contributor to TF/EE. The methodology for identifying wildfire-influenced data is provided in Appendix I and excluded data are summarized in Appendix II. The summers of 2014 and 2015 were characterized by hot, dry conditions and an above-average number of hectares burned compared to 2016. These fires created smoky conditions and periods of degraded air quality in several communities across the air zone.

Table 2 summarizes the as-measured concentrations for ground-level ozone and the management levels after consideration of TF/EE influences (none were identified). The Southern Interior Air Zone is assigned a “yellow” management level based on ozone concentrations in Kamloops and Kelowna. This indicates that ozone-related actions should focus on preventing further air quality deterioration.

Table 2. Summary of ozone concentrations as measured and air zone management levels for the Southern Interior Air Zone (based on 2014-2016 data).

Location	No. Valid Years	4 th Highest Daily 8-hour Maxima		Air Zone Management Level
		As Measured	TF/EE Influences Removed	
Castlegar	2	42	42	Goal: Preventing Further Deterioration
Kamloops	3	52	52	
Kelowna	3	53	53	
Vernon	3	50	50	

Table 3 summarizes PM_{2.5} concentrations as measured and with TF/EE influences removed for each monitoring site (see Appendix II for more information on excluded data). The impact of removing such data changed management levels for Castlegar from “red” to “orange”. As air zone levels are preferentially based on three complete years of data, the Southern Interior Air Zone is assigned an “orange” management level, based on 24-hour concentrations in Golden and Kamloops and annual concentrations in Golden, Kamloops and Vernon. This indicates that PM_{2.5}-related activities would be prudent to prevent future CAAQS exceedances.

Table 3. Summary of PM_{2.5} concentrations as measured and air zone management levels for the Southern Interior Air Zone (based on 2014-2016 data).

Location	Monitor Type	No. Valid Years	Daily Mean (98 th Percentile)		Annual Mean		Air Zone Management Level
			As Measured	TF/EE Removed	As Measured	TF/EE Removed	
Castlegar	FEM	3*	52	18	7.3	7.1	Goal: Preventing CAAQS Exceedance
Golden	FEM	3	23	21	8.5	8.0	
Grand Forks	TEOM	3	18	16	5.8	5.0	
Kamloops	FEM	3	22	20	8.4	8.2	
Kelowna	TEOM/ FEM	3	20	16	6.3	5.7	
Vernon	TEOM/ FEM	3	23	17	7.6	7.1	

*Incomplete year of data in 2015 was included in calculation of daily mean value but did not meet data requirements for calculation of annual mean value.

Actions to Protect Air Quality

The reduction of smoke-related PM_{2.5} emissions has been a priority in a number of communities across the Southern Interior Air Zone for several years. Strategies and actions to reduce PM_{2.5} emissions have been documented in local airshed plans that have been developed and implemented for the Central Okanagan Regional District,⁷ City of Kamloops,⁸ the Boundary airshed,⁹ and Merritt.¹⁰ Golden is host to an active air quality committee.¹¹

Between 2014-2016, wood stove change-out programs were supported in the Regional Districts of Kootenay-Boundary, Central Kootenay and Okanagan-Similkameen as well as in Kimberly, Cranbrook and Golden to encourage residents to change out their older, smoky wood stoves for low-emission appliances. Funding support is also being provided to the District of Coldstream in 2017.

A description of other activities underway in B.C. air zones can be found in the “Air Zone Management Response for British Columbia” (see: www.gov.bc.ca/bcairquality).

7

http://www.regionaldistrict.com/media/217275/RDCO_2015_Clean_Air_Strategy_Final_DRAFT_2015_02_03_final.pdf

⁸ <http://www.kamloops.ca/environment/pdfs/13-05-AirshedManagementPlan.pdf>

⁹ <http://www.grandforks.ca/air/aqmplans/GrandForksAQMP-Oct22.pdf>

¹⁰ http://www.env.gov.bc.ca/epd/bcairquality/reports/pdfs/merritt_aqmp.pdf

¹¹ <http://www.goldenairquality.ca/>

Appendix I – Approach to Identify Wildfire-influenced Data

Summertime air quality in British Columbia is periodically influenced by wildfire smoke – from local fires as well as long-range transport from outside of the province. The wildfire season in B.C. typically occurs between May and September, when warm and dry conditions prevail.

A myriad of different pollutants are emitted from wildfires, including PM_{2.5} and gases that include nitrogen oxides and volatile organic compounds that can react in the atmosphere to form ground-level ozone and additional PM_{2.5}.

Given that smoke-affected areas may be extensive, and that smoke may linger for days before being fully dispersed from an airshed, the current analysis has focussed on those periods when wildfire smoke may have contributed to an exceedance of the CAAQS levels for PM_{2.5} levels. Criteria used to flag and evaluate wildfire-influenced data included the following:

- 24-hour PM_{2.5} concentrations exceeded the CAAQS level of 28 µg/m³ or 8-hour daily maximum ozone concentrations exceeded the CAAQS level of 63 ppb between May and September,
- Wildfires of interest were identified based on data from B.C. Wildfire Management Branch,
- Wildfire smoke advisories had been issued by the Ministry of Environment & Climate Change Strategy during the period of interest,
- MODIS satellite images indicated smoke impacts over the region,
- Back-trajectory analyses indicate that air parcel over area may have passed over wildfires,
- Multiple monitoring sites in the area of concern exhibited similar air quality characteristics, suggesting a common source or contributing source, and
- Modelling studies identify enhanced pollutant concentrations due to wildfire smoke.

Wildfire-influenced data were excluded from the calculation of air zone management levels. Excluded data are as summarized in Appendix II.

Appendix II – Wildfire-influenced Data in the Southern Interior Air Zone (2014-2016)

Ozone and PM_{2.5} data from 2014-2016 for the Southern Interior Air Zone were evaluated based on the criteria set out in Appendix I for TF/EE influences. Wildfire-influenced PM_{2.5} data are summarized in Table II-1. Supporting evidence included the following:

- There were a number of wildfires contributing to smoke in the air zone in 2014 and 2015. Of particular note was the Stickpin fire (21,965 ha) discovered on August 11, 2015 about 5km south of the US/Canada border near Grand Forks, and which burned for several weeks. Other major fires included the Rock Creek wildfire (4,417 ha) discovered August 13, 2015, the Testalinden Creek wildfire (5,133 ha) discovered August 14, 2015 6 km west of Oliver.
- Days flagged as potentially wildfire-influenced generally coincided or preceded with wildfire smoke-related advisories issued by the Ministry of Environment & Climate Change Strategy (see Table II-1).
- The single-day event in Grand Forks on Sept. 12, 2015 was associated with smoke from Washington wildfires (see. NASA MODIS satellite image at: https://lance3.modaps.eosdis.nasa.gov/imagery/subsets/?subset=AERONET_Kelowna.2015255.terra.1km.jpg).

Table II-1. Wildfire-influenced PM_{2.5} data.

Location	Date	24-Hr PM _{2.5} (µg/m ³)	Wildfire Smoke-related Air Quality Advisory?
Kamloops Federal Building	2014-07-16	61.4	Y
Vernon Science Centre	2014-07-16	37.7	Y
Kamloops Federal Building	2014-07-17	53.5	Y
Vernon Science Centre	2014-07-17	37.6	Y
Creston PC School	2014-07-18	39.8	Y
Castlegar Zinio Park	2014-07-19	34.4	Y
Creston PC School	2014-07-19	48.4	Y
Kamloops Federal Building	2014-08-06	28.2	Y
Castlegar Zinio Park	2014-08-07	34.5	
Castlegar Zinio Park	2014-08-08	29.6	Y
Grand Forks City Hall	2014-08-10	42	
Grand Forks City Hall	2014-08-11	33.8	Y
Grand Forks City Hall	2014-08-12	37.8	Y
Kelowna College	2014-08-12	32.2	Y
Vernon Science Centre	2014-08-12	29.5	Y
Kelowna College	2015-07-07	30.1	Y
Vernon Science Centre	2015-07-07	28.9	Y
Kamloops Federal Building	2015-07-10	38.3	Y
Creston PC School	2015-08-14	31.3	
Grand Forks City Hall	2015-08-14	29.5	

Table II-1 (continued)

Location	Date	24-Hr PM _{2.5} (µg/m ³)	Wildfire Smoke-related Air Quality Advisory?
Castlegar Zinio Park	2015-08-15	47.7	Y
Creston PC School	2015-08-15	33.5	Y
Castlegar Zinio Park	2015-08-19	41.6	
Grand Forks City Hall	2015-08-19	55	
Castlegar Zinio Park	2015-08-20	95.3	Y
Creston PC School	2015-08-20	91.1	Y
Grand Forks City Hall	2015-08-20	84.6	Y
Grand Forks City Hall	2015-08-22	28.5	Y
Castlegar Zinio Park	2015-08-23	53.6	Y
Kamloops Federal Building	2015-08-23	78.9	T
Kelowna College	2015-08-23	294.5	Y
Vernon Science Centre	2015-08-23	142.7	Y
Castlegar Zinio Park	2015-08-24	117	Y
Creston PC School	2015-08-24	77.3	Y
Kamloops Federal Building	2015-08-24	45.4	Y
Kelowna College	2015-08-24	183.5	Y
Vernon Science Centre	2015-08-24	103.8	Y
Castlegar Zinio Park	2015-08-25	165.8	Y
Creston PC School	2015-08-25	76.7	Y
Grand Forks City Hall	2015-08-25	311.8	Y
Kelowna College	2015-08-25	43.3	Y
Vernon Science Centre	2015-08-25	31.8	Y
Castlegar Zinio Park	2015-08-26	126.7	Y
Creston PC School	2015-08-26	70.3	Y
Grand Forks City Hall	2015-08-26	241.7	Y
Kelowna College	2015-08-26	79.6	Y
Vernon Science Centre	2015-08-26	72.8	Y
Castlegar Zinio Park	2015-08-27	125.5	Y
Grand Forks City Hall	2015-08-27	250.8	Y
Kelowna College	2015-08-27	53.7	Y
Vernon Science Centre	2015-08-27	68.7	Y
Castlegar Zinio Park	2015-08-28	137	Y
Grand Forks City Hall	2015-08-28	165.1	Y
Kelowna College	2015-08-28	29.2	Y
Vernon Science Centre	2015-08-28	41.6	Y
Castlegar Zinio Park	2015-08-29	47.2	Y
Kelowna College	2015-08-29	36.9	Y
Vernon Science Centre	2015-08-29	34.2	Y
Grand Forks City Hall	2015-09-12	29.6	

