

# Prince George Ambient Air Monitoring Program 2000 Air Quality Data Summary

## INTRODUCTION

The 2000 Prince George Air Quality Data Summary summarizes the data of the joint government/industry program to monitor reduced sulphur (TRS), sulphur dioxide (SO<sub>2</sub>), nitrogen oxides (NO<sub>x</sub>), particulate less than ten microns (PM<sub>10</sub>), particulate less than two and half microns (PM<sub>2.5</sub>), and ground level ozone (O<sub>3</sub>) in Prince George.

Interpretation of the data is not provided in this summary report, but will be available in the final Annual Air Quality Report. Anyone wishing further information on the data in this summary or on the monitoring network should contact one of the people listed at the bottom of this page.

## HIGHLIGHTS

- a) The total reduced sulphur (TRS) Provincial Level A one-hour objective (7µg/m<sup>3</sup>) was exceeded a maximum of 13.1% of time, at the Jail station, and a minimum of 1.9% at the Lakewood station. TRS levels in the Prince George area in 2000 exceeded the Provincial Level B one hour objective (28 µg/m<sup>3</sup>) from 0% of the time at the Lakewood station to 0.4% of the time at the Jail station. Annual averages for four out of the last five years at the Jail were less than 2.7 ug/m<sup>3</sup> and were the four lowest annual averages ever recorded at this site since monitoring began at the Jail. Plaza also recorded its four lowest annual averages in the past five years.
- b) The downtown-Plaza 400, and the Jail site, recorded annual averages of 7.4 µg/m<sup>3</sup> and 9.0 µg/m<sup>3</sup>, respectively, levels similar to those in 1999. The annual average SO<sub>2</sub> concentration recorded at the Jail was the lowest in nine years. The CBC station recorded the highest one-hour average concentration of (519 µg/m<sup>3</sup>).
- c) All four non-continuous PM<sub>10</sub> stations showed lower annual averages in 2000 compared to the previous year, and were the lowest since monitoring began. Lakewood recorded the lowest annual average of 14.1 µg/m<sup>3</sup>, whereas the CNR site recorded the highest annual average of 21.3 µg/m<sup>3</sup>. The BCR site recorded the greatest number of exceedances of the Level B daily criteria (50 µg/m<sup>3</sup>) for the continuous monitoring sites. However, the BCR site was the only continuous site that showed a decrease in 2000 whereas the other three sites showed an increase. It should be noted that PM<sub>2.5</sub> measured at the Downtown-Plaza station showed a sharp increase over the previous year.

### Contacts:

S. J. Lamble  
Air Resources Officer  
Environmental Section  
Omineca Peace Region  
250-565-6454

D. C. Fudge  
Air Quality Meteorologist  
Environmental Section  
Omineca Peace Region  
250-565-4210

## 1.0 TOTAL REDUCED SULPHUR (TRS) RESULTS

TRS describes a group of sulphur compounds containing gases made up primarily of four compounds; dimethyl disulphide (DMDS), dimethyl sulphide (DMS), methyl mercaptan (MESH), and hydrogen sulphide (H<sub>2</sub>S).

These compounds are largely by-products of the pulping process, that is, these compounds are not added directly but form because of the use of other sulphides in the pulping process. Pulp mills using the kraft process emit the largest amounts of TRS (in Prince George) followed by oil and gas processing plants (refining) and to a small extent automobiles (catalytic converters). Natural sources include swamps, bogs, and marshes.

Tables 1.1 to 1.4 list monthly summaries of TRS concentrations for each of the Prince George TRS ambient air stations and their respective annual summary for 2000.

Table 1.1  
2000 Monthly Summary of TRS Data at Jail

	Monthly Average ( $\mu\text{g}/\text{m}^3$ )	No. of 1-Hour Values		Maximum Hourly Average ( $\mu\text{g}/\text{m}^3$ )	No. of Daily Values		Maximum 24 Hour Average ( $\mu\text{g}/\text{m}^3$ )
		$>7\mu\text{g}/\text{m}^3$	$>28\mu\text{g}/\text{m}^3$		$>3\mu\text{g}/\text{m}^3$	$>6\mu\text{g}/\text{m}^3$	
<b>January</b>	3.2	121	1	28	13	2	18.1
<b>February</b>	3.3	148	0	24	13	6	11.3
<b>March</b>	1.7	57	1	41	8	0	10.7
<b>April</b>	1.3	35	0	23	5	0	5.8
<b>May</b>	1.6	55	0	21	6	0	4.9
<b>June</b>	1.6	64	0	24	7	2	8.1
<b>July</b>	1.1	37	0	26	3	0	7.6
<b>August</b>	0.7	12	0	18	1	0	4.4
<b>September</b>	3.6	118	23	62	11	8	17.4
<b>October</b>	3.8	168	9	38	13	7	18.4
<b>November</b>	2.8	105	0	26	8	6	10.7
<b>December</b>	4.1	159	3	40	17	7	13.8
<b>Annual</b>	<b>2.4</b>	<b>1079</b>	<b>37</b>	<b>62</b>	<b>105</b>	<b>38</b>	<b>18.4</b>

B.C. Ambient Air Quality Objectives  
Total Reduced Sulphur (TRS)

Level A

Level B

One Hour Average  
24 Hour Average

$7\mu\text{g}/\text{m}^3$   
 $3\mu\text{g}/\text{m}^3$

$28\mu\text{g}/\text{m}^3$   
 $6\mu\text{g}/\text{m}^3$

Table 1.2  
2000 Monthly Summary of TRS Data at Plaza

	Monthly Average ( $\mu\text{g}/\text{m}^3$ )	No. of 1-Hour Values		Maximum Hourly Average ( $\mu\text{g}/\text{m}^3$ )	No. of Daily Values		Maximum 24 Hour Average ( $\mu\text{g}/\text{m}^3$ )
		$>7\mu\text{g}/\text{m}^3$	$>28\mu\text{g}/\text{m}^3$		$>3\mu\text{g}/\text{m}^3$	$>6\mu\text{g}/\text{m}^3$	
January	2.7	94	6	35	8	3	21.8
February	3.0	116	0	23	12	4	11.3
March	0.7	26	0	20	2	0	7.1
April	0.6	13	0	10	1	0	3.2
May	0.6	17	0	16	0	0	3.5
June	0.9	40	0	18	4	0	6.8
July	0.9	27	0	20	3	0	5.2
August	0.4	13	0	13	0	0	3.9
September	2.1	71	8	45	7	5	12.2
October	2.1	84	2	35	9	3	11.2
November	1.2	51	0	24	4	1	7.0
December	3.2	123	0	26	12	5	18.3
Annual	1.5	675	16	45	62	21	21.8

Table 1.3  
2000 Monthly Summary of TRS Data at Lakewood

	Monthly Average ( $\mu\text{g}/\text{m}^3$ )	No. of 1-Hour Values		Maximum Hourly Average ( $\mu\text{g}/\text{m}^3$ )	No. of Daily Values		Maximum 24 Hour Average ( $\mu\text{g}/\text{m}^3$ )
		$>7\mu\text{g}/\text{m}^3$	$>28\mu\text{g}/\text{m}^3$		$>3\mu\text{g}/\text{m}^3$	$>6\mu\text{g}/\text{m}^3$	
January	1.3	37	0	21	4	1	12.8
February	1.0	25	0	16	2	2	7.8
March	0.1	0	0	6	0	0	1.8
April	0.1	0	0	3	0	0	1.2
May	0.2	0	0	4	0	0	0.9
June	0.1	1	0	7	0	0	2.2
July	0.1	0	0	6	0	0	1.2
August	0.1	0	0	4	0	0	1.0
September	1.0	23	0	23	3	0	6.1
October	0.5	12	0	14	0	0	3.1
November	0.3	0	0	6	0	0	1.5
December	1.4	59	0	20	6	2	9.7
Annual	0.5	157	0	23	15	5	12.8



Monthly averages ranged from a high of 4.1  $\mu\text{g}/\text{m}^3$  at Jail (December) to a low of 0.1  $\mu\text{g}/\text{m}^3$  at Lakewood. Jail had the highest annual average followed by Plaza and then Lakewood.

### 1.1 TRS Annual Trends

The TRS data from the Plaza (1980-2000), the Jail (1981-2000), and Lakewood (1982-2000) stations were examined to reveal trends in Prince George. The following tables (Table 1.4 - Table 1.6) summarize the TRS data from the three stations over the last nineteen to twenty-one years.

A major improvement in TRS levels have occurred from 1988 to 1990. Since 1990 TRS levels have declined slightly from year to year. The annual average concentrations for Plaza and Jail are one of the lowest since the monitoring program started. Annual averages in 2000 at the Plaza and Jail stations are approximately 75% lower than the 1986-1989 averages.

Table 1.4  
Annual Trend Summary of TRS Data at Jail

Year	Annual Average ( $\mu\text{g}/\text{m}^3$ )	No. (%) of 1-Hour Values		Maximum Hourly Average ( $\mu\text{g}/\text{m}^3$ )	No. (%) of Daily Values		Maximum Daily Average ( $\mu\text{g}/\text{m}^3$ )	No. of hours Instrument Operated
		$>7\mu\text{g}/\text{m}^3$	$>28\mu\text{g}/\text{m}^3$		$>3\mu\text{g}/\text{m}^3$	$>6\mu\text{g}/\text{m}^3$		
1981	7.2	1673 (21.7%)	638 (8.3%)	177	151 (47.0%)	121 (37.7%)	59.6	7701
1982	8.8	1169 (23.8%)	519 (10.6%)	149	125 (61.0%)	99 (48.3%)	53.5	4915
1983	11.0	2051 (27.3%)	1066 (14.2%)	156	184 (58.8%)	151 (48.2%)	70.2	7511
1984	6.5	1341 (17.6%)	593 (7.8%)	170	136 (42.8%)	109 (34.3%)	54.8	7638
1985	6.4	1483 (19.1%)	546 (7.0%)	149	145 (44.8%)	117 (36.1%)	44.9	7773
1986	9.8	2404 (29.1%)	1028 (12.4%)	347	197 (57.1%)	169 (49.0%)	68.0	8268
1987	11.6	2268 (29.0%)	1306 (16.7%)	382	200 (54.8%)	175 (47.9%)	86.0	7817
1988	11.6	2307 (27.9%)	1343 (16.2%)	312	197 (53.8%)	166 (45.4%)	94.6	8267
1989	9.2	1852 (25.8%)	793 (11.0%)	212	174 (47.7%)	134 (36.7%)	68.3	7183
1990	5.4	927 (14.3%)	215 (3.3%)	177	91 (24.9%)	66 (18.1%)	46.7	6494
1991	3.1	1195 (14.4%)	138 (1.7%)	126	117 (32.1%)	61 (16.7%)	39.2	8316
1992	3.7	1103 (14.9%)	71 (1.0%)	78	109 (29.9%)	66 (18.1%)	17.7	7387
1993	3.8	1343 (16.7%)	201 (2.5%)	69	123 (33.7%)	86 (23.6%)	37.7	8061
1994	3.1	1214 (15.1%)	108 (1.3%)	57	119 (32.6%)	67 (18.4%)	19.7	8017
1995	3.3	1274 (15.2%)	73 (0.9%)	64	157 (43.0%)	64 (17.5%)	23.0	8359
1996	2.6	1043 (12.5%)	42 (0.5%)	62	112 (30.7%)	53 (14.5%)	18.8	8359
1997	2.5	1157 (13.9%)	42 (0.5%)	55	119 (32.6%)	37 (10.1%)	21.3	8347
1998	3.1	1488 (17.9%)	57 (0.7%)	51	138 (37.8%)	62 (17.0%)	26.1	8314
1999	2.5	1085 (13.4%)	44 (0.5%)	123	107 (29.3%)	43 (11.8%)	26.5	8117
2000	2.4	1079 (13.1%)	37 (0.5%)	62	105 (28.8%)	43 (11.8%)	18.4	8214

Table 1.5  
Annual Trend Summary of TRS Data at Plaza

Year	Annual Average ( $\mu\text{g}/\text{m}^3$ )	No. (%) of 1-Hour Values		Maximum Hourly Average ( $\mu\text{g}/\text{m}^3$ )	No. (%) of Daily Values		Maximum Daily Average ( $\mu\text{g}/\text{m}^3$ )	No. of hours Instrument Operated
		$>7\mu\text{g}/\text{m}^3$	$>28\mu\text{g}/\text{m}^3$		$>3\mu\text{g}/\text{m}^3$	$>6\mu\text{g}/\text{m}^3$		
1980	5.9	1085 (16.0%)	449 (6.6%)	163	127 (45.0%)	85 (30.1%)	62.4	6772
1981	6.0	1276 (16.9%)	566 (7.5%)	241	136 (43.2%)	101 (32.1%)	53.8	7558
1982	5.3	1241 (15.1%)	445 (5.4%)	198	156 (45.5%)	111 (32.4%)	41.3	8235
1983	8.6	1670 (20.7%)	817 (10.1%)	177	180 (53.4%)	150 (44.5%)	90.5	8082
1984	4.7	978 (11.9%)	372 (4.5%)	227	121 (35.2%)	84 (24.4%)	51.7	8246
1985	4.8	875 (12.3%)	406 (5.7%)	297	106 (35.8%)	78 (26.4%)	81.6	7115
1986	7.9	1295 (20.1%)	677 (10.5%)	289	126 (46.8%)	102 (37.9%)	61.4	6456
1987	9.0	1806 (21.8%)	972 (11.7%)	218	174 (47.7%)	143 (39.2%)	69.0	8280
1988	7.3	1633 (19.7%)	792 (9.5%)	194	159 (43.4%)	121 (33.1%)	69.4	8297
1989	5.9	1376 (16.9%)	594 (7.3%)	184	140 (38.4%)	111 (30.4%)	41.8	8166
1990	2.7	905 (11.0%)	187 (2.3%)	122	100 (27.4%)	57 (15.6%)	47.0	8234
1991	2.2	925 (11.2%)	43 (0.5%)	58	84 (23.0%)	46 (12.6%)	19.4	8256
1992	2.1	882 (10.6%)	34 (0.4%)	69	89 (24.4%)	49 (13.4%)	18.4	8300
1993	2.4	956 (11.4%)	92 (1.1%)	72	89 (24.4%)	54 (14.8%)	29.8	8360
1994	1.9	790 (9.5%)	22 (0.3%)	67	84 (23.0%)	34 (9.3%)	19.1	8295
1995	1.9	757 (9.2%)	20 (0.2%)	44	93 (25.5%)	32 (8.8%)	20.5	8205
1996	1.7	592 (8.3%)	28 (0.4%)	103	67 (18.4%)	25 (6.8%)	27.2	7168
1997	1.6	654 (7.8%)	13 (0.2%)	40	65 (17.8%)	22 (6.0%)	12.9	8371
1998	2.2	958 (11.6%)	9 (0.1%)	50	88 (24.1%)	32 (8.8%)	19.5	8245
1999	1.7	784 (9.5%)	13 (0.2%)	44	71 (19.5%)	24 (6.6%)	18.5	8247

<b>2000</b>	1.5	675 (8.1%)	16 (0.2%)	45	58 (15.9%)	23 (5.5%)	21.8	8314
-------------	-----	------------	-----------	----	------------	-----------	------	------

Table 1.6  
Annual Trend Summary of TRS Data at Lakewood

Year	Annual Average ( $\mu\text{g}/\text{m}^3$ )	No. (%) of 1-Hour Values		Maximum Hourly Average ( $\mu\text{g}/\text{m}^3$ )	No. (%) of Daily Values		Maximum Daily Average ( $\mu\text{g}/\text{m}^3$ )	No. of hours Instrument Operated
		$>7\mu\text{g}/\text{m}^3$	$>28\mu\text{g}/\text{m}^3$		$>3\mu\text{g}/\text{m}^3$	$>6\mu\text{g}/\text{m}^3$		
1982	0.9	176 (3.0%)	22 (0.4%)	85	22 (9.1%)	13 (5.4%)	19.7	5785
1983	2.6	615 (7.9%)	147 (1.9%)	99	79 (24.5%)	43 (13.4%)	40.7	7737
1984	0.9	199 (2.4%)	15 (0.2%)	64	34 (9.9%)	10 (2.9%)	15.4	8269
1985	1.0	216 (2.9%)	28 (0.4%)	50	29 (9.3%)	12 (3.9%)	21.6	7456
1986	1.7	439 (5.6%)	110 (1.4%)	71	51 (15.7%)	32 (9.9%)	27.1	7807
1987	1.6	461 (6.0%)	106 (1.4%)	92	51 (14.0%)	34 (9.3%)	35.7	7680
1988	1.1	270 (3.6%)	38 (0.5%)	64	38 (10.4%)	23 (6.3%)	17.5	7462
1989	0.7	170 (2.8%)	33 (0.5%)	92	22 (6.0%)	8 (2.2%)	28.0	6057
1990	0.3	66 (1.2%)	3 (0.05%)	79	8 (2.2%)	1 (0.3%)	10.5	5484
1991	0.6	173 (2.3%)	3 (0.04%)	37	19 (5.2%)	4 (1.1%)	18.7	7667
1992	0.6	214 (2.6%)	0 (0.00%)	27	22 (6.0%)	9 (2.5%)	11.7	8351
1993	0.9	308 (4.4%)	6 (0.09%)	86	32 (8.8%)	13 (3.6%)	19.4	6952
1994	0.6	147 (1.8%)	2 (0.02%)	74	21 (5.8%)	4 (1.1%)	11.5	8196
1995	0.9	270 (3.2%)	2 (0.02%)	37	37 (10.1%)	11 (3.0%)	15.0	8358
1996	0.7	227 (2.7%)	8 (0.1%)	44	27 (7.4%)	7 (1.9%)	15.0	8281
1997	0.7	174 (2.1%)	0 (0.0%)	27	17 (4.7%)	4 (1.1%)	9.8	8211
1998	0.7	215 (2.6%)	0 (0.0%)	18	21 (5.6%)	5 (1.3%)	10.7	8204
1999	0.6	172 (2.1%)	1 (0.01%)	28	14 (3.8%)	5 (1.3%)	9.5	8191
2000	0.5	157 (1.9%)	0 (0.0%)	23	15 (4.1%)	3 (0.8%)	12.8	8348

\* Instrument Operated June - December 1982

## 2.0 SULPHUR DIOXIDE (SO<sub>2</sub>) RESULTS

Sulphur dioxide is formed primarily by the combustion of material containing sulphur. It is a colourless gas and at concentrations above  $900 \mu\text{g}/\text{m}^3$  (hourly Level B objective) has a pungent odour. At present, major sources of SO<sub>2</sub> in the Prince George area include the Husky refinery and the pulp mills.

The sulphur dioxide (SO<sub>2</sub>) data are summarized in Tables 2.1- 2.4. The average annual ambient SO<sub>2</sub> levels were within the Provincial objectives at all stations. Only the CBC transmitter site recorded more than one Level A hourly exceedance. This station also recorded the highest one-hour average concentration ( $519 \mu\text{g}/\text{m}^3$ ). Both Plaza and Jail stations did not record any exceedance of the Provincial objectives.

B.C. Ambient Air Quality Objectives  
Sulphur Dioxide (SO<sub>2</sub>)

Level A

Level B

One Hour Average

$450 \mu\text{g}/\text{m}^3$

$900 \mu\text{g}/\text{m}^3$

24 Hour Average  
Annual Average

160  $\mu\text{g}/\text{m}^3$   
25  $\mu\text{g}/\text{m}^3$

260  $\mu\text{g}/\text{m}^3$   
50  $\mu\text{g}/\text{m}^3$

Table 2.1  
2000 Monthly Summary of SO<sub>2</sub> Data at Jail

	Monthly Average ( $\mu\text{g}/\text{m}^3$ )	No. of 1-Hour Values		Maximum Hourly Average ( $\mu\text{g}/\text{m}^3$ )	No. of Daily Values		Maximum 24 Hour Average ( $\mu\text{g}/\text{m}^3$ )
		>450 $\mu\text{g}/\text{m}^3$	>900 $\mu\text{g}/\text{m}^3$		>160 $\mu\text{g}/\text{m}^3$	>260 $\mu\text{g}/\text{m}^3$	
January	13.7	0	0	216	0	0	59.4
February	14.1	0	0	186	0	0	49.5
March	8.0	0	0	168	0	0	62.9
April	9.4	0	0	176	0	0	42.4
May	7.5	0	0	184	0	0	35.8
June	10.3	0	0	230	0	0	62.3
July	7.4	0	0	194	0	0	39.9
August	3.0	0	0	69	0	0	21.8
September	4.3	0	0	144	0	0	45.8
October	13.3	0	0	415	0	0	96.3
November	6.9	0	0	99	0	0	69.7
December	10.6	0	0	138	0	0	56.0
Annual	9.0	0	0	415	0	0	96.3

Table 2.2  
2000 Monthly Summary of SO<sub>2</sub> Data at Plaza

	Monthly Average ( $\mu\text{g}/\text{m}^3$ )	No. of 1-Hour Values		Maximum Hourly Average ( $\mu\text{g}/\text{m}^3$ )	No. of Daily Values		Maximum 24 Hour Average ( $\mu\text{g}/\text{m}^3$ )
		>450 $\mu\text{g}/\text{m}^3$	>900 $\mu\text{g}/\text{m}^3$		>160 $\mu\text{g}/\text{m}^3$	>260 $\mu\text{g}/\text{m}^3$	
January	8.4	0	0	149	0	0	45.0
February	17.0	0	0	168	0	0	49.9
March	7.1	0	0	176	0	0	38.2
April	4.8	0	0	181	0	0	29.7
May	3.2	0	0	80	0	0	16.0
June	6.4	0	0	170	0	0	38.2
July	5.4	0	0	189	0	0	29.4
August	3.8	0	0	290	0	0	31.9
September	6.8	0	0	117	0	0	34.4
October	11.1	0	0	341	0	0	58.7
November	6.0	0	0	80	0	0	23.2
December	9.3	0	0	136	0	0	41.0



<b>Annual</b>	<b>7.4</b>	<b>0</b>	<b>0</b>	<b>341</b>	<b>0</b>	<b>0</b>	<b>58.7</b>
---------------	------------	----------	----------	------------	----------	----------	-------------

Table 2.3  
2000 Monthly Summary of SO<sub>2</sub> Data at CBC Transmitter

	Monthly Average ( $\mu\text{g}/\text{m}^3$ )	No. of 1-Hour Values		Maximum Hourly Average ( $\mu\text{g}/\text{m}^3$ )	No. of Daily Values		Maximum 24 Hour Average ( $\mu\text{g}/\text{m}^3$ )
		>450 $\mu\text{g}/\text{m}^3$	>900 $\mu\text{g}/\text{m}^3$		>160 $\mu\text{g}/\text{m}^3$	>260 $\mu\text{g}/\text{m}^3$	
<b>January</b>	20.8	2	0	519	1	0	160.7
<b>February</b>	19.0	0	0	280	0	0	98.2
<b>March</b>	8.8	0	0	250	0	0	46.6
<b>April</b>	11.3	0	0	368	0	0	89.3
<b>May</b>	8.2	0	0	178	0	0	39.0
<b>June</b>	10.8	0	0	336	0	0	56.4
<b>July</b>	11.5	0	0	389	0	0	75.1
<b>August</b>	5.4	0	0	149	0	0	43.0
<b>September</b>	-	-	-	-	-	-	-
<b>October</b>	-	-	-	-	-	-	-
<b>November</b>	9.3	0	0	170	0	0	61.5
<b>December</b>	11.6	0	0	192	0	0	63.4
<b>Annual</b>	-	2	0	519	1	0	160.7

## 2.2 SO<sub>2</sub> Annual Trends

Table 2.4 summarizes the annual trend of SO<sub>2</sub> data collected at the Jail station. Prior to 1986, data was obtained from a Philips SO<sub>2</sub> continuous analyzer which was subject to interference from various TRS compounds resulting in erroneous data. In 1986 the Philips analyzer was replaced with a TECO analyzer utilizing newer technology. The SO<sub>2</sub> average for 1986 is based upon Philips data for Jan. - Feb. and TECO monitor data for Mar. - Dec. The lower annual averages after 1986 are attributable to the improved accuracy of data provided by the new monitor.

The trend in ambient SO<sub>2</sub> levels recorded at the Jail site from 1990 to 1996 had shown an increase. However, since 1996, SO<sub>2</sub> levels have been decreasing at this site. The annual concentration at the Jail site in 2000 has been the lowest recorded since 1991.

Table 2.4  
Annual Trend Summary of SO<sub>2</sub> Data at Jail

Year	Annual Average ( $\mu\text{g}/\text{m}^3$ )	No. of 1-Hour Values		Maximum Hourly Average ( $\mu\text{g}/\text{m}^3$ )	No. of Daily Values		Maximum Daily Average ( $\mu\text{g}/\text{m}^3$ )	No. of Hours Instrument Operated
		>450 $\mu\text{g}/\text{m}^3$	>900 $\mu\text{g}/\text{m}^3$		>160 $\mu\text{g}/\text{m}^3$	>260 $\mu\text{g}/\text{m}^3$		
1981	10.7	1	0	506	0	0	75.2	7,016
1982	25.5	1	0	719	0	0	161.4	6,535
1983	31.8	1	0	533	0	0	143.5	8,310
1984	22.9	6	1	1198	3	0	242.8	7,551
1985	21.0	0	0	399	0	0	119.3	7,860
1986*	16.9	4	0	538	0	0	152.5	8,278
1987	12.8	2	0	666	0	0	123.0	8,348
1988	13.8	8	0	586	1	0	173.8	8,412
1989	12.0	1	0	659	0	0	99.6	8,165
1990	5.5	0	0	283	0	0	56.7	6,719
1991	8.7	1	0	492	0	0	118.1	8,236
1992	12.1	1	0	450	0	0	104.0	8,363
1993	10.9	0	0	363	0	0	81.8	8,225
1994	12.5	1	0	469	0	0	123.0	8,162
1995	20.5	1	0	530	0	0	118.8	8,354
1996	23.7	10	0	735	2	0	189.8	8,368
1997	20.5	4	0	759	0	0	143.2	8,350
1998	18.6	1	0	469	0	0	129.0	7,580
1999	10.0	0	0	306	0	0	103.9	8,358
2000	9.0	0	0	415	0	0	96.3	8,362

\* Philips analyzer was replaced with a TECO analyzer

Tables 2.5 and 2.6 summarizes the annual trend of SO<sub>2</sub> data collected at the Plaza, and CBC sites, respectively. Annual average SO<sub>2</sub> concentrations recorded at the Jail were the lowest in nine years and at the CBC site, the lowest annual average since monitoring began.

Table 2.5  
Annual Trend Summary of SO<sub>2</sub> Data at Plaza

Year	Annual Average ( $\mu\text{g}/\text{m}^3$ )	No. of 1-Hour Values		Maximum Hourly Average ( $\mu\text{g}/\text{m}^3$ )	No. of Daily Values		Maximum Daily Average ( $\mu\text{g}/\text{m}^3$ )	No. of Hours Instrument Operated
		>450 $\mu\text{g}/\text{m}^3$	>900 $\mu\text{g}/\text{m}^3$		>160 $\mu\text{g}/\text{m}^3$	>260 $\mu\text{g}/\text{m}^3$		
1995	-	2	0	474	0	0	156.3	5,323*

<b>1996</b>	15.8	13	1	913	1	0	167.9	8,229
<b>1997</b>	14.6	8	0	663	0	0	118.3	8,363
<b>1998</b>	11.9	0	0	399	0	0	93.5	8,220
<b>1999</b>	7.2	1	0	490	0	0	105.3	8,703
<b>2000</b>	7.4	0	0	341	0	0	58.7	8,071

\* Instrument installed May 1995

Table 2.6  
Annual Trend Summary of SO<sub>2</sub> Data at CBC

Year	Annual Average ( $\mu\text{g}/\text{m}^3$ )	No. of 1-Hour Values		Maximum Hourly Average ( $\mu\text{g}/\text{m}^3$ )	No. of Daily Values		Maximum Daily Average ( $\mu\text{g}/\text{m}^3$ )	No. of Hours Instrument Operated
		>450 $\mu\text{g}/\text{m}^3$	>900 $\mu\text{g}/\text{m}^3$		>160 $\mu\text{g}/\text{m}^3$	>260 $\mu\text{g}/\text{m}^3$		
<b>1995</b>	-	4	0	597	0	0	101.1	5,096*
<b>1996</b>	23.9	10	0	794	4	0	189.3	8,370
<b>1997</b>	20.1	9	0	778	2	0	197.7	8,312
<b>1998</b>	22.6	10	0	597	1	0	224.0	8,361
<b>1999</b>	13.0	11	0	751	1	0	227.1	8,560
<b>2000</b>	-	2	0	519	0	0	142.4	6,958

\* Instrument installed May 1995

### 3.0 NITROGEN DIOXIDE (NO<sub>2</sub>) RESULTS

Nitrogen dioxide is a reddish-brown gas with a pungent and irritating odour over 250  $\mu\text{g}/\text{m}^3$ . The NO<sub>2</sub> absorbs short wave blue light, leaving longer wavelengths that cause it to appear reddish-brown.

NO<sub>2</sub> is emitted as a primary pollutant (emitted directly from high temperature combustion sources) or a secondary pollutant (produce from the oxidation of nitric oxide). Nitric oxide is formed during high temperature combustion, primarily from motor vehicle exhausts and stationary combustion (such as industrial processes, waste incineration, and fuel combustion for heating homes and buildings). Emissions of nitric oxide are greater during winter months when there is an increase in the use of heating fuels and in the idling of cars.

The nitrogen dioxide data from the Plaza site is summarized in Tables 3.1. Ambient NO<sub>2</sub> levels were well within the Provincial objectives, with no Level A hourly or daily exceedances. Operational problems with the instrument were noted in March, April, September, and October of 2000.

Table 3.1.  
2000 Monthly Summary of NO<sub>2</sub> Data at Plaza

	Monthly Average (µg/m <sup>3</sup> )	No. of 1-Hour Values		Maximum Hourly Average (µg/m <sup>3</sup> )	No. of Daily Values		Maximum 24 Hour Average (µg/m <sup>3</sup> )
		>400µg/m <sup>3</sup>	>1000µg/m <sup>3</sup>		>200µg/m <sup>3</sup>	>300µg/m <sup>3</sup>	
<b>January</b>	45.1	0	0	140	0	0	88.3
<b>February</b>	51.3	0	0	149	0	0	95.2
<b>March</b>	-	-	-	-	-	-	-
<b>April</b>	-	-	-	-	-	-	-
<b>May</b>	18.9	0	0	80	0	0	33.2
<b>June</b>	14.5	0	0	71	0	0	36.4
<b>July</b>	11.5	0	0	59	0	0	27.7
<b>August</b>	17.4	0	0	56	0	0	34.3
<b>September</b>	11.1	0	0	42	0	0	17.9
<b>October</b>	-	-	-	-	-	-	-
<b>November</b>	24.8	0	0	47	0	0	33.9
<b>December</b>	34.5	0	0	119	0	0	76.2
<b>Annual</b>	-	<b>0</b>	<b>0</b>	<b>149</b>	<b>0</b>	<b>0</b>	<b>101.6</b>

B.C. Ambient Air Quality Objectives ..... Level B  
Nitrogen Dioxide (NO<sub>2</sub>)

Level C

One Hour Average	400 µg/m <sup>3</sup>	1000 µg/m <sup>3</sup>
24 Hour Average	200 µg/m <sup>3</sup>	300 µg/m <sup>3</sup>
Annual Average	60 µg/m <sup>3</sup>	100 µg/m <sup>3</sup>

### 3.1 NO<sub>2</sub> Annual Trends

A trend discussion has not been available previously due to the shortage of data particularly in 1992 and 1995. A trend in NO<sub>2</sub> levels does not appear to be apparent as shown in Table 3.2. Annual average NO<sub>2</sub>

levels in 2000 were the highest to be recorded out of the eight full years of monitoring. The highest maximum daily NO<sub>2</sub> average ever recorded in Plaza occurred on March 12 2000.

Table 3.2  
Annual Trend Summary of NO<sub>2</sub> Data at Plaza

Year	Annual Average ( $\mu\text{g}/\text{m}^3$ )	No. of 1-Hour Values		Maximum Hourly Average ( $\mu\text{g}/\text{m}^3$ )	No. of Daily Values		Maximum Daily Average ( $\mu\text{g}/\text{m}^3$ )	No. of Hours Instrument Operated
		>400 $\mu\text{g}/\text{m}^3$	>1000 $\mu\text{g}/\text{m}^3$		>200 $\mu\text{g}/\text{m}^3$	>300 $\mu\text{g}/\text{m}^3$		
1992	-	0	0	61	0	0	97.0	4,825*
1993	25.4	0	0	149	0	0	83.3	8,072
1994	23.7	0	0	130	0	0	70.7	7,852
1995	-	0	0	101	0	0	61.5	4,957**
1996	25.9	0	0	151	0	0	84.6	7,787
1997	24.3	0	0	186	0	0	91.5	8,242
1998	28.7	0	0	143	0	0	67.1	8,051
1999	25.6	0	0	143	0	0	99.1	8,218
2000	-	0	0	149	0	0	101.6	5,871**

\* Note: NO<sub>2</sub> TECO Analyzer Installed in June 1992

\*\* Note NO<sub>2</sub> TECO Operational Problems

#### 4.0 OZONE (O<sub>3</sub>) RESULTS

Ground level ozone is produced when oxides of nitrogen (NO<sub>x</sub>) react with volatile hydrocarbons in the presence of sun light. Ozone is termed a secondary pollutant because it is produced from a photochemical reaction involving primary pollutants (NO<sub>x</sub> and volatile hydrocarbons).

Tables 4.1 summarizes the ozone data recorded at the Plaza station. The maximum one-hour level (132  $\mu\text{g}/\text{m}^3$ ) was recorded on May 2<sup>nd</sup>. The greatest number of exceedances of the one-hour objective occurred in April. Exceedances of the one-hour Level A objective in 2000 occurred between the months of March and July. Exceedances of the 24-hour Level A objective occurred eight or more days during every month in 2000. July and October was the only months in which there were no exceedance of the 24-hour Level B objective. There is some debate whether the daily objective is below the background level for ozone.

Table 4.1  
2000 Monthly Summary of O<sub>3</sub> Data for Plaza

	Monthly Average ( $\mu\text{g}/\text{m}^3$ )	No. of 1-Hour Values		Maximum Hourly Average ( $\mu\text{g}/\text{m}^3$ )	No. of Daily Values		Maximum 24 Hour Average ( $\mu\text{g}/\text{m}^3$ )
		>100 $\mu\text{g}/\text{m}^3$	>160 $\mu\text{g}/\text{m}^3$		>30 $\mu\text{g}/\text{m}^3$	>50 $\mu\text{g}/\text{m}^3$	
January	27.9	0	0	78	14	4	60.6
February	26.8	0	0	98	9	3	90.6
March	53.3	7	0	103	29	17	86.1
April	71.5	90	0	120	28	25	98.7
May	66.4	70	0	132	29	27	107.3
June	44.6	8	0	120	24	9	68.4
July	29.6	5	0	118	14	0	45.5
August	28.3	0	0	98	15	1	59.2
September	24.7	0	0	78	10	0	59.1
October	19.4	0	0	78	4	0	51.1
November	23.3	0	0	70	9	0	62.1
December	20.5	0	0	68	10	1	62.1
<b>Annual</b>	<b>36.4</b>	<b>180</b>	<b>0</b>	<b>132</b>	<b>195</b>	<b>87</b>	<b>107.3</b>

B.C. Ambient Air Quality Objectives  
Ozone (O<sub>3</sub>)

## Level A

## Level B

One Hour Average  
24 Hour Average  
Annual Average

100  $\mu\text{g}/\text{m}^3$   
30  $\mu\text{g}/\text{m}^3$

160  $\mu\text{g}/\text{m}^3$   
50  $\mu\text{g}/\text{m}^3$   
30  $\mu\text{g}/\text{m}^3$

4.1 O<sub>3</sub> Annual Trends

A discussion in the trend has not been available previously due to the shortage of data since installation occurred in 1995. The annual average O<sub>3</sub> level in 2000 was lower than the annual average of the previous two full years of data, but there were more daily exceedances of the Level A Hourly objective in 2000.

Table 4.2  
Annual Trend Summary of O<sub>3</sub> Data at Plaza

Year	Annual Average ( $\mu\text{g}/\text{m}^3$ )	No. of 1-Hour Values		Maximum Hourly Average ( $\mu\text{g}/\text{m}^3$ )	No. of Daily Values		Maximum Daily Average ( $\mu\text{g}/\text{m}^3$ )	No. of Hours Instrument Operated
		>100 $\mu\text{g}/\text{m}^3$	>160 $\mu\text{g}/\text{m}^3$		>30 $\mu\text{g}/\text{m}^3$	>50 $\mu\text{g}/\text{m}^3$		
1995	-	28	0	112	108	29	75.2	5,548*
1996	35.2	21	0	120	202	83	88.3	8,290
1997	35.2	54	0	120	217	88	81.8	8,358
1998	37.2	159	0	156	214	93	100.0	8,284
1999	37.8	140	0	140	222	86	102.0	8,638
2000	36.4	180	0	132	193	95	107.3	8,309

\* Note: O<sub>3</sub> API Analyzer installed in late April 1995

## 5.0 PARTICULATE MATTER (PM<sub>10</sub>) RESULTS

Particulate matter refers to small particles ranging in size from 0.001 µm (micrometres) to 100 µm. Particles range in chemical composition, size, shape, and physical properties. Sources of particles are either natural sources such as pollen, dust from soil erosion, volcanoes or man-made sources such as soot, flyash and smoke. In Prince George sources of particulate are industries, wood waste burners, road dust, sawmills, motor vehicles, burning of fuels for heating, and some natural sources.

Larger particulate may cause a nuisance or irritation problem but smaller particulate (less than 10 µm) cause the greatest health effect because they are inhaled deep into the lung cavities. Particles in this size range usually come from man-made sources, internal combustion engines, industrial processes, burning, and road dust.

Particulate matter less than 10 micrometres (PM<sub>10</sub>) is measured either non-continuously (once every sixth day when samplers are run for 24 hours at a time) (seven locations in Prince George), or continuously, using a tapered element oscillating microbalance technology (TEOM).

Table 5.1 summarizes the PM<sub>10</sub> data from both the continuous and non-continuous monitors in Prince George for 2000. Annual averages ranged from a low of 9.9 µg/m<sup>3</sup> at Glenview to a maximum of 21.3 µg/m<sup>3</sup> at the CNR site. The maximum 24-hour average, 86 µg/m<sup>3</sup> was recorded at the BC Rail site.

Table 5.1  
2000 Airshed Summary of PM<sub>10</sub>

Station	Annual Average (µg/m <sup>3</sup> )	No. (%) of Daily Values			Maximum Daily (µg/m <sup>3</sup> )	Minimum Daily (µg/m <sup>3</sup> )	No. of Values
		>50µg/m <sup>3</sup>	>100µg/m <sup>3</sup>	>150µg/m <sup>3</sup>			
Plaza	19.0	1 (1.7%)	0	0	65	4	59
Plaza - (TEOM)	18.0	9 (2.4%)	0	0	74	2	362
Lakewood	14.1	0 (0%)	0	0	50	4	60
Van Bien	17.4	2 (3.5%)	0	0	84	4	57
CNR Site	21.3	3 (5.0%)	0	0	61	3	59
BCR Site #2	20.0	13 (3.6%)	0	0	86	1	364
Gladstone	13.7	4 (1.1%)	0	0	55	2	360
Hart-Glenview	9.9	0 (0%)	0	0	49	2	364

B.C. Ambient Air Quality Objectives for PM<sub>10</sub>: ..... Level B  
24 Hour ..... 50 µg/m<sup>3</sup>



## 5.1 PM<sub>10</sub> Continuous Monitors

PM<sub>10</sub> is continuously sampled on top of the Plaza 400 building, Gladstone Elementary, Glenview School and the BCR Warehouse with a TEOM (tapered element oscillating microbalance) 1400a PM<sub>10</sub> sampler. The principle of operation is to draw a constant sample of air through a filter, continuously weighing the filter (every two seconds) and calculating real time (10 minute) mass concentrations. The instrument also calculates 30 minute, one hour, eight hour, and twenty-four hour averages of mass concentration. The initial weight of the filter is compared to the present weight to produce the total mass of the collected particulate.

### 5.1.1 PM<sub>10</sub> Results

Tables 5.2 to 5.5 summarize the continuous PM<sub>10</sub> monitoring at the four monitoring sites in Prince George. Monthly concentrations ranged from 8.1 µg/m<sup>3</sup> (at Glenview on July) to 24.3 µg/m<sup>3</sup> (at Plaza on February). The maximum hourly concentration of 288 µg/m<sup>3</sup> and the maximum 24-hour level (86 µg/m<sup>3</sup>) were both recorded at the BCR site in April.

Table 5.2  
2000 Monthly Summary of Continuous PM<sub>10</sub> Data at Plaza

	Monthly Average (µg/m <sup>3</sup> )	Maximum Hourly (µg/m <sup>3</sup> )	No. of Daily Values		Maximum 24 Hour (µg/m <sup>3</sup> )	Minimum 24 Hour (µg/m <sup>3</sup> )
			>50µg/m <sup>3</sup>	>100µg/m <sup>3</sup>		
<b>January</b>	19.4	112	1	0	62	4.1
<b>February</b>	24.3	128	1	0	60	5.5
<b>March</b>	24.1	174	1	0	60	3.8
<b>April</b>	19.2	175	2	0	71	3.5
<b>May</b>	15.2	77	0	0	31	4.2
<b>June</b>	14.7	77	0	0	42	4.1
<b>July</b>	13.8	91	0	0	36	4.3
<b>August</b>	13.2	82	0	0	29	4.1
<b>September</b>	18.6	108	0	0	42	3.1
<b>October</b>	19.6	130	0	0	61	4.2
<b>November</b>	17.4	104	0	0	45	4.4
<b>December</b>	17.1	82	0	0	50	2.1
<b>Annual</b>	<b>18.0</b>	<b>175</b>	<b>5</b>	<b>0</b>	<b>71</b>	<b>2.1</b>

Table 5.3

2000 Monthly Summary of Continuous PM<sub>10</sub> Data at Gladstone

	Monthly Average ( $\mu\text{g}/\text{m}^3$ )	Maximum Hourly ( $\mu\text{g}/\text{m}^3$ )	No. of Daily Values		Maximum 24 Hour ( $\mu\text{g}/\text{m}^3$ )	Minimum 24 Hour ( $\mu\text{g}/\text{m}^3$ )
			>50 $\mu\text{g}/\text{m}^3$	>100 $\mu\text{g}/\text{m}^3$		
January	17.3	110	1	0	55	3.1
February	17.9	92	1	0	55	3.6
March	16.3	148	0	0	51	2.6
April	13.6	114	1	0	51	3.7
May	11.0	68	0	0	20	3.0
June	10.6	88	0	0	29	3.7
July	9.7	57	0	0	24	3.8
August	9.4	53	0	0	22	3.6
September	11.7	62	0	0	26	2.4
October	14.7	63	0	0	37	3.1
November	15.2	80	0	0	38	2.9
December	15.5	74	0	0	42	0.5
Annual	<b>13.6</b>	<b>148</b>	<b>3</b>	<b>0</b>	<b>55</b>	<b>0.5</b>

Table 5.4

2000 Monthly Summary of Continuous PM<sub>10</sub> Data at BCR

	Monthly Average ( $\mu\text{g}/\text{m}^3$ )	Maximum Hourly ( $\mu\text{g}/\text{m}^3$ )	No. of Daily Values		Maximum 24 Hour ( $\mu\text{g}/\text{m}^3$ )	Minimum 24 Hour ( $\mu\text{g}/\text{m}^3$ )
			>50 $\mu\text{g}/\text{m}^3$	>100 $\mu\text{g}/\text{m}^3$		
January	21.2	137	1	0	74	5.7
February	21.6	110	2	0	56	3.1
March	22.2	192	2	0	59	5.3
April	23.2	288	3	0	86	5.5
May	17.7	135	0	0	45	4.3
June	15.4	154	0	0	44	3.1
July	16.7	117	0	0	43	2.9
August	16.1	164	0	0	37	4.9
September	21.9	159	3	0	71	3.2
October	20.5	142	1	0	56	3.5
November	22.6	123	1	0	63	6.8
December	20.4	171	0	0	65	1.4
Annual	<b>19.9</b>	<b>288</b>	<b>13</b>	<b>0</b>	<b>86</b>	<b>1.4</b>

Table 5.5  
2000 Monthly Summary of Continuous PM<sub>10</sub> Data at Hart Highlands - Glenview

	Monthly Average ( $\mu\text{g}/\text{m}^3$ )	Maximum Hourly ( $\mu\text{g}/\text{m}^3$ )	No. of Daily Values		Maximum 24 Hour ( $\mu\text{g}/\text{m}^3$ )	Minimum 24 Hour ( $\mu\text{g}/\text{m}^3$ )
			>50 $\mu\text{g}/\text{m}^3$	>100 $\mu\text{g}/\text{m}^3$		
<b>January</b>	11.3	73	0	0	42	2.6
<b>February</b>	10.1	78	0	0	30	2.3
<b>March</b>	8.8	48	0	0	27	3.6
<b>April</b>	9.4	76	0	0	27	2.5
<b>May</b>	8.9	48	0	0	18	2.6
<b>June</b>	8.7	97	0	0	23	2.9
<b>July</b>	8.1	153	0	0	17	3.0
<b>August</b>	8.4	62	0	0	18	3.8
<b>September</b>	10.4	45	0	0	24	2.1
<b>October</b>	11.2	85	0	0	41	3.5
<b>November</b>	12.5	81	0	0	43	2.9
<b>December</b>	11.1	95	0	0	49	2.0
<b>Annual</b>	<b>9.9</b>	<b>153</b>	<b>0</b>	<b>0</b>	<b>49</b>	<b>2.0</b>

### 5.1.2 PM<sub>10</sub> Annual Trends (Continuous Monitors)

In 2000, of the four continuous TEOM monitors only the BCR site recorded a lower annual average than the previous year. The increase at the other three stations ranged from 0.2  $\mu\text{g}/\text{m}^3$  to 1.0  $\mu\text{g}/\text{m}^3$ . The TEOM monitor at Plaza was sent for upgrades at the end of 1993. The annual average of the TEOM has usually been lower (1993 and 1994) than the discontinuous monitor and 2000 continued that trend with a difference of about 1.0  $\mu\text{g}/\text{m}^3$ .

The TEOMs located at Gladstone School, the BC Rail Warehouse, and Glenview School (Tables 5.7, 5.8, and 5.9 respectively) have been operating since late December 1995, October 1996, and June 1998 respectively, but have not operated long enough for any trend discussion. However, of the three sites, only the BCR site annual average decreased in 2000 from the previous year.

Table 5.6  
Annual Trend Summary of the Continuous PM<sub>10</sub> Data at Plaza

Year	Annual Average (µg/m <sup>3</sup> )	Maximum Hourly Average (µg/m <sup>3</sup> )	No. (%) of Days > 50 µg/m <sup>3</sup>	No. (%) of Days > 100 µg/m <sup>3</sup>	Maximum Daily Average (µg/m <sup>3</sup> )	Hours Instrument Operated
1992	21.6	488	18 (6.1%)	0 (0%)	80	7,083
1993	22.3	171	11 (4.5%)	0 (0%)	75	5,828
1994*	22.3	284	19 (5.6%)	4 (1.2%)	117	8,162
1995	24.2	291	33 (9.1%)	2 (0.6%)	108	8,686
1996	20.5	373	10 (2.8%)	3 (0.8%)	152	8,567
1997	20.2	208	13 (3.6%)	0 (0%)	76	8,719
1998	22.7	319	20 (5.5%)	2 (0.5%)	114	8,719
1999	17.8	211	10 (2.7%)	0 (0%)	98	8,609
2000	18.0	175	9 (2.5%)	0 (0%)	71	8,695

\* Monitor returned in January after upgrades

Table 5.7  
Annual Trend Summary of the Continuous PM<sub>10</sub> Data at Gladstone

Year	Annual Average (µg/m <sup>3</sup> )	Maximum Hourly Average (µg/m <sup>3</sup> )	No. (%) of Days > 50 µg/m <sup>3</sup>	No. (%) of Days > 100 µg/m <sup>3</sup>	Maximum Daily Average (µg/m <sup>3</sup> )	Hours Instrument Operated
1996	15.0	253	7 (1.9%)	1 (0.3%)	102	8,688
1997	14.8	155	7 (1.9%)	0 (0%)	61	8,673
1998	17.2	207	7 (1.9%)	0 (0%)	74	8,642
1999	12.7	127	0 0	0 (0%)	47	8,663
2000	13.7	148	4 (1.1%)	0 (0%)	55	8,686

Table 5.8  
Annual Trend Summary of the Continuous PM<sub>10</sub> Data at BCR

Year	Annual Average (µg/m <sup>3</sup> )	Maximum Hourly Average (µg/m <sup>3</sup> )	No. (%) of Days > 50 µg/m <sup>3</sup>	No. (%) of Days > 100 µg/m <sup>3</sup>	Maximum Daily Average (µg/m <sup>3</sup> )	Hours Instrument Operated
1996*	21.3	125	5 (1.9%)	0 (0.3%)	69	2,257
1997	24.2	401	32 (8.9%)	2 (0.6%)	107	8,673
1998	29.1	582	49 (13.4%)	4 (1.1%)	127	8,702
1999	21.3	380	19 (5.2%)	0 (0%)	82	8,718
2000	20.0	288	13 (3.6%)	0 (0%)	86	8,739

\* analyzer installed in October 1996

Table 5.9  
Annual Trend Summary of the Continuous PM<sub>10</sub> Data at Glenview

Year	Annual Average (µg/m <sup>3</sup> )	Maximum Hourly Average (µg/m <sup>3</sup> )	No. (%) of Days > 50 µg/m <sup>3</sup>	No. (%) of Days > 100 µg/m <sup>3</sup>	Maximum Daily Average (µg/m <sup>3</sup> )	Hours Instrument Operated
1998	11.0	117	0 (0%)	0 (0%)	38	4,852*
1999	9.2	221	0 (0%)	0 (0%)	34	8,670
2000	9.9	153	0 (0%)	0 (0%)	49	8,731

Note: Monitoring started June 1998

## 5.2 PM<sub>10</sub> Annual Trends (Non-continuous Monitors)

Unlike the continuous TEOM monitors, all non-continuous stations showed lower annual average PM<sub>10</sub> levels in 2000 compared to the previous year, and were the lowest since monitoring began. Tables 5.10 - 5.16 show the annual summary of PM<sub>10</sub> data from all the monitoring sites.

Table 5.10  
Annual Trend Summary of Non-Continuous PM<sub>10</sub> Data at Plaza

Year	Annual Average (µg/m <sup>3</sup> )	No. (%) of Daily Values > 50 µg/m <sup>3</sup>	No. (%) of Daily Values > 100 µg/m <sup>3</sup>	Maximum Daily Value (µg/m <sup>3</sup> )	Number of Samples
1990	22.8	2 (7.7%)	0	90	26
1991	29.4	7 (11.7%)	1 (1.7%)	217	60
1992	28.6	6 (10.2%)	0	92	59
1993	30.2	10 (16.7%)	0	92	60
1994	24.1	6 (10.0%)	0	85	61
1995	23.6	4 (6.6%)	0	85	61
1996	21.9	2 (3.3%)	0	61	61
1997	20.8	1 (1.7%)	0	56	61
1998	26.9	5 (8.2%)	1 (1.6%)	111	61
1999	21.4	2 (3.4%)	0	73	59
2000	19.0	1 (1.7%)	0	65	59

Table 5.11  
Annual Trend Summary of Non-Continuous PM<sub>10</sub> Data at Van Bien

Year	Annual Average (µg/m <sup>3</sup> )	No. (%) of Daily Values > 50 µg/m <sup>3</sup>	No. (%) of Daily Values > 100 µg/m <sup>3</sup>	Maximum Daily Value (µg/m <sup>3</sup> )	Number of Samples
1990	30.0	7 (26.9%)	0	62	26
1991	27.6	8 (13.6%)	0	84	59
1992	27.0	7 (12.5%)	0	89	56
1993	27.9	6 (10.2%)	0	99	59
1994	21.8	3 (6.3%)	0	69	48
1995	25.2	6 (10.2%)	1 (1.7%)	106	59

<b>1996</b>	23.4	3 (7.3%)	0	58	41
<b>1997</b>	20.8	3 (5.3%)	0	74	57
<b>1998</b>	28.3	6 (10.9%)	0	91	55
<b>1999</b>	22.7	3 (5.2%)	0	56	58
<b>2000</b>	17.4	2 (3.5%)	0	84	57

Table 5.12  
Annual Trend Summary of Non-Continuous PM<sub>10</sub> Data at Lakewood

<b>Year</b>	<b>Annual Average (<math>\mu\text{g}/\text{m}^3</math>)</b>	<b>No. (%) of Daily Values &gt; 50 <math>\mu\text{g}/\text{m}^3</math></b>	<b>No. (%) of Daily Values &gt; 100 <math>\mu\text{g}/\text{m}^3</math></b>	<b>Maximum Daily Value (<math>\mu\text{g}/\text{m}^3</math>)</b>	<b>Number of Samples</b>
<b>1990</b>	17.4	0	0	40	24*
<b>1991</b>	21.8	5 (8.5%)	0	89	59
<b>1992</b>	24.8	7 (11.9%)	0	74	59
<b>1993</b>	27.3	4 (7.6%)	1 (1.9%)	129	53
<b>1994</b>	18.9	1 (1.7%)	0	56	58
<b>1995</b>	18.7	4 (7.4%)	0	63	54
<b>1996</b>	16.3	0	0	50	59
<b>1997</b>	15.9	1 (1.7%)	0	51	59
<b>1998</b>	19.6	1 (1.9%)	0	61	54
<b>1999</b>	17.4	1 (1.8%)	0	74	56
<b>2000</b>	14.1	0	0	50	60

\* Note: High volume samplers converted to PM<sub>10</sub> (August 1, 1990). Therefore PM<sub>10</sub> data is for a 5 month duration.

Table 5.13  
Annual Trend Summary of Non-Continuous PM<sub>10</sub> Data at the CNR Site

<b>Year</b>	<b>Annual Average (<math>\mu\text{g}/\text{m}^3</math>)</b>	<b>No. (%) of Daily Values &gt; 50 <math>\mu\text{g}/\text{m}^3</math></b>	<b>No. (%) of Daily Values &gt; 100 <math>\mu\text{g}/\text{m}^3</math></b>	<b>Maximum Daily Value (<math>\mu\text{g}/\text{m}^3</math>)</b>	<b>Number of Samples</b>
<b>1991</b>	30.9	5 (13.2%)	2 (5.3%)	124	38
<b>1992</b>	28.3	6 (10.3%)	1 (1.7%)	103	58
<b>1993</b>	33.3	14 (23.0%)	1 (1.6%)	104	61
<b>1994</b>	25.2	8 (13.6%)	0	75	59
<b>1995</b>	27.5	8 (13.3%)	1 (1.7%)	110	60

<b>1996</b>	28.1	5 (8.3%)	1 (1.7%)	110	60
<b>1997</b>	22.8	2 (3.6%)	0	65	55
<b>1998</b>	34.0	8 (13.8%)	2 (3.4%)	124	58
<b>1999</b>	28.0	8 (13.6%)	1 (1.7%)	106	59
<b>2000</b>	21.3	3 (5.1%)	0	61	59

Table 5.14

Annual Trend Summary of Non-Continuous PM<sub>10</sub> Data at the BCR Site #1 (Discontinued)

<b>Year</b>	<b>Annual Average (µg/m<sup>3</sup>)</b>	<b>No. (%) of Daily Values &gt; 50 µg/m<sup>3</sup></b>	<b>No. (%) of Daily Values &gt; 100 µg/m<sup>3</sup></b>	<b>Maximum Daily Value (µg/m<sup>3</sup>)</b>	<b>Number of Samples</b>
<b>1990</b>	41.6	8 (21.6%)	3 (8.1%)	132	37
<b>1991</b>	49.3	8 (50.0%)	1 ((6.3%)	122	16*
<b>1992</b>	31.3	2 (9.1%)	0	76	22**
<b>1993</b>	43.2	17 (28.8%)	2 (3.4%)	147	59
<b>1994</b>	38.4	12 (21.1%)	2 (3.5%)	143	57
<b>1995</b>	40.9	16 (31.4%)	3 (5.9%)	181	51
<b>1996</b>	43.9	9(24.3%)	2 (5.4%)	153	37***

\* Note: Station relocated to CNR site May 6, 1991.

\*\* Note: Monitoring started July 29, 1992.

\*\*\*Note: Station closed August 20 1996 and relocated to BCR #2 site

Table 5.15

Annual Trend Summary of Non-Continuous PM<sub>10</sub> Data at the Current BCR Site #2

<b>Year</b>	<b>Annual Average (µg/m<sup>3</sup>)</b>	<b>No. (%) of Daily Values &gt; 50 µg/m<sup>3</sup></b>	<b>No. (%) of Daily Values &gt; 100 µg/m<sup>3</sup></b>	<b>Maximum Daily Value (µg/m<sup>3</sup>)</b>	<b>Number of Samples</b>
<b>1996</b>	31.8	4 (11.4%)	0	79	35*
<b>1997</b>	26.0	4 (7.1%)	0	75	61
<b>1998</b>	31.7	10(16.4%)	0	92	61

\* Note: Monitoring started June 2, 1996.

Table 5.16

Annual Trend Summary of Non-Continuous PM<sub>10</sub> Data at Gladstone (Discontinued)

<b>Year</b>	<b>Annual Average (µg/m<sup>3</sup>)</b>	<b>No. (%) of Daily Values &gt; 50 µg/m<sup>3</sup></b>	<b>No. (%) of Daily Values &gt; 100 µg/m<sup>3</sup></b>	<b>Maximum Daily Value (µg/m<sup>3</sup>)</b>	<b>Number of Samples</b>
<b>1992</b>	19.5	1 (3.2%)	0	54	31*
<b>1993</b>	25.8	5 (8.8%)	1 (1.8%)	122	57



<b>1994</b>	16.8	0	0	50	60
<b>1995</b>	16.7	2 (3.3%)	0	63	60
<b>1996</b>	15.4	0	0	49	59
<b>1997</b>	15.3	1 (1.7%)	0	57	61
<b>1998</b>	20.2	2 (3.3%)	0	57	61

\* Note: Monitoring started June 5, 1992.

#### 5.4 PM<sub>2.5</sub> Results

The Ministry operates one continuous PM<sub>2.5</sub> (particulate matter less than 2.5 micrometres) monitor, which is located at the Plaza site. The PM<sub>2.5</sub> monitor operates in the same manner as the TEOM PM<sub>10</sub> monitors. The only difference is the design of the sample inlet which restricts particles greater than 2.5µm being sampled. Similar to the PM<sub>10</sub> data, the lowest monthly average occurred in August. The maximum PM<sub>2.5</sub> monthly average occurred in February, the same month of the maximum PM<sub>10</sub> monthly average.

Table 5.17  
2000 Monthly Summary of Continuous PM<sub>2.5</sub> Data at Plaza

	Monthly Average (µg/m <sup>3</sup> )	Maximum Hourly (µg/m <sup>3</sup> )	No. of Daily Values		Maximum 24 Hour (µg/m <sup>3</sup> )	Minimum 24 Hour (µg/m <sup>3</sup> )
			>25µg/m <sup>3</sup>	>50µg/m <sup>3</sup>		
<b>January</b>	16.7	83	6	0	51.4	5.5
<b>February</b>	19.3	98	7	0	50.3	3.5
<b>March</b>	13.1	66	2	0	33.2	2.3
<b>April</b>	11.7	77	2	0	39.3	3.1
<b>May</b>	10.0	54	0	0	20.3	3.5
<b>June</b>	10.3	59	1	0	29.5	2.9
<b>July</b>	9.9	76	1	0	27.7	3.5
<b>August</b>	8.9	56	0	0	18.5	4.1
<b>September</b>	11.4	78	1	0	28.0	2.5
<b>October</b>	13.4	72	4	0	36.5	2.4
<b>November</b>	13.0	45	1	0	31.8	2.8
<b>December</b>	14.0	65	3	0	40.7	1.3
<b>Annual</b>	<b>12.6</b>	<b>98</b>	<b>27</b>	<b>0</b>	<b>51.4</b>	<b>1.3</b>

#### 5.5 PM<sub>2.5</sub> Annual Trend (Non-continuous Monitor)

The Ministry operates one non-continuous PM<sub>2.5</sub> monitor, which is located at the Plaza site. PM<sub>2.5</sub> is measured by drawing air through a size selective inlet prior to collection on a teflon fibre filter at a rate of about 0.017 cubic metres per minute. The sampler is run for 24 hours at a time on a 6 day cycle (i.e., once every sixth day). Table 5.18 shows the annual trend summary of PM<sub>2.5</sub> data for the last seven years at Plaza, which shows a steady decrease in levels from 1995 to 1998, but a sharp increase in 2000.

Table 5.18  
Annual Trend Summary of Non-Continuous PM<sub>2.5</sub> Data at Plaza

Year	Annual Average (µg/m <sup>3</sup> )	No. (%) of Daily Values > 25 µg/m <sup>3</sup>	No. (%) of Daily Values > 50 µg/m <sup>3</sup>	Maximum Daily Value (µg/m <sup>3</sup> )	Number of Samples
------	--	---	---	---	-------------------

<b>1994*</b>	13.5	1 (4.5%)	1 (4.5%)	52	22
<b>1995</b>	13.3	5 (8.8%)	1 (1.8%)	54	57
<b>1996</b>	12.7	4 (6.7%)	0	40	60
<b>1997</b>	12.3	6 (9/8%)	0	43	61
<b>1998</b>	11.4	5 (8.3%)	1 (1.7%)	52	60
<b>1999</b>	9.9	2 (3.6%)	0	38	55
<b>2000</b>	12.0	6 (10.5%)	1 (1.8%)	52	57

\* Instrument installed August 1994