

River Bulrush (*Bolboschoenus fluviatilis*)

Cyperaceae (Sedge Family)

RANGE

- Widespread in southern North America, from Alberta to the Maritime provinces, south to Maryland and west to Oregon (USA)
- In British Columbia, known from Osoyoos Lake area in the south Okanagan Valley and from one occurrence near Hesquiat on Vancouver Island




Figure 1 Thompson Okanagan Region distribution of *Bolboschoenus fluviatilis* (BC CDC 2013)

HABITAT

- Open marshes alongside rivers, oxbows and lakeshores in the Bunchgrass and Coastal Western Hemlock Biogeoclimatic Zones
- May also be found in fresh to brackish marshes in quiet water, sometimes >1 m deep
- Common associates include hard-stemmed bulrush (*Schoenoplectus acutus*), wetland grasses and sedges (*Carex* spp.)



Figure 2 Marsh habitat north of Osoyoos Lake, B.C.



Figure 3 Close-up of triangular stems on sterile plant

LIFE HISTORY

- Perennial species that mainly grows vegetatively by rhizomes, although colonization of open areas by seed may occur
- Flowers in late spring and achenes mature in summer, although flowering stems are uncommon and patches may be completely vegetative




Figure 4 Illustration of *Bolboschoenus fluviatilis* by Jeanne R. Janish (Hitchcock et al. 1969)

***Bolboschoenus fluviatilis* (continued)**

DESCRIPTION

General

- Tall, usually from 1 to 2 m, perennial wetland plant with sharply triangular stems
- Grows from tubers that are present along underground rhizomes, sometimes forming extensive, dense patches
- Flower production within patches is uncommon; patches are nearly completely vegetative

Leaves

- Several long, flat leaves, up to 40 cm long and from 7 to 18 mm wide

Flowers

- Flowers produced at tops of stems and surrounded by a series of well developed leaf-like bracts
- Numerous flowers clustered together forming egg or spindle shaped, usually pointed spikelets
- Most spikelets borne singly or in small clusters at ends of 3 to 8 cm long branches that flare out from top of the stem; other spikelets sometimes clustered in a compact group at base of bracts

Fruits

- Three-sided achenes produced by each flower

IDENTIFICATION TIPS

- Characterized by sharply triangular stems, distinctly triangular achenes, and open inflorescences with clusters of spikelets extended away from the stem
- Easily confused with seacoast bulrush (*B. maritimus* var. *paludosus*), a more common species also with triangular stems and clusters of spikelets at tops of the stems, some of which may also be extended away from stem on short branches; seacoast bulrush is usually smaller in stature with narrower leaves, two-sided or lens-shaped achenes, and frequently produces sessile spikelets that are tightly clumped at the ends of the stem; this species is also most common in wetland habitats with higher salinity, particularly along the edges of saline depressions and ponds
- Readily separated from hard-stemmed bulrush (*Schoenoplectus acutus*) and American bulrush (*S. pungens* var. *longispicatus*) by the numerous leaves along its stems and the series of leaf-like bracts at the base of the flowers
- Also readily separated with small-flowered bulrush (*Scirpus microcarpus*), which has well developed stem leaves but its spikelets are much smaller and always at ends of small branches



Figure 5 Close up of single flowering plant in otherwise sterile patch

GENERAL THREATS AND GUIDANCE

- **Avoid development in areas with known occurrences of *Bolboschoenus fluviatilis* through project relocation or redesign**
- Protect open riverine, oxbow and lakeshore marsh habitats from disturbance and development, including exclusion of livestock through fencing, and consider restoration including invasive plant removal following professional advice
- Because patches are often mostly sterile, careful searching for flowering stems is essential for identification
- Follow provincial methods for when and how to conduct plant species at risk surveys
- Follow provincial policy and guidance on how to avoid, minimize, restore and offset impacts to plant species at risk and their habitats
- Report any sightings to the B.C. Conservation Data Centre (cdccdata@gov.bc.ca) and FLNR Ecosystems Section (josie.symonds@gov.bc.ca)

REFERENCES

- B.C. Conservation Data Centre. 2013. <http://a100.gov.bc.ca/pub/eswp/>
- Douglas et al. (editors). 2001. *Illustrated Flora of British Columbia*. Vol. 6. B.C. Min. Environ., Lands and Parks, and Min. For., Victoria, B.C.
- Hitchcock et al.. 1969. *Vascular Plants of the Pacific Northwest*, Univ. Washington Press, Seattle, W.A.
- NatureServe. 2011. <http://www.NatureServe.org/explorer>

ACKNOWLEDGEMENTS

Bolboschoenus fluviatilis Plant Species at Risk Fact Sheet developed by Josie Symonds, based on content produced under contract by Terry McIntosh and review comments from Brenda Costanzo and Orville Dyer