

## Honeybee Disease Detection

Recognizing disease symptoms is an essential part of beekeeping management. Early detection allows for prompt remedial action and helps to prevent serious disease outbreak and economic losses. Please consult additional references for information about control products and methods.

This factsheet offers information about field symptoms and laboratory diagnoses of the most common honeybee diseases. For information about Varroa and tracheal mites, please refer to **Bulletin #219 – Tracheal Mites in Honeybee Colonies** and **#222 – Varroa Mite Detection Methods**.

### *American Foulbrood (AFB)*

#### *Field Diagnosis*

- Hive may show less than normal bee flight with dead bees on the bottom board. The colony may appear weak after opening the hive.
- Capped brood is uneven with puncture holes in the caps of brood cells.
- Colonies with heavy infestation often display irritable behaviour.
- AFB has a distinct “foul” odour that can help in alerting the beekeeper to a disease problem.
- With a toothpick, lift punctured cap and remove content of brood cell. The larval remnant may be a light brown mass sunk onto the bottom side of the brood cell. If the mass is **ropy** when withdrawing the toothpick from the cell, there is a strong indication of American Foulbrood Disease (AFB).
- Place toothpick in a small plastic bag or plastic wrap and mail to the Apiculture office for confirmation.
- Over time, the larval remains in the cell will dry and harden into a dark brown leathery scale on the bottom side of the brood cell. A single scale contains millions of spores that remain viable for decades. Bees cannot remove scales from cells.
- AFB scales can be readily detected in the field by holding the brood frame at an angle of approximately 15 degrees. Scales should be easily visible.



View of brood comb held at approximately a 15 – 20 degree angle. Note the darkened bottom sides of the cells containing scale. Capped brood is unevenly distributed (shotgun appearance). Cappings are often punctured by adult bees.

### **Laboratory Diagnosis**

- AFB is caused by *Paenibacillus larvae*, a spore-forming bacterium.
- A microscope slide can be prepared by dissolving a small part of an AFB scale. Stir the scale with a toothpick in a droplet of water placed on a slide and apply a cover slip.
- Under 400X magnification, the AFB spores are visible and characterized by their slightly oblong, uniform size and shape. The spores “jiggle” in a characteristic Brownian movement.
- *P. larvae* is competitive and does not tolerate growth of other bacteria in the parasitized bee larva. As a result, most microscopic slides will show a predominance of *P. larvae* spores. This is not always the case with poor samples or those left in the collection bag for too long.

### **Control and Treatment**

- Antibiotic-resistant AFB (r-AFB) has become established in BC. Antibiotics must be used for treatment purposes only. Do not use antibiotics as a prophylactic (=preventive) measure.
- Become thoroughly familiar with visual detection of brood diseases.
- Inspect regularly, especially when disease has been reported in the area or after the colony has been placed in crop pollination.
- For frames with suspect signs of brood disease, collect a sample and submit to the Apiculture office.
- When AFB has been confirmed, kill the bees and burn all the equipment. For alternative options, contact your Apiary Inspector or Apiculture Office.
- Use antibiotics only as recommended. Never use the product after its expiry date, and follow preparation instructions carefully (refer to **Bulletin #204**).
- Reduce the exchange of hive equipment between hives and apiaries.
- Replace 20% of all brood frames each year. Eventually, no brood frame is older than five years.
- Don't open-barrel feed or leave used hive equipment exposed to foraging bees.
- Apply hygienic management practices, including clean clothing, hive tools, and gloves.

### **European Foulbrood (EFB)**

#### **Field Diagnosis**

- European Foulbrood is less serious than AFB. EFB shows up when the colonies have been under stress due to other diseases, colonies nearby, poor management and weather.
- EFB is easily controlled with standard antibiotic treatments.
- EFB affects bee brood similarly to AFB except that the disease affects open brood (the larvae die before they are capped).
- Affected cells show discoloured larvae, often in twisted positions, with visible tracheal tubes.
- The brood has a sour odour, distinctly different from AFB.
- EFB scales are easily removed from the cell (compared to AFB scales).
- When scales are detected, collect cell scrapings and submit for laboratory examination.

#### **Laboratory Diagnosis**

- EFB is caused by *Melissococcus pluton*, but the secondary invader *Pseudobacillus alvei* is mostly observed when samples are examined microscopically.
- Samples are prepared the same way as AFB samples.
- At 400X, *P. alvei* is readily identified by its long spindle shaped spores.
- The spores do not jiggle but float by in the solution.
- Unlike AFB, EFB microscopic samples generally display a wide variety of microbes.

#### **Control and Treatment**

- Inspect brood frames regularly and be familiar with field symptoms.
- Remove all frames with significant numbers of affected cells.
- Spray or sprinkle antibiotics (oxytetracycline) dissolved in 250 ml of sugar syrup over the colony every 3-4 days for 10 days. Refer to **Factsheet #204** for handling of antibiotics.
- Requeening provides a distinct break in the brood cycle of the colony, allowing the bees to clean up existing disease. It may also provide new bees with stronger cleaning impulse and be less susceptible.
- Minimize robbing by preventing sugar spillage. Do not barrel feed.
- Apply hygienic management practices. Clean hive tools, smoker and gloves after inspection of each apiary. Clean clothes regularly.
- Replace brood frames after 5 years.

## **Chalkbrood Disease**

### **Field Diagnosis**

- Chalkbrood disease affects bee larvae and is caused by the fungus *Ascosphaera apis*.
- Chalkbrood incidence increases in the fall and spring. Mummified larvae in front of the hive and on the bottom board are easily detected. Mummies on the bottom board may not necessarily indicate a serious problem, but confirm hygienic bee behaviour.
- Remove heavily mummified comb and scrape bottom board to reduce source of infection.
- There is no control product available. High incidence of Chalkbrood mostly indicates poor hygienic behaviour and stress due to weather, poor management or diseases.
- When there is a persistent Chalkbrood problem, replace the queen.

### **Laboratory Diagnosis**

- Mummified larvae are generally white in colour. The mycelium of the fungus infiltrates the larval tissue that eventually hardens. The white colour is the result of asexual reproduction while sexual reproduction produces black or grey coloured mummies.

## **Sacbrood**

- Sacbrood is caused by a virus. The infected larva dies and the tissue disintegrates into a brown watery solution held by the larval outer skin. The skin sac can be removed intact from the cell. The cell is often uncapped but may also be closed and the cap punctured similarly to AFB cells.
- Sacbrood occurs when colonies have been under stress. Bees generally clear up the disease and remove affected brood themselves.
- There are no control products available.
- Positive identification requires PCR diagnostic methodologies.

## **Nosema**

### **Field Diagnosis**

- Nosema disease is caused by a spore-forming microsporidian fungus belonging to the genus *Nosema*. Two species, *N. apis* and *N. ceranae* have been identified in British Columbia.
- Nosema incidence in honeybee colonies peaks in early spring and early fall.
- The disease only affects adult bees by parasitizing their midgut. Adult bees have difficulty with controlling their fecal discharge. In heavy infestations, hive bodies are smeared with fecal deposits.
- The disease is often not detected because affected bees are either inside the colony (in winter) or in the field, where they die.
- Nosema impairs the digestive process and causes bee starvation.
- Nosema is often confused with dysentery caused by a virus which produces similar symptoms.

### **Laboratory Diagnosis.**

- For Nosema detection and confirmation, adult bees must be examined microscopically.  
Standard detection method: collect 25 dead bees and place in mortar with 25 ml of water (i.e. 1 ml water for every adult bee). Grind up and collect one droplet of solution and place on slide. Cover with coverslip.
- Examine at 100X magnification. Nosema spores are large, oblong and highly uniform in shape.
- For determining the level of infestation, a haemocytometer is used to calculate spores per bee.
- To submit a sample for analysis, collect at least 50 bees in tissue paper (no plastic) or paper bag, freeze for several days and mail to the Apiculture office..

### **Control and Treatment**

- Nosema disease incidence increases when colonies are under stress and poorly managed. The condition is often exacerbated when there is moisture build up and poor air circulation in the hive.
- The antibiotic fumagillin, sold under the trade name Fumadil B is effective. The best natural defense is a strong healthy colony with a prolific queen and sufficient food stores. Rapid replacement of adult bees will minimize or eliminate the disease.
- Higher Nosema incidence has often been reported with tracheal mite infestations.

## **Various “Disease-like” Brood Comb Conditions**

- **Wax Moth**

Larvae of several local species (Lesser Wax Moth, Dried Fruit Moth) damage unoccupied combs as they tunnel through, leaving webbing and droppings. If comb is only lightly damaged, the bees will clean and repair the comb. The Greater Wax Moth causes more extensive damage, but it has only been incidentally reported in southern parts of BC.

- **Bee Starvation**

A patch of dead adult bees, with their heads fully inserted into cells, is a telltale sign of starvation. Wintered colonies may still have food reserves that were not accessible to the bees. Comb with a patch of dead bees can be placed in a new colony for cleanup, providing it is free of disease.

- **Drone Brood in Worker Cells**

Multiple eggs per cell indicate laying workers. This occurs when the colony is queenless or when the queen is failing. When drone brood emerges from worker brood comb, the queen is a “drone layer” by laying unfertilized eggs. In either case, the colony needs a new queen. A new queen can be purchased or raised from other healthy colonies.

- **Chilled Brood**

Bee brood may die when sudden cold spells occur in spring. This happens more often in smaller colonies where the adult bee population is not large enough to maintain brood temperature. Placement of the frame in the center of the colony will force the bees to remove and clean all the cells.

- **Moldy Pollen**

Stored pollen in comb is preserved when covered by honey. If the honey is removed and bees are no longer present, the pollen will become moldy. No disease is involved and the frame can be placed back into a strong colony for cleaning and use.

---

**Plant & Animal Health Branch**  
1767 Angus Campbell Road  
Abbotsford, BC V3G 2M3

**Toll Free: 1-800-661-9903**  
Phone: 604-556-3003  
Fax: 604-556-3010