

Elm Seed and “Tuxedo” Bugs

New Invasive Seed Bugs in British Columbia

Introduction

The elm seed and tuxedo bugs are native to Europe and the Mediterranean region. They were first reported in Canada in Kelowna, British Columbia in 2016. There were more reports of elm seed bug infestations by homeowners in Kelowna in 2017. The seed bugs are not agricultural pests but can be a nuisance in high numbers because they enter homes and businesses. Elm seed bugs emit unpleasant odours when crushed and their fecal droppings on structures such as doors and windows can be unsightly. Elm seed and tuxedo bugs do not bite people.



Figure 1. Adult elm seed bug. Note triangle enclosed in a rusty-coloured rectangle. Photo credit, Ward Strong, Kalamalka Forestry Centre, Vernon, B.C.

Identification

Elm seed bug: Adults are 6.5 - 7 mm (about 1/3 inch) long, black and rusty red colour with black triangle bordered by a rusty coloured triangle on the back (Figure 1). The outer margins of the abdomen have contrasting black and white bands. Immature stages (nymphs) have a black head and a red abdomen; older nymphs have wing buds and two black spots on the back of the abdomen (Figure 3).

Tuxedo bug: Adults are about 5 - 6 mm (1/4 inch) long, dark brown to black, with a triangle outlined in white, distinct paired white spots at the top and lower part of the body and a white spot at the tip of the body (Figure 2).



Figure 2. Adult tuxedo bug. Note distinctive white stripes on the sides of the triangle and white spots on the upper and lower parts of the body. Photo credit, Ward Strong, Kalamalka Forestry Centre, Vernon, B.C.

Life cycle

The life cycle of elm and tuxedo seed bugs in British Columbia has not been determined.

Elm seed bug: In Europe and the United States, elm seed bug has one generation per year. Elm seed bugs overwinter as adults in and around structures and emerge in the spring to lay eggs on elms. Nymphs feed on elm seeds from May-June and adults are present in the summer. There are five immature stages and because of the long egg laying period, life stages may overlap with adults and immature stages being present in June and July. In Interior British Columbia, adults were collected in June in 2016. In 2017, mainly nymphs and a few adults were present at three infested residences in Kelowna in June.

Tuxedo bug: Tuxedo bugs overwinter as adults, which lay eggs in the ground or soil in early spring. Nymphs feed on developing seeds of plants in the mint family. There is one generation per year in the United States, two in England, and three in Russia. In Idaho, adults are active in early to mid-spring and from mid-summer into late fall. In British Columbia, adults were present in August in 2016.

Hosts and Damage

Elm seed bugs feed on elm seeds and leaves but do not cause much damage to the trees. Tuxedo bugs feed on plants in the mint family (e.g. black horehound, lambs ears, white mullein). Elm seed and tuxedo bugs are not known to feed on agricultural crops. They do not pose a health risk to humans or pets and do not bite. They are a nuisance when they invade homes and structures in large numbers. This can be a source of discomfort and anxiety for homeowners. Elm seed bugs stink when crushed and their fecal droppings on structures are an eyesore.



Figure 4. Aggregation of tuxedo bugs in a home.



Figure 3. Top; Elm seed bug adult. Note triangle with rust-coloured borders and light bands on the outer margins of the body. Middle and bottom, elm seed bug nymphs.

Control

Prevent entry into homes or buildings by sealing off any access points in windows, doors and screens.

Vacuum bugs in and around homes. For large numbers, use a shop vacuum with 1-2 inches of soapy water in the bottom to drown the bugs.

Remove volunteer elm trees. Where practical prune elm trees to reduce food source for elm seed bugs.

Clean up elm seeds and debris around the home and structures.

Use sticky traps for trapping bugs around window sills.

Inspect firewood for overwintering adults before bringing into the home.

The use of insecticides for controlling seed bugs in the home is not recommended.

For high numbers, treating immature stages outside the home with a barrier spray along foundations, patios, doors and windows will help prevent bugs from entering homes.

Products registered for home use containing permethrin, carbaryl and malathion will provide control.

Homeowners can hire a commercial pesticide applicator for more control options.

Acknowledgement

I would like to thank Dr Michael Schwartz, Agriculture and Agri-Food Canada, Insect Identification Service, Ottawa for identification of species.

Susanna Acheampong PhD
Ministry of Agriculture
1690 Powick Road
Kelowna, BC V1X 7G5
Phone: 250 861-7681
Email: Susanna.Acheampong@gov.bc.ca

December 2017



Figure 5. Top; Tuxedo bugs overwintering under the bark of a willow tree. Photo credit, Robert Geismayr, Kelowna. Middle and bottom, elm seed bugs on a sticky trap.

References

Acheampong S., Strong W.B., Schwartz M.D., Higgins R.J., Thurston M.A., Walker E.J., Roberts J. 2016. First Canadian records for two invasive seed-feeding bugs, *Arocatus melanocephalus* (Fabricius, 1798) and *Raglius alboacuminatus* (Goeze, 1778), and a range extension for a third species, *Rhyparochromus vulgaris* (Schilling, 1829) (Hemiptera: Heteroptera) J. Ent. Soc. B.C. 113: 74-78.

<https://journal.entsocbc.ca/index.php/journal/article/viewFile/940/992>

Bechinski, E. J., J. Barbour, and F. Merickel. 2012. Elm seed bug, *Arocatus melanocephalus*, a new bug in Idaho. <http://vickisgardentips.com/wpcontent/uploads/2013/01/Elm-seed-bug-handout.pdf>

Bechinski, E. J. and F. Merickel. 2007. Tuxedo bug. A new home-invading insect in Idaho. University of Idaho, College of Agricultural and Life Sciences, Extension publication CIS 1147. <http://www.cals.uidaho.edu/edcomm/pdf/cis/cis1147.pdf>

Davis, R. S. 2017. Elm seed bug. Utah State University Extension publication ENT-186-17. http://digitalcommons.usu.edu/cgi/viewcontent.cgi?article=2788&context=extension_curall

Idaho State Department of Agriculture. 2013. Elm seed bug, *Arocatus melanocephalus*: an exotic invasive pest new to the U.S. http://extension.oregonstate.edu/malheur/sites/default/files/spring_2013_esb_fact_sheet.pdf

Stokes, B.S., E. J. Bechinski and P. J. Castrovillo. 2016. Managing elm seed bugs around your home. University of Idaho Extension publication CIS 1223. <http://www.cals.uidaho.edu/edcomm/pdf/CIS/CIS1223.pdf>
