

Protocol for

Evaluating the Health of Wetlands

(Wetland Management Routine Effectiveness Evaluation)

Version 2.1

January 2021



Wetland Management Routine Effectiveness Evaluation – January 2021

Citation:

Fletcher, N.F., Tripp, D.B., Hansen, P.L., Nordin, L.J., Porter, M., and Morgan, D. 2021. *Protocol for the Wetland Health Management Routine Effectiveness Evaluation*. Forest and Range Evaluation Program, B.C. Ministry of Forests, Lands, Natural Resources Operations and Rural Development, Victoria, B.C.

Prepared by:

Neil Fletcher, BC Wildlife Federation
Derek Tripp, Tripp & Associates Consulting Ltd.
Paul Hansen, Ecological Solutions Group LLC
Lisa Nordin, B.C. Ministry of Forests, Lands, Natural Resources Operations and Rural Development
Marc Porter, ESSA Technologies
Don Morgan, B.C. Ministry of Environment & Climate Change Strategy
Maureen Nadeau, BC Wildlife Federation
Jason Jobin, BC Wildlife Federation
Tobias Roehr, BC Wildlife Federation

Additional Input Provided by:

Suzanne Bailey, University of Alberta
Nora Billy, St'at'imc Government Services
Wade Brunham, ERM
Darwyn John, St'at'imc Government Services
Alicia Krupek, Splitrock Environmental
Karen Kubiski, Carrier Chilcotin Tribal Council
Natasha Lukey, Okanagan Nation Alliance
Kim North, Splitrock Environmental
Daryn Scotchman, Splitrock Environmental
Crystal Wallace, Lower Nicola Indian Band

Special thanks to members of the Steering Technical Committee of the Skeena Environmental Sustainability Initiative for offering additional input into an earlier version of this protocol.

For more information on BC Forest and Range Evaluation Program (FREP) publications, visit our web site at:<http://www2.gov.bc.ca/gov/content/industry/forestry/managing-ourforest-resources/integrated-resource-monitoring/forest-range-evaluation-program/frepreports-extension-notes>.

© 2021 Province of British Columbia

When using information from this or any other FREP publication, please cite fully and correctly.

Table of Contents

Introduction	6
Recommended Steps for a Wetland Health Evaluation	11
Recording Data.....	22
Health Scoring Questions.....	23
Vegetation.....	23
1. Vegetative cover	23
2. Invasive and/or noxious spp	23
2.a) Invasive and/or Noxious Plant Canopy Cover	23
2.b) Invasive and/or Noxious Plants distribution.....	24
3. Coverage of disturbance-caused undesirable species	24
4. Vegetation compared to a healthy unmanaged wetland and riparian plant communities	25
4.a) Wetland plant community structure	25
4.b) Wetland plant recruitment, form and vigor	26
4.c) Upland plant community structure	27
4.d) Uplands plant recruitment, form and vigor	27
4.e) Long-term trajectory of the vegetation community.....	28
5. Sufficient vegetation to minimize windthrow, maintain adequate screening, visual cover and LWD supply	29
5.a) Retention of non-merchantable conifer, understory deciduous trees, shrubs and herbaceous vegetation within 20m of the wetland edge.....	27
5.b) Retention in the CDF, PP, BG, CWHxm, dm, ds and IDFxh, xw, and xm BEC units	29
5.c) Retention in the ESSF, MS, ICH, MH, CWHvm, mm, ms, ws and IDFdm, dk1, and dk2 units	30
5.d) Retention in the SWB, SBS, SBPS, BWBS, CWHvh and IDFww, mw, dk3, and dk4 units	30
5.e) Percent live woody vegetation removed from wetland, other than browsing.	29
6. Heavy browsing and grazing	28
6.a) Heavy browsing.....	28
6.b) Heavy grazing.....	29
6.c) Seedlings or saplings of palatable tree and shrub species.....	29
Soils	32
7. Bare and compacted ground.....	32
7.a) Bare and compacted ground.....	33

7 b) Bare soil within and/or hydrologically connected	33
Morphology.....	34
8. Physically altered site.....	34
8.a) Proportion of altered topography.....	34
8.b) Severity of the physical alteration	35
9. Intact wetland woody debris processes	35
9.a) Standing dead trees (snags) and decadent trees (dying trees) density.....	36
9.b) Standing dead tree diversity	37
9.c) Stable coarse woody debris.....	38
9.d) Diversity of coarse woody debris.....	38
10. Windfirm buffer	39
10.a) Riparian windthrow in RMA without an RRZ or designated WTP	37
10.b) Riparian windthrow in RRZ or WTP.....	40
10.c) Wildlife Trees.....	38
Hydrology.....	38
11. Changes in the hydrologic regime.....	38
11.a) Severity of hydrologic changes	38
11.b) Dead trees or shrubs at the wetland edge?.....	39
11.c) Younger age class plants or trees encroachment	39
11.d) Stream channel incisement.....	39
11.e) Natural surface or subsurface wetted areas.....	43
12. Threats to water levels in the wetland	43
12.a) Outlet structure	43
12.b) Active headcuts below or within the wetland	43
12.c) Disturbed shoreline of the wetland	43
12.d) Deep binding rootmass.....	44
Water Quality.....	44
13. Water quality of the wetland	45
13.a) Excessive nutrient loading.....	45
13.b) Basic water quality parameters	45
Landscape	46
14. Surrounding habitat	46
14.a) Total riparian area disturbance.....	46

14.b) Visible shoreline	47
14.c) Right-of-ways within 100 m	47
14.d) Percent mature and old forest within two kilometers	45
15. Surface and subsurface flows to the wetland intact.....	49
15.a) Mapped and unmapped streams diversion	49
15.b) Contributing basin intercepted by roads or ROWs	47
Appendix 1 – Recommended Field Sampling Sequence	51
Appendix 2 - Common invasive and noxious plants within BC	53
Appendix 3 – Select list of Culturally Significant Plants to Indigenous Communities in BC	56
Appendix 4 – Wetland Indicator Status Categories	60
Appendix 5 - Minimal Targets for Mature and Old Forest Coverage.....	61

List of Figures

Figure 1. A key to the different freshwater wetland ecological classes	9
Figure 2. Wetland boundaries for health assessment	10
Figure 3. Key to wetland riparian classification	13
Figure 4. Placement of transects and plots, and associated calculations.....	16
Figure 5. Distribution Codes for invasive species	24
Figure 6. Visual appearance codes for wildlife trees	37
Figure 7. Edatopic grid for abiotic factors and their relationship with different aquatic classes.....	46
Figure 8. Map of Right of Ways within 500m buffer to identify potential vantage points.....	47
Figure 9. Example of Road impinging within 100m of wetland area	48
Figure 10. Example Spatial query within a 2 km buffer of the wetland	49
Figure 11. Contributing Basin Assessments	47

List of Tables

Table 1. Riparian management distances	13
Table 2. Description of Dominant Upland vegetation Strata Around Perimeter of Wetland.....	16
Table 3. Transect Information.....	19
Table 4. Organic Soils Classification Table based on the Von Post Scale.....	22
Table 5. Example of a completed vegetation plot table	23
Table 6. List of Common disturbance-increaser species in BC	26
Table 7 Structures and Vegetation Form, Vigor, and Recruitment within wetland	28
Table 8. Structures and Vegetation Form, Vigor, and Recruitment upland	28
Table 9. Snags and Coarse Woody Debris Indicators.....	37
Table 10. Tree Diameter Classes	38
Table 11. Decay Classes for Coarse Woody Debris.....	39

Introduction

This guidance document provides supplementary details to the BC Ministry of Forests, Lands, Natural Resource Operations and Rural Development (FLNRORD) field form for evaluating the health of wetlands. Field technicians are expected to read this protocol prior to using the wetland health evaluation form in the field. This document provides information on site selection, sampling protocol, and interpretation of the questions in the form. Technicians are encouraged to refer to this document when evaluating their first few wetlands of the year. Reviewing the document on a regular basis will help calibrate interpretation skills.

The Resource Planning and Assessment Branch within the Ministry of Forests, Lands, Natural Resource Operations and Rural Development (FLNRORD) and the Ecosystems Branch in the Ministry of Environment and Climate Change Strategy (ENV) developed this Wetland Health Assessment Form and Protocol to complement their current suite of monitoring tools within the Forest and Range Evaluation Program (FREP). One example of a well-established FREP tool is the [Riparian Management Routine Effectiveness Evaluation Field Form](#) with the associated [Protocol](#), which assesses stream/riparian condition. The objective of this FREP Wetland Health Assessment Form and Protocol is to allow for persons with basic working knowledge of wetlands to evaluate the health of wetlands within or in proximity to industrial development activities (e.g., forestry). Field evaluators may be forest and range practitioners, First Nation's stewardship members, consultants, land managers, or other land users.

This Wetland Protocol is a coarse-level filter for assessing the health of wetlands. The assessment form is intended to be completed mainly in the field, promote consistency among users, gather pertinent data to inform the health of the wetland, and be cost effective as a tier 2 approach for monitoring (i.e., relatively quick to use in the field (1-4 hours)). One of the advantages of the protocol is that it is a relatively low-cost method to assess wetlands in the field and can enable comparative analysis among numerous sites. A wetland flagged with poor health may require further inspection, or a more detailed analysis may be required to investigate a wetland more thoroughly or to meet other objectives of land managers.

A healthy wetland is a measure of its capacity to perform a number of functions in the environment. Society assigns value to many wetland functions as they can support both healthy landscapes and healthy communities. Although not all wetlands perform the same functions, some of the functions include: sediment trapping, shoreline maintenance, wave energy dissipation, water storage, aquifer or stream recharge, maintenance of biotic diversity, carbon cycling and storage, nutrient cycling and absorption, and primary productivity. The values we derive from these functions may include: flood control, clean water, and provision of food and medicinal products. In order for a wetland to perform these functions effectively, it requires healthy vegetation, intact soils, specific hydrologic regimes, and an appropriate connectivity to the broader landscape. This protocol helps categorize the health of a wetland as: properly functioning; functioning, but at risk; functioning, but at high risk; or not properly functioning.

There are many wetland evaluation protocols in North America that quantify a wetland's functioning condition or assess its health, but none that have been formally and broadly adopted by the Province of BC. This protocol represents the first to be developed at a provincial scale for measuring wetland health. For clarity, the protocol does not quantify specific measurable processes of a wetland (e.g., the amount of

flood retention of a particular wetland), which requires a separate inventory framework. Instead, it focuses on observed or potential impacts that would affect a wetland's ability to perform functions. The development of this protocol is adapted from several existing protocols. Contributing or supporting literature includes:

- Riparian Area Management: A User Guide to Assessing Proper Functioning Condition and Supporting Science for Lentic Areas. TR 1737-16 1999 Revised 2003. USDA and Bureau of Land Management.
 - This guide provides a solid foundation for assessing wetland health and recommends a variety of tools and approaches. The lead author, Don Pritchard, helped establish and popularize the concept of Proper Functioning Condition. However, the indicators within the document rely on a team of experts reaching consensus in order to determine the health of the site. Many of the indicators recommended in this document made their way into the FREP riparian protocol, but additional thresholds and associated guidance were developed to ensure consistency among users.
- Riparian Management Effectiveness Evaluation Field Form and Protocol. 2017. B.C. Ministry of Forests, Lands, Natural Resource Operations and Rural Development.
 - This protocol, developed by the Forest and Range Evaluation Program (FREP), was closely reviewed for elements to transfer. It is expected that field crews completing the riparian protocol will also complete the wetlands protocol. The format for the FREP wetlands form is similar to the riparian version for ease of use and to promote consistency. Several of the indicators that relate to riparian areas are the same or slightly modified for the wetlands protocol.
- Alberta Lentic Wetland Health Assessment Survey; and Alberta Lotic Wetland Health Assessment Survey for Streams and Small Rivers. 2014. Alberta Riparian Habitat Management Society.
 - These two protocols, managed in Alberta by the Alberta Riparian Habitat Management Society, were developed by Paul Hansen and other experts working with the Ecological Solutions Group LLC. Modified versions of the protocol have been applied in Saskatchewan and numerous US states by land managers, as well as by Canadian and US federal agencies. The indicators and protocols have been refined over 20 years, where thresholds and weighting of indicators were developed by a group of experts using the Delphi method. In lieu of a BC wetland health assessment standard, these protocols have been used ad-hoc at a regional scale in BC by FLNRORD, BC Parks, the Nature Conservancy of Canada, and several consultants. Most of the applications have been on range land in the interior; however, according to the lead author, Paul Hansen, the forms were designed to be used for broad application and have been tested on the BC coast with satisfactory transferability. In several cases, where the thresholds for indicators in the FREP form did not exist, a threshold for a Yes or No was derived from the Alberta protocols. At the time of the development of this current FREP protocol, three separate initiatives in BC were attempting to integrate the Alberta Lentic and Lotic protocols into a BC framework. A key difference is that questions in the FREP version also consider natural disturbances, as opposed to only human-caused impacts, so data on the natural variability of wetland health may be determined. With permission from Paul Hansen, several of the guidance question indicators were replicated, and modified as needed, from these two Alberta protocols.

Wetland Delineation

The delineation of the wetland polygon is a critical step in the FREP Wetland Health Assessment as many of the indicator questions are heavily influenced by the defined polygon unit. For example, the percent cover of invasive species will change dramatically with substantial changes in the assessment area. The boundaries can be determined pre-field and adjusted in the field. A copy of recent high-resolution aerial imagery with an overlay of the boundary will improve the assessment in the field. The evaluator may choose the measuring tool on Google Earth imagery, ArcMap, FREPmap (for government staff) or QGIS to draw a polygon boundary and calculate the area. To get started, provincial wetland layers are available from the BC Data Catalogue, such as wetlands in the [Freshwater Atlas](#) and in the [Vegetation Resource Inventory](#). As a word of caution, these wetland boundaries may poorly resemble what is found on the ground, as they are generally created from remote imagery. If available, wetlands may be better delineated on imagery or the forest site plan for a cutblock, and the boundaries refined after field verification. FLNRORD stewardship officers may also be able to calculate wetland areas using the online FREPmap tool that includes recent SPOT imagery. On-the-fly calculations are possible with mobile map apps, such as Avenza. In the near future, drones may serve as a very helpful tool to gather real time data (i.e., both wetland delineation and determining values for some of the indicators). Ensure to save screenshots of any variances in wetland boundaries, that you can then include in your data submission.

In some cases, the evaluator may find that the boundaries need to shift once they get out to the field. The natural boundaries between an upland area and a wetland can generally be determined by a distinct change in vegetation of water-dependent (obligate) or water-tolerant (facultative) species to more terrestrial plant species. The presence of hydric soils (i.e., gleying or mottling of mineral soils within the first 30 cm, the presence of poorly decomposed organic soils (e.g., peat), or a near surface water table can also help delineate the terrestrial/wetland boundary. At the interface of the wetland and deeper water, such as a lake, the wetland is delineated as the vegetated zone with greater than 10% rooted vegetation and less than 2m depth in mid-summer. Any modifications of the boundaries in the field should be included with the assessment.

In some cases, where heavy disturbance has occurred, the boundaries of the wetland may not match the natural observable boundaries. The evaluation relies on understanding the potential area of wetland prior to any disturbances. If there has been soil deposited into the wetland or a decline in water level (and thus a retraction of the wetland area), the evaluator will need to consider the historical context of where the wetland would have been prior to the disturbance to evaluate the impacts. This is one of the more challenging parts of the protocol to consider, especially in heavily modified areas. Old aerial imagery, traditional or community knowledge, or onsite evidence of disturbance may help to determine the historic extent of the wetland. The evaluator will need to take detailed notes and provide a rationale for any changes in the boundary of the assessment area.

The evaluator may choose to only complete an assessment of a portion of a wetland. One reason to do so may be to subsample a wetland that is too large to feasibly assess. Again, the evaluator will need to provide details and document their rationale for modifying the boundary.

The interpretation of the various ecological classes and plant associations within the wetland will help the evaluator answer some of the questions (e.g., Question 4. “Is the vegetation of the entire polygon generally characteristic of what the healthy unmanaged wetland and riparian plant communities are

normally?”). Field crews should become familiar with using the book, *Wetlands of British Columbia: a guide to identification* (2004), which provides substantial detail of many of BC’s most common wetland plant associations and often indicates typical progressions between plant associations of wetter and drier sites. However, the book does not contain all wetlands and the assessor will encounter sites that do not conform to the descriptions. Some wetlands may be in transition between two plant associations, while others are not documented. More recent field guides to ecosystem classification may provide additional detail to plant associations in specific regions.

At a minimum, the field evaluator should become familiar with the various freshwater ecosystem classes: bogs, fens, marshes, swamps, and shallow water. Figure 1 provides a key to the different wetland ecological classes. Patterns of vegetation can often be observed from remote imagery, and differences in patterns often indicate a change between plant associations or wetland ecological classes. For the purposes of this assessment, the evaluator may not need to break the wetland into the various classes, but knowledge of the various types within it can aid in interpretation. See Figure 2 for examples of established wetland boundaries.




Figure 1. A key to the different freshwater wetland ecological classes (Source: Mackenzie 2012. Biogeoclimatic ecosystem classification of non-forested ecosystems in British Columbia. B.C. Min. For., Res. Br., Victoria, B.C. Tech. Rep. 068.). The “surface tier” mentioned in the chart is 40 cm thick excluding any loose litter, crowns of sedges and reeds, or living mosses. Shallow organic soils over mineral soil or

bedrock may have only a surface tier. Of note: in rare cases, some swamps may also contain mesic soils (e.g., Ws05).




Figure 2. Examples of Wetland boundaries established for the health assessment with wetland ecological classes identified. For the purposes of this assessment, in most cases, the boundaries should be established to the extent they would be pre-disturbance. In the example shown at the top of this figure, a road was constructed over a portion of the wetland, and the original extent of the wetland was estimated on the northeast section of the wetland.

Recommended Steps for a Wetland Health Evaluation

The following guidance is provided to help complete the first few sections of the field form. Review the recommended field sampling sequence in Appendix 1 for an efficient order of progressing through tables and checklists once in the field.

STEP 1 – OFFICE REVIEW AND PREPARATION:

Conduct an office review of the wetland polygon area before going out into the field. Complete the information in Tables 1 to 3 (**T.1-T.3**) of the field form to the best of your ability. Many of the questions in this section can be completed in the office but may need to be revised once you arrive in the field.

Useful GIS layers include topographic information (to delineate the contributing basin of the wetland), stream layers, tenure boundaries (range, aggregate, recreation, utilities, etc.), biogeoclimatic information, forest age classes (i.e., from the Vegetation Resource Inventory), cutblocks, and right-of-ways (e.g., roads, pipelines, etc.). Good quality remote imagery will also be useful to complete some sections, such as identifying any mass wasting that may have occurred.

Contact the licensee of the cutblock associated with the wetland to inform them of the intention of entering their operating area, invite them to attend, and request their site plan and any other related information to help with the assessment. The licensee may be able to also inform you of any gates or decommissioned roads that would be important for planning your site visit. It is important to read over the forestry site plan for the wetland of interest, as well as the land use plan if possible. This will ensure a general overview of what management strategies were prioritized and how the wildlife/land was considered.

Wetland Information (T.1.1)

Wetland ID – A unique identifier for the wetland (found by querying the spatial information).

Source of wetland polygon – Select or describe the source that best reflects the most accurate boundaries of the wetland (you may need to confirm this once in the field). Sources may include the Freshwater Atlas (FWA), site plans, the Vegetation Resource Inventory, digitized layers defined by the user, etc.

Biogeoclimatic Ecosystem Classification (BEC) – record the BEC for the area.

Natural Disturbance Type (NDT) – record the natural disturbance type (e.g., NDT1). This will be needed to help answer Q14, and can be found by cross-referencing the BEC sub-zone and variant in the *Biodiversity Guidebook*. (<https://www.for.gov.bc.ca/hfd/library/documents/bib19715.pdf>)

If unknown, both BEC and NDT can be found by following these 5 steps:

1. Launch iMap BC: <https://maps.gov.bc.ca/ess/hm/imap4m/>
2. Zoom to wetland location of interest.

Note: If you are unsure where your wetland is, open up the wetlands layer within the Freshwater Atlas which “may” include the wetland of interest or at least help better orient yourself. Do this by clicking under tab “Data Sources”, select “Add Provincial Layers”. In the pop-up “Add/Remove Map Information”, click to expand options under “Base Maps”, then click to expand options within “Freshwater Atlas”, then select “FWA – Wetlands – Colour Theme” and press OK

3. Select tab “Data Sources” and select “Add Provincial Layers”
4. In the pop-up “Add/Remove Map Information”, click to expand options under “Forest Grasslands and Wetlands”, then click to expand options within “BEC Analysis – Zones/Subzones/Variants – All”, then select “BEC Analysis – Zones/Subzones/Variants – Coloured Themes” and press OK
5. Under the “Find Tab”, select the icon labeled “Point” (it has an icon of a cross-hairs with a blue circle with the letter “i”), then hover your cursor over the wetland of interest, and then left click. A side bar will appear labeled Identify Results. Select the “BEC Analysis – Zones/Subzones/Variants – Colour Theme”. The BGC for the area you selected will appear (e.g., Interior Douglas Fir), click on the symbol “>”, next to the listed BGC, and details for the BGC will appear. Record both the BGC Label including the variant (e.g., IDFxh1), and the Natural Disturbance Type (e.g., NDT4). Put these in the appropriate fields of the form.

Total Wetland Size – Record in square meters (note: 1 ha = 10,000 m²)

Hydrogeomorphic System – Select from estuarine, fluvial, basins & hollows, seepage slopes, lacustrine, ponds & potholes, and marine. Refer to [Mackenzie and Moran 2004](#), (p41) for a descriptor of each type.

Wetland is hydrologically connected to a stream? – Record yes or no.

Helpful tip: Calculate wetland size information on page 3 of the field form. The field evaluator must document Wetland Size, Total Polygon Assessed, and Total Upland Area (10 m buffer) (See pg 16 of the protocol for further information). These values represent the actual wetland size and assessed area as the field evaluator may be required to make adjustments once they visit the site and determine the true boundaries and any access issue. It is helpful, however, to measure these values using desktop tools prior to going into the field, and then adjust if modifications are necessary. This will allow the field evaluator to make many of the subsequent calculations in the field that require these values.

First Nations Information (T.1.2)

List all First Nations territory in which the wetland is located and contact the chief(s) or band offices (as appropriate). The Province of BC’s [Consultative Area Database](#) can help you determine which First Nations may have an interest in a particular wetland. Explain the objective of conducting a wetland health assessment and invite interested individuals to participate in the field-based component.

Opening Information (T.1.3)

Complete this section if the wetland is within 2 riparian management area (RMA) widths of a cutblock.

District – The official three-letter code for the natural resource district (e.g., DCR for Campbell River Natural Resource District).

Opening ID – The unique 5-7 digit code from RESULTS (used each year in the random selection of cutblocks for sampling).

Licensee – The company that holds the forest license for the block.

Forest License – The forest license recorded for the cutblock in RESULTS.

Block – The designation used by the licensee on their logging plan or site plan map.

Harvest Year – The year timber in the selected block was harvested along the sample reach.

Harvest Location – The location of harvesting in the block relative to the sample wetland; record north, east, south, and/or west, as applicable.

Wetland Riparian Class on Plans – Record the wetland riparian class (W1-W5 or unclassified wetland) indicated on the logging plan and/or site plan map.

Wetland Riparian Class in Field – In most cases, this is the same as the wetland riparian class on the logging plan. Record a different wetland riparian class only if it was misclassified. Wetland riparian classes are defined in Figure 3. Note: A wetland complex (W5) must consist of two or more individual wetlands with overlapping riparian management areas and a combined wetland area of 5 ha or more.

Record the riparian management area (RMA), riparian reserve zone (RRZ), and riparian management zone (RMZ) in metres (see Table 1).




Figure 3. Key to wetland riparian classification (Source: Forest Practices Code Guidebook: Riparian Management Area Guidebook 1995).

Wetland Riparian Class	RMA (m)	RRZ (m)	RMZ (m)
NCW	0	0	0
W1	50	10	40
W2	30	10	20
W3	30	0	30
W4	30	0	30
W5	50	10	40

Table 1. Associated riparian management distances from edge of wetland based on wetland riparian class from the B.C. Riparian Management Area Guidebook. Note: NCW = non-classified wetland.

Riparian/Buffer Retention Information (T.1.4)

Riparian/Buffer Retention – Record the following attributes related to the wetland buffer. The buffer can have 3 potential treatments: full retention, partial retention, or no retention. If you feel the information on riparian retention in **T.1.4** does not adequately describe the condition of the riparian area, please add a sketch of the riparian area on page 16 of the field form. A photograph or two of the riparian area can also be invaluable in describing the riparian conditions.

Length of wetland perimeter within assessment polygon (m): Record in meters the length around the wetland edge.

Length along wetland perimeter with full treed retention (m): Record in meters the length of the perimeter where the riparian area has had no harvesting.

Average width of full treed retention present (from wetland edge) (max 100m): Record the width of full retention or record 100 m if the width is greater than 100 m.

Length along wetland perimeter with partial retention (m): Record in meters the length of the perimeter where the riparian area has had recent partial/selective harvesting.

Average width of treed retention present with partial retention (from wetland edge - max 100 m): Record the width of partial retention or record 100 m if the average width is greater than 100 m.

Average treed retention present with partial retention (% of basal area): Record % of basal area remaining.

Length along wetland perimeter with no retention (m): Record in meters the length of the perimeter where clearcut harvesting is adjacent to the wetland boundary.

Description of dominant upland vegetation strata around perimeter of wetland (T.1.5)

Table **T.1.5** of the field form is used to stratify abutting upland areas in order to place sampling transects and to describe the associated upland characteristics. The evaluator will want to stratify the perimeter to represent major differences in the land-base abutting the wetland that may positively or negatively influence the functioning condition of the wetland.

Upland Descriptor for Vegetation Strata and Age – Provide a descriptor for dominant vegetation types within 100 m perpendicular to the wetland perimeter. For example, if the first 30 m of the upland area from the wetland edge is treed retention and the next 70 m is recent cutblock, record recent cutblock. Descriptors may include: recent cutblock; young/mature/old coniferous; young/mature/old deciduous; young/mature/old mixed forest; lake/pond abutting; stream/river abutting; grassland, other.

Disturbance: Record disturbance type. For example, if the upland descriptor was “young forest” , it may be because it was burned. If that is the case, record “recent fire” for disturbance type. Descriptors may include: recent cutblock, old cutblock, recent fire, insect infestation, log sort – landing, road within 100 m, other.

Width – Record the approximate distance the dominant vegetation type extends beyond the wetland perimeter. If it extends beyond 100 m, then record 100 m.

Length Along Wetland Perimeter – Record the length of the wetland perimeter occupied by the dominant vegetation type.

Fraction along wetland perimeter- This is the fraction that is composed of your upland type. If your total perimeter is 800m and the cutblock is along 200m, the fraction is 0.25 (or 25%).

Number of Transects – Record the number of transects to be documented within the wetland for each strata. For wetlands <5 ha, place up to three transects. If the dominant vegetation type is homogenous around the wetland edge (e.g., a cutblock surrounds the entire perimeter), then you may choose to select only two transects. For wetlands with two dominant vegetation types both with >45% of the perimeter occupied, place two transects in the wetland – one in each type. For wetlands with two dominant vegetation types, but one type occupies less than 45% of the wetland perimeter, then place two transects in the dominant upland type and one in the less dominant. For wetlands with greater than three dominant vegetation types, place a transect in each vegetation type that occupies >10% of the perimeter.

Location of Transect Along Perimeter Section – To place transects within the stratified vegetation types, use a random number generator that will mark the location of the transect along the wetland perimeter. The FREP Wetland Filemaker App will automatically do this for you. Otherwise, if the cutblock portion occupies 204 metres of the wetland perimeter, then select a random number between 0 and 204. Using this example, a random number can be generated very quickly in Microsoft Excel by typing in a cell “=randbetween(0,204)”, which will randomly generate a number between the two numbers specified. For consistency, set smaller numbers to larger numbers in a clockwise direction around the wetland perimeter.

Upland Descriptor for Vegetation Strata and Age	Disturbances (s) (e.g., insects, road, cutblock)	Width (m)	Length along perimeter (m)	Fraction along wetland perimeter	Number of Transects	Location(s) of transects along perimeter section		
Mature Coniferous Forest	Road	100	370	0.54	1	345	x	x
Cutblock	Recent Cutblock	80	212	0.31	1	198	x	x
Young Mixed Forest	Old Cutblock	100	105	0.15	1	32	x	x

Table 2. Example of T.1.5: Description of Dominant Upland vegetation Strata Around Perimeter of Wetland.

Other Developments (T.1.6)

Describe the contributing basin upstream of the wetland and estimate the percent that has been developed. You can use the tools provided in ArcGIS, QGIS, Google Earth, FREPmap (government only), or

a georeferenced pdf to draw polygons and calculate the areas of both the watershed and the development to help estimate this value. Include road right-of-ways, agriculture pastures, existing cutblocks, transmission lines, and any other man-made features when calculating the area of development.

Number of Road Crossings – Three numbers are asked for: the number of road crossings within the polygon being assessed (including the 10m upland area), the number of road crossings in the wetland, and the number of crossings on any upstream tributaries that flow into the wetland.

Percent of Watershed Developed Upstream – Delineate the watershed area above the wetland polygon being assessed using topographic map layers and/or imagery and estimate the total percent area developed to date.

Main Development – Record the main human activity present in the watershed area above the sample polygon (e.g., roads, forestry (except roads), agriculture, recreation, mining, oil and gas, transportation, utilities, other, none).

Landscape Indicators (T.2)

Prior to heading out to the field, several of the health indicator questions can be estimated in the office. Refer to the landscape section starting on page 44 in this document for completing section **T.2.** on the form.

Supplementary Management Observations (T.3)

Section **T.3** in the form contains supplemental management observations that can be observed either in the office and/or while on site.

Do the boundaries on the site plan for the wetland coincide with observations in the field?– Record Y or N.

Was there retention around other wetlands observed on the block? Record Y or N – this may be best observed from a combination of map and field observation.

Is four-wheel drive access blocked on roads within 100 m from the wetland edge? Record Y or N.

If rangeland is present, were measures taken to reduce/block livestock access to the wetland edge? Record Y or N. For example, are there any recent fences or logs that appear to purposefully block or control access to the wetland?

STEP 2 – FIELD BASED OBSERVATIONS

Field surveyors are strongly encouraged print, or take an electronic copy, of the 2-page instructions in Appendix 1 to take into the field for an efficient step-by-step process to gather field-based observations.

Once in the field, before starting any measurements, first confirm the wetland is a wetland, the wetland polygon boundary is accurate/appropriate, and the wetland is within two riparian management areas.

At the top of the FIELD SAMPLING INFORMATION page (pg 3):

Check the box for whether sampling is representative of the entire wetland or a portion of the wetland. If only a partial area of a wetland was selected, provide a description on page 18 as to where the wetland polygon was segmented, and provide a rationale for why the partial wetland was selected (e.g., large size, change in management regimes, access blocked, etc.). The entire wetland may not be feasible to assess due to access barriers (e.g., extremely dense brush, deep channels, other safety concerns). After reviewing the site from the office or the field, the evaluator may determine that the wetland poses access barriers and adjust the total polygon assessed.


Total Polygon Assessed – Record in m². For the purposes of this evaluation, the Wetland Polygon Assessed includes the 10 m spatial buffer for some questions. This measurement is the wetland assessed size plus the 10 m upland size. This can be calculated using QGIS or FREPmap (for government employees) by drawing a wetland polygon, then creating a 10m buffer and drawing a polygon around the outer boundary and calculating total area using the GIS tools.


10 m Upland Size – Record in m². This is the area of the 10 m buffer around the wetland to be assessed. Use the previous value for total assessed area and subtract the area that only represents the wetland to derive 10m upland size.

Wetland Assessed Size – Record in m² (will be different if only a portion of the wetland is assessed as part of this evaluation).

Transect Placement

As per instructions on page 14 of this guide, place up to three transects in the wetland. At least one of the transects should enter the wetland upslope of the cutblock to be evaluated. If there is variability in upland seral or ecosystem communities (e.g., young, old, mixed forest, grassland, other) that are greater than 10% along the perimeter, distribute the other transects to best represent the next two most dominant communities (See Figure 4a). Less dominant ecosystems can be “clumped” if needed. Selection of where to place the transects can often be determined from reviewing aerial imagery but may need to be adjusted if there are barriers in the field.





Total Polygon Assessed = Wetland Area + 10 m Adjacent Upland Area – Open Water

Figure 4. Placement of transects and plots, and associated calculations. A) Simple wetland where the entire wetland is part of the total polygon assessed. B) Only a portion of the wetland is assessed due to a barrier to access. C) Placement of plots into the wetland are located to represent major changes in vegetation or seral stage.

For each transect, record the percent of dominant ecosystem type or seral stage that is represented in the upland section in table T.4.1 in the field form. This should match what was recorded in T.1.5. on page 1 of the form. Use this percentage and the percent length of the transect (i.e., in comparison to all transects at your site) to estimate the weighted fraction that will be needed for calculating weighted averages of transect data in T.7.1

UTM Coordinates at Wetland Edge				Transect Bearing into Wetland (0-360°)	Transect is representing wetland influenced from upland strata area composed of:	End Type ¹	Trans. Length	Fraction (Fa) = TL/TTL	Upland Fraction (Fb) From T.1.5.	Weighted Fraction WF = (Fa+Fb)/2	
	UTM Zone	Easting	Northing								
T1	10U	665396	5601259	12°	<i>Mature Coniferous Forest</i>	Ww	53	0.35	0.54	0.44	
T2	" "	...401	...264	211°	<i>Cutblock</i>	Ow	30	0.20	0.31	0.25	
T3	" "	...311	...387	345°	<i>Young Mixed Forest</i>	E	70	0.45	0.15	0.3	
Total Transect Lengths (TTL) (m)							153	1	1		

Table 3. Example of completed T.4.1 Transect Information. Ww = Shallow open Water, Ow = Other, E = End of 50 m transect.

Beginning at the edge of the wetland, record the UTM coordinates for the transect in table T.4.2 on page 3 of the field form. **NOTE:** after writing the full coordinates for the first transect, additional GPS coordinates can be truncated so that there is only a need to record the last 3 or 4 digits of the Northing and Easting when the other numbers are not changing.

The first 10 m of the upland area surrounding the wetland is included in many of the evaluation questions. Walk at a right angle 10 m upslope from wetland edge. In the vegetation plot, T.4.3, record % invasive and % disturbance increasers (record plot as U1 for the first upland transect).

Importance of buffers around wetlands. Buffers around wetlands can provide visual screening for large ungulates and thus provide security from predators (including hunters), limit livestock accessing the site, improve habitat complexity for species that require multiple habitat needs (including many amphibians, flycatchers, cavity nesters, etc.), and reduce sedimentation from upstream erosion. As wetlands provide numerous resources and life needs, they are often part of wildlife movement corridors. Wildlife and human trails are often found around the perimeter of wetlands within the upland portion. Culturally modified trees to mark trails and other signs of human use are of historic and ongoing value to archaeologists and First Nation communities.

After recording buffer parameters, use a rotary tape and walk into the wetland at a right angle from the wetland edge. While completing the transect, walk either: 50 m (i.e., to the end of the rotary tape) or, until you are approximately midway into the wetland (which may occur in smaller wetlands), or until you are unable to go further because of a barrier. Barriers may include: shallow open water, deep water marshes, wetlands that are suspected to have thin mats of soil floating above open water (and pose a safety risk), or deep channels. If you encounter a barrier, record the transect end type in table **T.4.1** (i.e., C = channel, Ww = shallow open water, Ow = open water (where aquatic or submergent is <10% coverage), M = middle of wetland, Ot = other, ET = end of 50 m transect). Otherwise, record NA = not applicable.

As you walk out into the wetland, attempt to identify major changes in vegetation zonation that might indicate a change in wetland class or wetland plant association. Place a plot within each distinct zone you encounter. Your first plot (P1) will describe the first wetland zone encountered as you walk from the edge of the wetland. Begin to fill out information in the Wetland Plot Information table (**T.4.2**). Record the length of each zone under the row associated with the plot representing that zone. For bands of vegetation that are fully within the transect, walk to the center of the zone and record “C” as your plot location. If you reach the end of your 50 m transect, but you haven’t described the vegetation zone and the zone continues, then record “E”. If you encounter a barrier that is a shallow open water wetland (Ww) or a deep marsh (Wm), you will be able to document it by standing on the water’s edge and observing up to the first 10 m (or less if the band of vegetation is narrower), record “W” for water’s edge and record the width of the vegetation band assessed (up to 10m) as your zone length. In T.4.1, tally the transect length as the first 10 m of upland riparian buffer + the length of the transect assessed (which may include up to 10 m beyond the end of the rotary tape if you are at E or W). The largest possible length of a transect is 70 m. From these plot locations (C, E, or W), finish completing the Wetland Plot Information (**T.4.2**) and Vegetation Plot Information (**T.4.3**) sections with the guidance provided below. For sampling purposes, consider 10 m on either side of you, and up to 10 m in front or behind you (but limited to the extent of distinct vegetation for the zone you are trying to estimate).

Soil Samples – With a soil auger or small shovel dig a hole at least 40 cm deep. Determine if the soils are: (1) primarily mineral (i.e., sand, silts or clays), or humic organic, or (2) mesic or fibric organic. Humic organic soils are highly decomposed and it’s difficult to distinguish the origin of the plants that contributed to the soil (think of rich garden soil) aside from the live and/or woody roots that may occupy this layer. Conversely, fibric and mesic organic soils are poorly decomposed and it’s possible to make out some plant fragments within the soil. In order to test the level of decomposition of organic soils, you can use the von Post (VP) scale. Take a handful

of the organic soil and squeeze it in one hand as hard as you can. While doing so, pay attention to the colour and amount of water and peat that is extruded between your fingers, and then make observations about the remaining fragments using Table 4 on the next page. Select the description that best matches your soil sample and categorize it as per the VP Classification. Mark 'Y' in the appropriate soil column in table **T.4.2** of the field form. The determination of these soil properties aids in the classification of wetlands primarily between swamps/marshes and bogs/fens (see Figure 1 in this guide). Swamps and marshes typically have humic or mineral soils, whereas fens and bogs are often associated with mesic or fibric organic soils. Of note, some swamps can occasionally have mesic soils (e.g., Ws05).

Depth to water – If water is present in the soil pit, record the depth to water in cm. Use positive numbers if the water is below the surface. Alternatively, record a negative number (by putting a minus sign in front of the number) if the water is above the surface.

pH and temperature – Use a recently calibrated pH meter to measure pH and record the temperature in degrees Celsius.

Wetland class and wetland plant association - these determinations can be made by considering the soil type and pH in combination with the vegetation plot information in **T.4.3** of the field form. Use Figure 1 in this guide to determine the wetland class. Refer to *Wetlands of British Columbia: A Guide to Identification* (W.H. Mackenzie and J.R. Moran, 2004) to determine the wetland plant association. For bogs, fens, marshes and swamps, the plant associations can be determined by using the two reference tables that are at the start of their respective chapters in the aforementioned publication. The first table organizes wetland plant associations by biogeoclimatic zone and their relative frequency. The second table organizes wetland plant associations by typical percent cover of plants occupying a particular wetland plant association. If the evaluator cannot identify the appropriate class or association, then label “?” for undetermined.

VP Classification		Decomposition	Plant structures	Water	Peat Escape
Fibric Peat	1	None	Unaltered	Clear or light yellow-brown	No peat escapes
	2	Almost none	Distinct	Light yellow-brown	
	3	Very weak		Turbid brown	
Mesic Peat	4	Weak	Clear but becoming indistinct	Strongly brown turbid	Some peat escapes
	5	Moderate		Very strongly brown turbid. Some peat suspension	
	6	Moderately strong	Somewhat indistinct	Muddy with much suspended peat	Approx. 1/3 of peat escapes
Humic Peat	7	Strong	Indistinct but recognizable	Strongly muddy	Approx. 1/2 of peat escapes
	8	Very Strong. Remnants are resistant to decomposition (e.g., roots and wood).	Very indistinct	Thick mud with little water	Approx. 2/3 of peat escapes
	9	Almost complete	Almost unrecognizable	No water	Almost all peat escapes
	10	Complete	Unrecognizable		All peat escapes

Table 4. Organic soils classification based on the von Post scale.

Completing the Vegetation Plot Information Table

A full plant list is not necessary when filling out table **T.4.3** of the field form. In each plot, focus on a few of the most dominant wetland plants (so that you can determine the wetland plant association in **T.4.2.**), invasive and disturbance increaser species (so that you can answer indicator Questions 2a and 3), and plants of cultural importance. For each plant, record the percent cover. Mark “N” if there is poor form, vigor or recruitment. Poor form may include observations of atypical plant structure or physical damage. For instance, when sedge hummocks are over browsed, the crown may be damaged, and they may look irregular as opposed to well-rounded. Poor vigor would include atypical colour (e.g., browning or yellowing), stunted growth, or disease. Poor recruitment includes lack of younger age classes. Also, check off if the plant is an invasive weed (Appendix 2), a disturbance increaser (Table 6 in this guide), or plant of cultural value (Appendix 3). Your region may have lists of weeds that are more appropriate for your study area. Local First Nations may also wish to provide a revised list of culturally significant plants to monitor. Table 5 below provides an example of the completed table.

Transect (e.g., T1, T2)	Plot ID (e.g., U1 or P1)	Species Name For Upland Plots (U), just record invasive or disturbance increasers. For Wetland Plots (P), record dominant, invasive (pg. 53), disturbance (pg. 24), and plants of cultural significance (pg. 56) (if known)	% Cover	Mark N if there is poor form, vigor or recruitment			Check if applicable		
				Form	Vigor	Recruitment	Invasive	Disturbance/Inreater	Cultural Value
T1	U1	Dandelions	10				<input type="checkbox"/>	x	<input type="checkbox"/>
T1	P1	Cattail	90	Y	Y	Y	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
T1	P1	Yellow Flag Iris	5	Y	Y	Y	x	<input type="checkbox"/>	<input type="checkbox"/>
T1	P1	Marsh Cinquefoil	2	Y	N	Y	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
T1	P1	Bare compacted ground	3	NA	NA	NA	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
T1	P2	Willow Sp. 1. Sample taken.	40	N	Y	Y	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
T1	P2	Sedge Sp. 1. Sample taken.	30	Y	Y	Y	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
T1	P2	Mountain Alder	30	Y	Y	Y	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

Table 5. Example of a completed vegetation plot table.

Reading Table 5: The first 10 m of the riparian upland (U1) had 10% dandelions. Other vegetation parameters for the riparian upland are not necessary to document. In the first wetland plot (P1), the surveyor recorded 3% bare soil. In the second plot (P2), the surveyor was unable to identify two of the three species, but indicated that a sample was taken for later verification. The lack of identification to the species level may mean that the second plot will only be classified to wetland class, as opposed to wetland plant association; however, this will not affect the outcome of the assessment.

Completing the Transect Observations Summary (T.5.1)

For each transect, record the width (m) of retention upland from the wetland (perpendicular to the wetland edge) to the cutblock up to a maximum of 100 m. Record 100% if no selective cutting occurred within the area. If the retention of trees was partial (i.e., selective cutting occurred), estimate the percent of trees retained with respect to basal area.

Note: Remaining data within the field cards directly apply to Question Indicators 1 through to 15. Information related to these indicators is organized sequentially below.

Recording Data

Every space in T 5.1., through to T.9 should have a value if the indicator was measured, or “NA” if not. **There should be no blanks, nor other text or symbols (e.g., “estimate”, “>”, “<”) in these columns.** Confine text or symbols to the space provided for recording individual measurements or notes.

The number in the “Total” column for T.5.1., T.5.2., and T.7.1 should be the sum of the weighted averages (i.e., of the transects observations given in in % or #) plus any values (i.e., in % or #) recorded for large non-homogenous patches. The number recorded in the “Weighted Average” column can be calculated using equations provided in the workspace at the bottom of the tables, whereby transect observations

are weighted based on upland fractions (Fb) for T.5.1. & T.5.2., or upland and transect fractions combined (WF) for T.7.1. If using the FileMaker App, then these calculations are automated for you. The number recorded in the “Total” column should be a value that represents the entire polygon. A column for non-homogenous patches allows for the observer to add significant abnormalities for an indicator that are encountered outside of the transects but within the assessment polygon, and that may influence the total value for a particular indicator.

Health Scoring Questions

Vegetation

1. Is vegetative cover of the entire polygon (i.e., wetland portion of polygon AND upland portion of polygon representing 10 m from wetland edge that is within the assessment area) greater than 85%?

In addition to the presence of both water and soils near the surface, vegetation is a key ingredient for defining a wetland. Plants play an important role in carbon and nutrient cycling. Among other services for wildlife and fish, they provide hiding places, homes, and food sources. Woody species, robust emergent vegetation, and root systems can all help to prevent soil erosion. This question estimates how much of the polygon is covered by standing plant growth (both alive and dead). Estimate the percent cover of the entire polygon that is covered by rooted plant material, whether it be alive or dead. Do not consider the area that is covered by water in your estimation of cover (e.g., water between emergent plants) or areas with unrooted plant material (e.g., fallen wood or other plant litter). Use table **T.7.1** in the field form to calculate the weighted average of vegetation cover within the transects, plus any abnormalities outside the transect but within your polygon (e.g., to factor in a large bare patch that falls within the polygon but doesn't land within the transect).

2. Is the presence of invasive and/or noxious species minimal to non-existent in the entire polygon?

Invasive plants are non-native species that are likely to cause environmental or economic harm. They often lack natural predators to control their numbers and can spread quickly. Invasive plants displace native vegetation and are typically of less value to wildlife. After habitat loss, invasive species are considered the second largest threat to wetland ecosystems. Noxious species are native or non-native species that cause damage to economic resources, public health, or the environment. The presence of these species in the polygon indicates a disturbed or degrading ecosystem. Answer the following two sub-questions to score this question. If either of these are scored 'N', then record 'N' for this question.

2. a) Is invasive and/or noxious plant canopy cover less than 5% of the entire polygon?

A low threshold of only 5% is required to score “N” for this question. Once established, many invasive and/or noxious plants can spread quickly. Within the transects, estimate the percent cover of invasive species as a fraction of your answer to Question 1 (i.e., vegetated cover). In other words, evaluate the total percentage of the polygon area that is covered by the combined canopy of all rooted plants (live or dead) of all species of invasive plants. Do not include unrooted material, such as fallen wood or other plant litter. Do not consider area that is covered by water in your estimation of cover (e.g., water between emergent plants). Calculate the weighted average from the upland plot information in **T.4.3** of the field form using the workspace provided at the bottom of **T.7.1**. Then estimate the percent cover of invasive/noxious plants by adding the weighted average within the transects plus any large patches observed outside of the transects but within the polygon in **T.7.1**.

For a list of common invasive plants within BC, please see Appendix 2.

2. b) Is the distribution of invasive and/or noxious plants less than Code 4 in the entire polygon?

Choose the code from Figure 5 below that best fits what is observed in the polygon as the pattern and extent of invasive plant distribution. Due to its specific habitat limitations, including water needs or other tolerances such as shade, the spread of an invasive and/or noxious plant may occupy one or multiple vegetation zones (e.g., marsh, swamp, 10 m upland zone from wetland edge). Even if the plant only occupies one zone, use Figure 5 to score its distribution within that specific zone.

T.7.2. Select the density distribution code of Invasive Species that approximates their extent. Q2b (pg. 22) or mark NA <input type="checkbox"/>								
Code 1 <input type="checkbox"/>	Code 2 <input type="checkbox"/>	Code 3 <input type="checkbox"/>	Code 4 <input type="checkbox"/>	Code 5 <input type="checkbox"/>	Code 6 <input type="checkbox"/>	Code 7 <input type="checkbox"/>	Code 8 <input type="checkbox"/>	Code 9 <input type="checkbox"/>
Rare individual, a single occurrence	Few sporadically occurring individuals	Single patch or clump of a species	Several sporadically occurring individuals	A few patches or clumps of a species	Several well spaced patches or clumps of a species	Continuous uniform occurrence of well spaced individuals	Continuous occurrence of a species with a few gaps in the distribution	Continuous dense occurrence of a species

Figure 5. Distribution codes for invasive species. Source: Invasive Alien Plant Program – Field Forms. <https://www.for.gov.bc.ca/hra/plants/forms/FS1260.pdf>

3. Is the coverage of disturbance-caused undesirable species (e.g., domestic grasses, dandelions, pineapple weed, buttercups, etc.) less than 25% of total area in the riparian upland area 10 m from wetland edge?

Disturbance-caused undesirable species will often perform poorly, have shallow roots, and are less productive than other plants. They may result from a disturbance that has removed a more desirable species. Use tables T.4.3 and T.5.1. in the field form to answer this question. For T.4.3, you can estimate the percent cover of each species within your plots. For T.5.1, calculate the weighted average of the transects plus add any large patches that are outside the transect but within the upland portion of your polygon. Since you are only considering the upland portion of your transects, the transect lengths considered are equal (i.e., 10 m) and are not included in the weight. Your weighted fraction will only adjust the percent cover to account for the upland fraction (Fb), % of the wetland perimeter, that each transect represents (Refer to Table T.4.1). Thus for this question, the weighted average is calculated as $(T1\%dist.sp * T1Fb) + (T2\%dist.sp * T2Fb) + (T3\%dist.sp * T3Fb)$. Table 4 below lists common disturbance species in BC. Invasive plant species that have been considered in Question 2 are not reconsidered.

Common disturbance-increaser species in BC			
Common name	Latin name	Common name	Latin name
Strawberry	<i>Fragaria spp.</i>	Pineapple weed	<i>Matricaria matricariodes</i>
Cinquefoil	<i>Potentilla spp. (excl. Marsh Cinquefoil)</i>	Dock	<i>Rumex spp.</i>
Yarrow	<i>Achillea millefolium</i>	Pasture sage	<i>Artemisia frigida</i>
Baltic rush	<i>Juncus balticus</i>	Gumweed	<i>Grindelia squarrosa</i>
Dandelions	<i>Taraxacum spp.</i>	Pussytoes	<i>Antennaria spp.</i>
Sow thistles	<i>Sonchus spp.</i>	Buttercups	<i>Ranunculus spp.</i>
Foxtail barley	<i>Hordeum jubatum</i>	Bluegrasses	<i>Poa spp.</i>
Yellow salsify/ Western goat's beard	<i>Tragopogon dubius</i>	Plantains	<i>Plantago spp.</i>
Clovers	<i>Trifolium spp.</i>	Willowherbs	<i>Epilobium spp.</i>

Table 6. List of Common disturbance-increaser species in BC.

4. Is the vegetation of the entire polygon generally characteristic of what the healthy unmanaged wetland and riparian plant communities are normally?

This question aims to determine if there are any vegetation or structural components within the wetland or 10 m upland from the wetland edge that are missing or in poor condition. The evaluator should consider the following questions while completing transects and walking between sites. Questions 4a - 4d have associated fillable tables in the field form to help make calculations. These tables need only be used if there are observable impacts from recent treatments (e.g., logging) to various wetland or riparian plant communities and structural elements, where the evaluator needs to determine if the plant communities and structural elements drop below a percent threshold. The threshold for wetlands (4a and 4b) is set at 85% compared to 75% for upland areas. The rationale for setting a more sensitive threshold for wetlands is that wetlands can take much longer to recover and thus there should be less tolerance for impacts directly within wetlands.

4.a) Is greater than 85% of the layers and features in the wetland portion of the polygon intact?

Familiarity with the various wetland ecological classes and plant associations will help the evaluator better understand what to expect in terms of structure at these sites. This question should consider information gathered while walking the transects as well as observing the various wetland classes present at the site. There is often a high degree of variability within wetland sites – even within the same wetland class or plant association. *Wetlands of British Columbia: A Guide to Identification* (Mackenzie and Moran, 2014) describes a range of variability in terms of plants and structure for each plant association. Look for elements that are obviously missing due to recent disturbances. Only consider features that you can see or that you know are missing from observable impacts. If an element is missing, but there is uncertainty to whether it was present prior to a disturbance, do not record it as missing. Important components may include: hummocks, wetted depressions, wildlife trees, downed wood, and vegetation. Hummocks provide elevated microsites on slightly drier ground. Elevated microsites are valuable components of a wetland because they contain unique characteristics, features and conditions which can act as seed beds

for ectomycorrhizal forest trees, shrubs, plants and fungi. However, they are vulnerable to erosion or mechanical damage. Wetted depressions are important for a variety of wildlife to access open water but are vulnerable to sedimentation or overburden from the deposition of too much downed wood. Consider similar sites that are in close proximity but relatively less disturbed for comparison.

EXAMPLE: A wetland is evaluated that has both a marsh and a forested bog wetland ecological class. Recent windthrow has impacted approximately 25% of the spruce (both snags and overstory trees) present in the bog. As is often the case, the marsh naturally doesn't contain either of these elements, so only the bog is assessed for impacts to these layers. Indicate that snags and overstory are impacted and record that 75% remain for each category. The excessive downed wood has increased the CWD to where it is estimated to be over and above what is natural by about 20%. This translates to 80% for the CWD layer (proportion relative to what is normal). The excess of downed wood covers approximately 50% of the open water pools in the entire polygon. The remaining pools appear to have aquatic submergent and floating vegetation, and it's suspected that the open water pools that were impacted also had these elements, so Herbs – Aquatic is also scored at 50%. Other layers that are observed but do not have apparent impacts are rated 100%. Do not record anything for layers that are not present – these rows in the table marked NA, unless there is evidence that the entire layer has been impacted. Using this example, the total sum of percentages representing the impacted layers adds to 630%, with an average of 79% (see Table 7 in the next section). In this scenario, the evaluator would record 'N' for this question, as the value is below the 85% threshold.

4. b) Does greater than 85% of all expected layers and components show good recruitment, form and vigor in the wetland?

Use the form, vigor and recruitment columns in **T.4.3** on the field form to help complete the table in **T.8** to answer this question. List each vegetation layer where impacts are identified and give a "Y" or "N" answer in terms of its form, vigor and recruitment. Only record the layers that have been identified as present or 100% lost. Do not include layers that are missing if it is unknown whether they were present previously. If less than 85% of the responses are "Y", indicating poor form, vigor or recruitment, mark "No" for the question (see Table 5 below).

FORM, VIGOR AND RECRUITMENT WITHIN WETLAND							
Using the table below, estimate the percent (%) of all observed layers or those 100% lost due to a disturbance. Additionally, using Yes or No answers, determine if all expected layers and components show good form, recruitment and vigor. This estimation is based only on the wetland portion of the polygon. (Q4a+b)							
Layer/Feature	Typical Associated Wetland Class	Check if present OR evidence of 100% lost	NA	Remaining after a disturbance (%)	Form (Y/N)	Vigor (Y/N)	Recruitment (Y/N)
Snags	Wb, Ws	x	<input type="checkbox"/>	75	N	NA	Y
Over-story Trees	Wb, Ws	x	<input type="checkbox"/>	75	N	Y	Y
Under-story Trees	Wb, Ws	<input type="checkbox"/>	<input checked="" type="checkbox"/>	NA	NA	NA	NA
Tall Shrubs	Wb, Ws	<input type="checkbox"/>	<input checked="" type="checkbox"/>	NA	NA	NA	NA
Low Shrubs	Wb, Ws, Wf	<input type="checkbox"/>	<input checked="" type="checkbox"/>	NA	NA	NA	NA
Herbs – Terrestrial/ Emergent	Wb, Ws, Wf, Wm, Ww	x	<input type="checkbox"/>	100	Y	Y	Y
Herbs – Aquatic	Wm, Wf, Ww	x	<input type="checkbox"/>	50	N	N	Y
Elevated Microsites	Wb, Wf, Ws	x	<input type="checkbox"/>	100	Y	NA	Y
Mosses/Lichens	Wb, Wf, Ws	x	<input type="checkbox"/>	100	Y	Y	Y
CWD	Wb, Ws	x	<input type="checkbox"/>	80	Y	NA	Y
Open Water Pools (>4m ²)	Wb, Wf, Wm, Ws	x	<input type="checkbox"/>	50	N	NA	Y
				Total (Sum of %'s)	Total possible number of Yes answers		
				630	20		
				Average (%) (Q4c)	Actual number of Yes answers		
				79	15		
					% of cells with Yes answers (Q4b)		
					75		

Table 7. Example of structures and vegetation form, vigor and recruitment within the wetland portion of the polygon.

*Depending on the site you are evaluating, certain structural features may not be relevant. Wetland layers/features listed in column 1 are more common in those wetland classes listed in the adjacent cell (column 2). Note: Wb = bog, Wf = fen, Wm = marsh, Ws = swamp, Ww = shallow open water. Enter NA for features not observed at the site or where the evaluator is not confident if the layer is missing due to disturbance.

4.c) Is greater than 75% of the layers and features in the 10 m upland portion of the polygon intact?

Follow the same steps as 4 a) but assess the upland portion of the polygon (see Table 8 in next section).

4. d) Does greater than 75% of all expected layers and components show good recruitment, form and vigor in the upland portion of the polygon?

Evaluate each vegetation layer that is present or 100% removed and give a “Y” or “N” answer in terms of its form, vigor and recruitment.

Record the total number of layers receiving “Y” answers in the appropriate box. If less than 75% of the responses are “Y”, indicating poor form, vigor or recruitment, mark “No” for the question.

Using the table below, estimate the percent (%) of layers and features that have been 100% lost due to a disturbance. Additionally, using Yes or No answers, determine if the layers and features show good form, recruitment and vigor. This estimation is based only on the upland portion of the polygon. (Q4+d, pg. 25)						
Layer	Check if present OR evidence of 100% lost	NA	Remaining after a disturbance (%)	Form (Y/N)	Vigor	Recruitment (Y/N)
Snags	<input type="checkbox"/>	<input type="checkbox"/>			NA	
Over-story Trees	<input type="checkbox"/>	<input type="checkbox"/>				
Under-story Trees	<input type="checkbox"/>	<input type="checkbox"/>				
Tall Shrubs	<input type="checkbox"/>	<input type="checkbox"/>				
Low Shrubs	<input type="checkbox"/>	<input type="checkbox"/>				
Herbs	<input type="checkbox"/>	<input type="checkbox"/>				
Gaps	<input type="checkbox"/>	<input type="checkbox"/>			NA	
Mosses	<input type="checkbox"/>	<input type="checkbox"/>				
Lichens	<input type="checkbox"/>	<input type="checkbox"/>				
CWD	<input type="checkbox"/>	<input type="checkbox"/>			NA	
			Total (Sum of %'s)	Total possible number of Yes answers		
				Actual number of Yes answers		
			Average (%) (Q4c)	% of cells with Yes answers (Q4d)		

Table 8. Structures and vegetation form, vigor and recruitment within the upland portion of the polygon.

4. e) Is the % of the long-term trajectory of the vegetation community altered less than 15% for the entire polygon?

Questions 4a - 4d consider current structure and include both short- and long-term impacts on the vegetation structure. There is some overlap, but this sub-question considers only changes that will alter the trajectory of the vegetation community over the long term. For this sub-question, take into consideration the entire polygon (i.e., both the wetland and 10 m upland from the wetland edge). This question can be answered using aerial imagery combined with observations in the field.

Large tree growth is often confined to raised microsities in wetlands (especially wetland swamps with plant associations Ws07, Ws08, Ws10, Ws11, Ws53, Ws54, Ws55), but when harvesting removes the canopy, the water table can rise and reduce the microsities that are otherwise dry enough to support trees. In other circumstances, competing vegetation within the shrub layer may limit future tree establishment without brush control. When areas are cleared, they may become more frost prone and cause seedling mortality due to frosts during the growing season. Such observations should be recorded as affecting the long-term trajectory of the vegetation community at the site.

Other long-term impacts to vegetation may include the limitation for native vegetation communities to establish due to colonization by invasive/noxious non-native species. If, for example, reed canary grass encroaches on marsh vegetation, this is considered a dis-climax community (Mackenzie and Moran, 2004). In other circumstances, conversion of the site to lawns, agricultural fields, or planting of non-native trees can affect the site's long-term trajectory.

5. Has sufficient vegetation been retained to minimize windthrow and maintain adequate screening, visual cover, and LWD supply?

Although similar to question 4, this question considers the specific impacts of live woody species removal. Live woody plants provide important structure and help recruit CWD and snags for wildlife. Include all causes of live wood removal (e.g., insect outbreaks, fire or logging) with the exception of impacts from browsing. Browsing refers to woody material being harvested by animals and is addressed in Question 6.

5. a) On all wetlands, have most (95%) of non-merchantable conifers, understory deciduous trees, shrubs, and herbaceous vegetation been retained within 20 m of the wetland edge?

This metric is based on the *Riparian Management Area Guidebook* best practices (Tables 14-16), which recommend retention of most non-merchantable timber, understory deciduous trees, shrubs and herbaceous vegetation within the first 20 m of a wetland edge.

For the purposes of this assessment, non-merchantable is defined as follows:

Interior regions:

- Cedar – less than 15 cm diameter inside bark (DIB).
- All other tree species – less than 10 cm DIB.

Coastal regions:

- Harvesting within mature stands – less than 15 cm DIB, where mature stands are >121 years for coniferous and >41 years for deciduous stands.
- Harvesting within immature stands (e.g., for thinning) – less than 10 cm DIB, where immature stands are <121 years for coniferous stands and <41 years for deciduous stands.

DIB is estimated at approximately 1.3 m from the base of the tree or at the height of the stump below 1.3 m (Ministry of Forests 2005. Provincial Logging Residue and Waste Measurement Procedures Manual. Timber Pricing Branch).

5. b) For wetlands in the CDF, PP, BG, CWHxm, dm, ds and IDFxh, xw, xm biogeoclimatic units, have all wildlife trees, 70% of the mature co-dominant windfirm conifers, and all deciduous trees been retained in the Riparian Management Zone?

This metric is based on the *Riparian Management Area Guidebook* best practices for the riparian management zone adjacent to wetlands, Table 14. Codominant trees are defined as: "A tree whose crown helps to form the general level of the main canopy in even aged stands, or in uneven-aged stands, the main canopy of the tree's immediate neighbours, receiving full light from above and comparatively

little from the sides.” (Ministry of Forest and Range 2008. Glossary of Forestry Terms in British Columbia <https://www.for.gov.bc.ca/hfd/library/documents/glossary/Glossary.pdf>)

5. c) For wetlands in the ESSF, MS, ICH, MH, CWHvm, mm, ms, ws and IDFdm, dk1, dk2 biogeoclimatic units, have all wildlife trees, 40% of the mature co-dominant windfirm conifers, and all deciduous trees been retained in the Riparian Management Zone?

This metric is based on the *Riparian Management Area Guidebook* best practices for the riparian management zone adjacent to wetlands, Table 15.

5. d) For wetlands in the SWB, SBS, SBPS, BWBS, CWHvh and IDFww, mw, dk3, dk4 biogeoclimatic units, have all wildlife trees, 10% of the mature co-dominant windfirm conifers, and 30% of the deciduous trees been retained in the Riparian Management Zone?

This metric is based on the *Riparian Management Area Guidebook* best practices for the riparian management zone adjacent to wetlands, Table 16.

5. e) Percent live woody vegetation removed from the wetland, other than browsing.

Live trees and other woody vegetation provide important structure and are recruits for CWD and snags for wildlife. Using field observations, you will record any live woody vegetation removal in the wetland. Include all causes of removal (e.g. Insect outbreaks, fire or logging) except for impacts from browsing which is captured in a separate question. Do not consider the loss of volume from woody species that are purposefully removed because they are non-native and invasive (e.g., Russian olive). Check “Yes” for Question 5e if there is no removal of live woody vegetation.

If there are any “No” answers for the sub-questions, mark the “No” box for Question 5. Note that if a wetland is in a range area, retention should be greater than the minimum required, otherwise range barriers should be constructed to discourage access by livestock.

6. Is heavy browse and grazing absent?

There is a relationship between this question and Question 4, in that if Question 6 is scored “No”, it may also result in a “No” answer in Question 4. However, Question 6 is referring to impacts specifically from browsing and grazing by animals. Browse represents the impact on woody species, whereas grazing represents the impact to non-woody herbaceous species (i.e., mostly grass-like species). Both wildlife and livestock may over-utilize a site. If heavy browse and/or grazing is present and persistent over time, it can lead to unsustainable modifications of available food sources through a gradual transition to less palatable species.

6. a) Is less than 50% of second and older leaders of palatable species browsed in the entire polygon?

For the purposes of this evaluation, heavy browse is when more than half of second and older leaders are browsed. Attempt to make observations on at least three palatable shrubs or trees, such as *Thuja plicata* (Western redcedar), *Salix* species (willows), *Cornus stolonifera* (red-osier dogwood), or *Amelanchier alnifolia* (Saskatoon berry) within the polygon. Check the branches of palatable species that would be within reach of animals (e.g., up to a height of six feet) for evidence of browse. Estimate the amount of recent browse that surpasses this year’s growth (the new shoots from this year) and encroaches on last year’s growth or older (i.e., older branches are often more rigid and dull in colour). A shrub or tree that

appears to have unusually compact branches or is umbrella shaped is an indication of heavy browse. Consistent with the FREP riparian/stream protocol, also record 'N' for heavy browse if a stem is recently chewed by beaver.

In heavily browsed areas, there may be under-representation of palatable species and overrepresentation of less-palatable species, including:

- *Crataegus* spp. (hawthorn);
- *Elaeagnus commutata* (silverberry/wolf willow);
- *Dasiphora fruticosa* or *potentilla fruticosa* (shrubby cinquefoil);
- *Rosa* spp. (rose);
- *Symphoricarpos* spp. (buckbrush/snowberry); and
- Non-native species such as *Caragana* spp. (caragana) and *Elaeagnus angustifolia* (Russian olive).

If these less palatable species dominate the understory, look for adjacent sites at similar elevations to compare and determine if the site's vegetation community may have shifted due to overbrowsing. If not listed above, consider other palatable deciduous plants that may be evaluated for browse pressure.

6. b) Is 90% or more of the available grazing area free of heavy grazing?

Consider the polygon free of heavy grazing if 90% or more of the available forage has a stubble height greater than the recommended minimum (usually 5 cm or 2 inches for most ranges). Consider heavy grazing present if only short stubble remains, almost all plant biomass has been removed, and only the root systems and parts of the stems remain. Heavy grazing can lead to undesirable invaders inhabiting the site as the native plants are prevented from flowering and often extirpated. Do not count dead plants unless it is clear that death resulted from overgrazing.

6. c) Do seedlings or saplings of palatable tree and shrub species make up more than 5% of those species in the entire polygon?

Only consider the forested portion of the polygon. Mark NA if the site lacks potential for trees or shrubs (e.g., the site is an herbaceous wet meadow or marsh). Not all riparian areas can support trees and/or shrubs; however, on sites where such species occur, they play important roles. The root systems of woody species are excellent bank stabilizers, while their spreading canopies provide protection for soil, water, wildlife, and livestock. Young age classes of woody species are important for the continued presence of woody communities not only at a given point in time but into the future. On severely disturbed sites, the evaluator should seek clues to vegetation potential by observing nearby sites with similar landscape position. **NOTE:** Vegetation potential is commonly underestimated on sites with a long history of disturbance.

The following species are excluded from the evaluation (**those not listed are considered preferred**):

- *Crataegus* species (hawthorn);
- *Elaeagnus commutata* (silverberry/wolf willow);
- *Dasiphora fruticosa* or *potentilla fruticosa* (shrubby cinquefoil);
- *Rosa* species (rose);
- *Symphoricarpos* species (buckbrush/snowberry); and
- Non-native species such as *Elaeagnus angustifolia* (Russian olive) and *Caragana* species (caragana).

The above species may reflect long-term disturbance on a site, are generally less palatable to browsers, tend to increase under long-term moderate to intense grazing pressure, and rarely have any problem maintaining a presence on site. *Elaeagnus angustifolia* (Russian olive) and *Caragana* species (caragana) are considered especially aggressive, undesirable exotic plants.

The main reason for excluding the above plants from the evaluation is that they can be more abundant than desired species (e.g., *Thuja plicata* (Western redcedar), *Salix* species (willows), *Cornus stolonifera* (red-osier dogwood), *Amelanchier alnifolia* (Saskatoon serviceberry), and many other taller native riparian species). This means they could mask the ecological significance of these more desirable species, especially when the desired species are present in smaller quantities.

EXAMPLE: A polygon may have *Symphoricarpos occidentalis* (buckbrush/snowberry) with 30% canopy cover represented by young plants that will replace older ones, while also having a trace of *Salix exigua* (sandbar willow) present but represented only by older mature individuals. The failure of the willow to regenerate (even though there is only a small amount) is very important in the health evaluation, but by including the buckbrush/snowberry and willow together in the polygon, the condition of the willow would be hidden (i.e., overwhelmed by the larger amount of buckbrush/snowberry).

For shrubs in general, seedlings and saplings can be distinguished from mature plants as follows. For those species having a mature height generally over 1.8 m (6.0 feet), seedlings and saplings are those individuals less than 1.8 m tall. For species normally not exceeding 1.8 m, seedlings and saplings are those individuals less than 0.45 m (1.5 feet) tall or which lack reproductive structures and the relative stature to suggest maturity. Count plants installed by human planting if these are successfully established, which means they have survived at least one full year after planting. **NOTE:** Evaluators should take care not to confuse short stature resulting from intense browsing with that due to young plants.

Soils

7. Has bare and compacted ground been minimized in the entire polygon?

Wetlands are natural sediment traps that will eventually fill in over long periods of time. During this slow phase of sediment entrapment, water clarity for the most part is very good, while the substrate remains biologically active. Microbial, plant and invertebrate populations are typically unaffected as a result. Sudden, or chronic but larger than normal introductions of sediment can upset this balance, resulting in reduced productivity at almost all trophic levels, either indirectly through reductions in water clarity or directly when biologically active substrates are inundated with new sediments. The lifetime of the wetland is also reduced.

Bare ground includes any soil or fill with particles smaller than 2 mm (small peppercorn-sized) that is not covered by plants, litter, lichens, moss, downed wood, or coarse gravel that water can wash into an adjacent wetland or stream connected to a wetland. Examples include road cut-and-fill slopes, bladed trails, gouges and scalps due to yarding, tipped over root wads, and windthrow scars, slides and slumps. It also includes animal trails or recreation trails if mineral soil is exposed. Compacted ground is any ground that does not absorb water readily. It includes both gravel or paved roads, as well as skid trails, backspare trails, ATV trails, bike trails, or animal trails. Compacted ground also includes the pugging and hummocking found where cattle or other ungulates walk through heavy, saturated soils.

7. a) Is there less than 5% bare and compacted ground in the entire polygon?

Check that sediments do not cover the wetland polygon for any more than 5% of the total polygon. Look near inlets or where water may enter via sheet flow and examine these areas for recent evidence of deposition. Typically, shallow open water, marshes and swamps experience periods of flooding and have greater chances of receiving upstream inputs.

Measure and record the amount of bare and compacted ground present in the wetland, as well as in the first 10 m of the riparian area if there is a sign of sediment transport to the wetland. This includes all the hydrologically connected portions of active and inactive roads within the 10 m area. In estimating the amount of bare and compacted ground, remember to net out the vegetation, gravel, rocks, roots, debris, etc. that are not erodible.

EXAMPLE: The lower surface of a tipped-over root wad measures 3 m X 2 m, with about half of this area (0.5) comprised of erodible soils. The rest of the area is comprised of roots and coarse sediments. The root scar is of a similar area and consistency but it is in a depression and clearly not connected to the wetland. You see a sediment trail and accumulations at the edge of the wetland originating from the side of the root ball. Considering the connectivity scale as 0 (not connected) to 1.0 (fully hydrologically connected) you apply a connectivity factor of 0.5 to the total area of the root wad that is potentially connected to the wetland so the total net value = $-(3 \text{ m} \times 2 \text{ m} \times 0.5) \times 0.5 = 1.5 \text{ m}^2$. Do this type of calculation for each identifiable patch of ground with bare erodible soil in the first 10 m of the riparian area.

7. b) Is the amount of bare soil within and/or hydrologically connected to the entire polygon less than 15%?

In most cases, the landscape will be quite flat and sediment transport won't be an issue. However, if you encounter sediment deposits or evidence of sediment transfer to the wetland polygon and suspect it may be originating from upslope, further assessment is needed. It is possible to have disturbances in the uplands and still not see major changes in the magnitude, timing or duration of overland flows that negatively impact riparian/wetland areas. No upslope measurements are necessary if there is no evidence that erosion deposits from the uplands are affecting the wetland area.

If erosion or fine sediment deposits are observed in the wetland polygon and are suspected to have been a result of upslope activity, then calculate any hydrologically connected compacted and bare ground outside of the wetland polygon. Include all roads, landings, cutslopes, rootwads, and other sources of compacted and bare soil. Hydrologically connected ground is any compacted or bare soil that shows evidence of sediment transfer to the wetland polygon. Slopes >10% immediately adjacent to the wetland polygon are assumed to be hydrologically connected to a maximum distance of 30 m. Bare or compacted ground on slopes that are beyond 30 m from the wetland polygon should show some signs of surface water transport in order to be included in the calculation. After calculating the area (m²) of any bare or compacted ground hydrologically connected to the wetland polygon, multiply by the degree of hydrologic connectivity to get a net area of hydrologically connected ground.

To estimate how hydrologically connected the patch is to the riparian area, use simple values of 0.0, 0.2, 0.5, 0.8 and 1.0 for connectivity that are equal to visual estimates of none, a little, half, a lot and all, respectively.

Add up all the net bare and/or compacted ground area present within the wetland polygon, plus that which is hydrologically connected to the riparian area. Divide this number by total polygon area and then multiply this number by 100 to get the percent of the wetland polygon that is affected by bare and/or compacted ground, within and hydrologically connected to the polygon area.

It will be difficult to survey all potential bare and/or compacted ground in and around large wetlands. Focus instead on the most obvious sources such as roads, trails or concentrations of windthrow. For homogeneous disturbance, subsample three to six sections or plots (e.g., plots 10 m² in area) and extrapolate the results to the total wetland polygon area. Livestock or game trails if present are likely to be ubiquitous around the entire forested portion of the wetland, requiring information only on the number of trails on average on cross-sections across the riparian area, average trail width, and average percent coverage by fines and sands.

Morphology

8. Has less than 15% of the topography of the entire polygon been physically altered with subsequent impacts to vegetative communities and hydrologic function?

The purpose of this question is to assess physical change to the soil, bank/shore integrity, hydrology, etc. as it affects the ability of the natural system to function normally. This question seeks to assess the accumulated effects of all physical alterations (both natural disturbances and human caused). If both of the following questions have “No” answers, mark the “No” box for Question 8.

8. a) Is less than 15% of the topography of the entire polygon physically altered?

Estimate the percent impacted by both natural and human-caused impacts. Changes in shore and bank contour and any change in soil structure will alter infiltration of water, increase soil compaction, and cause increased sediment contribution to the water body.

Include all changes to the physical attributes of the site caused by human actions (e.g., logging, mining, housing development) or by agents of human management (e.g., livestock), wildlife use, as well as natural events. Examples of natural impacts include slides or flood deposition that may or may not be directly caused by human alterations. If uncertain about the cause or if it is indirect, such as climate change or sedimentation after wildfire, then consider these to be natural when filling out the causal factor checklist on page 16 of the field form.

The kinds of physical change that diminish or disrupt natural wetland functions on the site include, but are not limited to:

Soil Compaction – This kind of alteration includes livestock-caused hummocking and pugging, animal (wildlife or livestock) and recreation trails that obviously have compacted the soil, vehicle and machine tracks and ruts in soft soil, etc.

Plowing/Tilling – This is disruption of the soil surface for cultivation purposes. It does not include the alteration of drainage or topographic patterns, which are included in the **Topographic Change** category.

Hydrologic Change – Include in this category any area that is physically affected by removal or addition of water for human purpose. The physical effects to look for are erosion due to reduced or increased water, bared soil surface that had water cover removed, or flooded area that normally supports a drier vegetation type.

Human Impervious Surface – This includes roofs, hardened surfaces like walkways and roads, boat launches, rip-rapping of shores and banks, etc.

Topographic Change – This is the deliberate alteration of terrain and/or drainage patterns for human purposes. It may be for aesthetic (landscaping/beach clearing) or other reasons, including such structures as water diversions ditches and canals.

NOTE: Do not count the same area twice by including it as both a vegetative (Question 4) and a physical alteration, unless there clearly are both kinds of alteration. **EXAMPLE:** A cottage owner may clear vegetation to gain a view of the lake without causing topographic change to one area, whereas if sand is hauled in to enhance the beach, there is also physical alteration of the same site.) Decide into which category each particular effect should go.

8. b) Is the severity of the physical alteration slight or non-existent?

Four categories of alteration severity are described in terms of change to the site vegetation and hydrologic function. **NOTE:** This criterion uses vegetation change to indicate the degree of physical alteration, but the alteration must be physical in nature, not just vegetative change alone (e.g., disruption of soil, hydrology (including infiltration/interception of water), topography, etc.). Document with photos and commentary. Categories of severity of human-caused physical alteration are described below with conceptual guidelines. These guidelines are not comprehensive but are intended as a relative scale by which the observer can judge the site. Every case is different, and there is no absolute measuring stick to apply. Use the following comparative descriptions to choose a category of alteration on the site:

No alteration: No human-caused physical alteration observed on the polygon.

Slight: Physical site integrity is near natural. Alteration (including recovery from any past severe alterations) is apparent but reflects minimal impact to plant communities and hydrological function in the altered areas (e.g., the plant community shows little change from that on nearby sites lacking physical alteration; any changes to microtopography are slight and the site is well vegetated with appropriate species).

Moderate: As compared with nearby unaltered sites, human-caused physical alteration on the polygon (including recovery from any past severe alterations) has noticeably altered the physical site integrity to the point that plant communities and hydrological function on the altered areas show visible impacts. The plant community differs noticeably (by having introduced or missing components) from nearby sites on similar landscape positions that lack physical alterations. Changes to the microtopography of the soil profile are moderate in depth. Such alteration is either in the process of becoming revegetated with appropriate species or is well covered with a mix of less desirable and appropriate species.

Severe: Human-caused physical site alteration on the polygon has compromised the physical integrity of the altered areas (even if only a small area is altered). Old alterations have not recovered and are still affecting the vegetation or hydrological functions (e.g., the plant community differs radically from nearby sites in similar position that lack physical alterations, reflecting altered hydrologic and/or soil conditions). Disruption of the microtopography of the soil profile is severe in depth of disturbance. Alterations remain mostly bare of plant cover, are no longer supporting wetland habitat, or are becoming vegetated with invasive or undesirable species.

9. Are wetland woody debris processes intact 10 m upland of the wetland?

Coarse woody debris (CWD, or more simply “wood”) is a critical component of all forest ecosystems. Standing dead trees are one of the main sources of CWD in riparian areas. The other main sources include

the limbs, stems and root wads of live trees that get blown over or snapped off. Studies on where CWD originates in riparian areas of streams regularly show that 80-90% of the CWD originates from trees and snags located within 10 m of the stream edge. A similar relationship is expected for CWD around wetlands.

Wetland CWD processes to consider include: origin of CWD, wood type (e.g. size, decay class), and function. Changes in wetland wood processes can have a major impact on the habitat characteristics of wetlands and its use by fish and wildlife. For the purposes of this protocol, CWD is defined as any branch or stem with a minimum diameter of 7.5 cm. Length is not a factor.

In most cases, it will not be practical to survey the entire wetland polygon to determine if woody debris processes are within a normal, natural range of variation. Use table **T.7.1** in the field cards to record observations of dead and dying wood, and CWD during the site survey. An example of a completed section of **T.7.1** is shown in Table 9 below. Additional guidance is provided in the subsections that follow. To estimate snags, consider all snags within the transect, including 10m on all sides. To estimate CWD, only consider those pieces that intersect your rotary tape through the center of the transect.

FULL POLYGON ASSESSMENT OBSERVATIONS							
Question Indicator	TRANSECT SUMMARY INFO IN BOTH WETLAND AND 10 m UPLAND FROM WETLAND EDGE	T1	T2	T3	Transect Summary	Large non-homogenous patches outside transects but within assessment polygon (% of polygon)	Total = Transect summary plus large patches
9a	% Dead or Decadent trees of all trees (pg.37)	10	20	15	Weighted Average % 15	NA	15%
9c	Old (O) or New(N) Coarse Woody Debris (>7.5 cm diameter) – that crosses transect % old to new (pg.35) <i>Note: Tally O and N within each transect for CWD that intersects with rotary tape.</i>	O,N, O,O	O,N N	O,O, O	% Old of Total 70	NA	70%

Table 9. Snags and Coarse Woody Debris Indicators.

9. a) Does the density of standing dead trees (snags) and decadent trees (dying trees) in the upland area and forested wetland areas (i.e., forested bogs and swamps) appear within the range of natural variability, not counting catastrophic events such as wildfire, i.e. neither too much or too little?

The amount of decadent and dead woody material (snags) on a site can be an indicator of overall health. Large amounts of decadent and dead woody material may indicate a reduced hydroperiod due to either human or natural causes. De-watering of a site, if severe enough, may change the site vegetation potential from riparian/wetland species to upland species. In addition, decadent and dead woody material may indicate severe stress from over browsing. Finally, large amounts of decadent and dead woody material may indicate climatic impacts, or disease and insect damage. For instance, severe winters may cause extreme die back of trees and shrubs, and cyclic insect infestations may kill individuals in a stand. In all these cases, a high percentage of dead and decadent woody material reflects degraded

vegetative health, reducing production and other wildlife values, and, if along a channel, stream integrity. The most common usage of the term decadent may be for over mature trees past their prime and which may be dying, but we use the term in a broader sense. We count decadent plants, both trees and shrubs, as those with 30% or more dead wood in the upper canopy. In this item, scores are based on the percentage of total woody canopy cover which is decadent or dead, not on how much of the total polygon canopy cover consists of dead and decadent woody material. Only decadent and dead standing material is included, not that which is lying on the ground. The observer is to ignore (not count) decadence in poplars or cottonwoods which are decadent due to old age (rough and furrowed bark extends substantially up into the crowns of the trees), because cottonwoods/poplars are early seral species and naturally die off in the absence of disturbance to yield the site to later seral species. The observer is to consider (count) decadence in these species if apparently caused by de-watering, browse stress, climatic influences, or parasitic infestation (insects/disease). The observer should comment on conflicting or confounding indicators, and/or if the cause of decadence is simply unknown (but not due to old age). Do not count plants installed by human planting that are less than one year old as dead/decadent.

9. b) Are the standing dead trees composed of different diameter and decay classes?

Record all diameter and decay classes observed in each transect using **T.7.3** of the field form. See Figure 6 and Table 10 below for decay class and diameter size class tables respectively. Mark “Yes” if there are three or more decay classes and more than two diameter classes present.

T.7.3. Continuous observations: snag and coarse woody debris within wetland and 10 m upland											
Softwood						Hardwood					
Dead						Dead Fallen	Dead			Dead Fallen	
Hard →			Spongy	→ Soft			Not Sampled	Hard →	Spongy	→ Soft	Not Sampled
3	4	5	6 2/3 original height	7 1/2 original height	8 1/3 original height	9	3	4	5	6	
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	
Check all decay classes of snags observed within 10 m upland of the wetland edge											
Check all Diameter Sizes observed: 0 < 10 cm <input type="checkbox"/> ; 10 < 20 cm <input type="checkbox"/> ; 20 < 30 cm <input type="checkbox"/> ; 30 < 40 cm <input type="checkbox"/> ; 40 < 50 cm <input type="checkbox"/> ; 50 cm + <input type="checkbox"/>											

Figure 6. Visual appearance codes for wildlife trees. Copied from Forest and Range Evaluation Program Protocol for Stand-level Biodiversity Monitoring (Appendix 6).

Class	Diameter (cm)
1	0<10
2	10<20
3	20<30
4	30<40
5	40<50
6	50+

Table 10. Tree diameter classes.

9. c) Is more than half of the coarse woody debris that is present stable and well incorporated into the wetland, with no evidence of recent movement?

“Old” wood is wood that is stable and well incorporated into the wetland. It is usually mossy though this is not critical. “New” wood is any wood that is not stable or well incorporated into the wetland. New wood is usually recently deposited after road building and the most recent harvesting. However, new wood could also be old wood that was once stable, but has recently moved and is no longer stable. Most new wood will probably be wood that was introduced as a result of the treatment being examined. In most cases, this will be wood that has been introduced where there has been falling and/or yarding within or adjacent to the wetland. Windthrow is also new wood if it happened after the treatment (logging) and fell into the wetland. Blowdown from trees killed by insects, fire and/or self-thinning can also be new wood.

9. d) Does the coarse woody debris present show distinct multiple modes with regard to diameter and decay?

Record all diameter classes observed in **T.7.3** of the field guide. Also record all decay classes for CWD observed in each transect (see Table 9 below). Mark “Yes” if there are three or more decay classes and two or more diameter classes present.



9b: # decay classes of standing dead trees: _____		# Diameter Size Classes of standing dead trees: _____				
						
	Class 1	Class 2	Class 3	Class 4	Class 5	
Wood Texture	Hard	Sap rot (but still hard, thumbnail penetrates)	Advanced decay (spongy/large piece)	Extensive decay (crumbly-mushy)	Small pieces, soft portions	
Portion on Ground	Elevated on support points	Elevated but sagging slightly	Sagging or broken	Fully settled on ground	Partly sunken	
Branches	Hard branches with twigs	Soft branches	Branches/stubs absent	Absent	Absent	
Bark	Firm	Loose	Trace	Absent	Absent	
Wood Appearance	Fresh/recent	Colour fading	Fading colour	Light or brown	Reddish brown	
Wood Strength	Supports person	May not support person	Breaks easily. Pieces snap	Collapses with weight. Pieces do not snap	Feels firm like ground	
Invading Roots	None	None	In sapwood	In heartwood	In heartwood	
Check all decay classes of CWD observed:	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	
Check all Diameter Sizes observed:	0 < 10 cm <input type="checkbox"/> ; 10 < 20 cm <input type="checkbox"/> ; 20 < 30 cm <input type="checkbox"/> ; 30 < 40 cm <input type="checkbox"/> ; 40 < 50 cm <input type="checkbox"/> ; 50 cm + <input type="checkbox"/>					
9d: # decay classes of coarse woody debris: _____		# Diameter Size Classes of coarse woody debris: _____				

Table 11. Decay classes for coarse woody debris. Adapted from LMH 25: Describing Terrestrial Ecosystems in the field (Ch. 8, p27)

10. Has vegetation around the wetland been adequately protected from windthrow 10 m upland of the wetland?

Windthrow is defined as any live dominant or codominant tree that is sheared off (“wind snap”) or tipped over (“blow down”) by a strong wind, and no longer self-supporting. If the stem is snapped off, the stump that the stem was attached to should still be identifiable. Dead trees, snags or understory trees that get blown over or sheared off do not count as windthrow. Excessive windthrow in riparian areas can cause a number of problems, including lower terrain stability, greater bank or shoreline erosion, increased sedimentation, blockages to large animal movements, higher fuel loads, or more sites for insect or fungal infestations, to name just a few.

The answers to the indicators in Question 10 may seem obvious, for example, the riparian area of concern may be clearcut in which case there is no windthrow. Or it may be obvious that the amount of windthrow is clearly under (no windthrow) or over (new windthrow >50%) the thresholds. In either case, there is no need for additional information.

At other times, it may be necessary to calculate the difference between old and new windthrow. Do this by estimating or counting the number of old and new windthrown trees present in the upland area of the transects. Use the transect data to determine percent new (“post-treatment”) windthrow, percent old (“pre-treatment”) windthrow, and the difference between them, as follows:

$\% \text{ new windthrow} = (\text{number of new windthrow}) / (\text{number of new windthrow} + \text{number of standing trees}) \times 100$

$\% \text{ old windthrow} = (\text{number of old windthrow}) / (\text{number of old windthrow} + \text{number of new windthrow} + \text{number of trees still standing}) \times 100$

Complete these calculations for each transect, and then apply a weighted average (using Fb) to determine the amount of new and old windthrow within the upland area around the wetland. Use the following equation to determine the percent windthrow over and above what occurs naturally.

$\% \text{ windthrow over and above what occurs naturally} = \% \text{ new windthrow} - \% \text{ old windthrow.}$

Use the workspace at the bottom of T.5.2. to complete all the necessary equations.

To count as old (“pre-treatment”) windthrow, as opposed to “ancient windthrow”, the tipped over stumps should still have a distinct rootwad with at least a few main branch roots visible. If the stem is snapped off, the stump that the stem was attached to should also still be identifiable. Additionally, only count the windthrow with a rootwad within the transect that is being considered. **NOTE:** Only consider mature trees that were alive at the time of windthrow.

Based on information provided above, answer either 10 a or 10 b as applicable, depending on if the wetland contains an Riparian Reserve Zone (RRZ).

10. a) Is the incidence of post-treatment windthrow 20 m upland around small (0-5 ha) wetlands with no RRZ less than 10% of the living stems over and above what occurs naturally in the area?

10. b) Is the incidence of post-treatment windthrow in the RRZ of the riparian area around large (>5 ha) wetlands or small (0-5ha) with a RRZ less than 5% of the living stems present over and above what occurs natural in the area?

10. c) Are wildlife trees (e.g., nest sites, bear dens) still standing, or if not, still functioning as wildlife trees?

If any wildlife trees, are observed that were damaged during the treatment (e.g., logging), then document the impact, and record the number of trees impacted in T.7.1., and mark N for this question. This question requires direct observations of impacts to trees that showed evidence of supporting wildlife species (e.g., nesting cavities, nests, etc.)

Hydrology

11. Is vegetation in the wetland and its riparian area free of any impacts due to changes in the hydrologic regime?

The presence of water, and the depth, timing and duration within the rooting zone strongly influence the type of wetlands that occupy a site. Even slight changes in water drawdown or inundation over the long term can modify the wetland plant association, whereas extreme changes in the hydrological regime can even lead to wetland loss.

11. a) Are hydrologic changes minor or non-existent?

Although water levels naturally fluctuate on a seasonal basis in most systems, many wetland systems are affected by human-caused (artificial) additions or withdrawals. This artificial change of water level rarely follows a temporal regime that maintains healthy, native wetland plant communities. The result is often a barren band of shore exposed or inundated for much of each growing season. This causes shore material to destabilize, and often provides sites for weeds to invade. Such conditions are extremely detrimental to healthy riparian function. Not all wetlands evaluated will have surface water potential, but any wetland may have its water table degraded by draining, pumping or diverting its surface or subsurface supply. On such lentic wetlands as marshes and wet meadows, look for evidence of drainage ditching, pumping, and the interruption of normal surface drainage inputs by livestock watering dugouts, cross slope ditches, or upslope dams. For this indicator, the evaluator is asked to categorize the degree to which the system is subjected to artificially rapid or unnaturally timed fluctuations in water level. Reservoirs intended for storage of water for power generation, irrigation, and/or livestock watering typically exhibit the most severe effects, but water may also be diverted or pumped from natural systems for many other reasons (domestic, agricultural or industrial use). This indicator requires the evaluator to make a subjective call by choosing as a best fit one of the categories of drawdown severity described below. **NOTE:** Be careful to consider the scale of the water body as it relates to the scale of change. Pumping a small dugout full of water for livestock might severely impact a 0.8 ha (2 acre) slough but be negligible to a lake covering a larger section of land. Be sure to document the grounds for your estimate. If there is no way to know with any reasonable degree of certainty how much water is being added or removed to gauge the level of severity, it may be better to describe the situation but conservatively answer “Yes” to this question.

Severity Categories of Lentic Water Level Manipulation

Not Subjected – The water body or wetland is not subjected to artificial water level change (e.g., drawdown, addition, stabilization, etc.). This category may include very small amounts of change that cause no detectible fluctuation in water level.

Minor – The water body or wetland is subject to no more than minor artificial water level change. The shore area remains vegetated, and withdrawal of water is limited or slow enough that vegetation is able to maintain growth and prevent soil exposure. A relatively narrow band affected by the water level fluctuation may support only annual plants.

Moderate – The water body or wetland is subject to moderate quantities, speed and/or frequency of artificial water level change. Where water is removed, it is done in a way that allows pioneer plants to vegetate at least half of the exposed area resulting from drawdown. Where water is added, some flooding may occur at levels or times not typical to the area/season.

Extreme – The water body or wetland is subjected to extreme changes in water level due to volume (extent), speed and/or frequency of artificial water addition or removal. Frequent or unnatural levels of flooding occur where water is added, including extensive flooding into riparian and/or upland areas, or no natural annual drawdown is allowed to occur. In extreme artificial drawdown situations, a wide band of exposed bottom remains unvegetated.

11. b) Are recent dead trees or shrubs absent from the wetland edge?

The presence of dead trees or shrubs may indicate increased inundation, especially for species that are typically more terrestrial and were originally above the water table. Alternatively, this could be due to excessive drying out. Mark “Yes” if there is not a distinct band of dead shrubs or trees within or around the edges of a wetland. If the water level has risen, the observer may also identify younger, more water hydrophytic plants establishing in close proximity.

11. c) Is the wetland free of progressively younger age class plants or trees extending into the wetland from the drier edges of the wetland or adjacent upland area?

Encroachment of upland plants may occur if the wetland experiences a drawdown in the water table, or if the wetland becomes filled in (e.g., from sediment deposits). Mark “Yes” if there is not a clear progression of younger age class plant species extending into the wetland from the drier edges of the wetland or adjacent upland area. Wetland associated plants may display stress and less vigor in close proximity as water is taken up by these new propagations.

11. d) If the wetland has a defined stream flowing through it, does any incisement of the channel have only a minor to non-existent effect on the wetland vegetation?

An incised stream channel has experienced vertical downcutting of its bed. Incisement can lower the water table enough to change vegetation site potential. It can also increase stream energy by reducing sinuosity, reduce water retention/storage, and increase erosion. A stream becomes critically incised when downcutting lowers the channel bed so that the two-year flood event cannot overflow the banks.

Some typical downcutting indicators include: headcuts, exposed cultural features (pipelines, bridge footings, culverts, etc.), lack of sediment deposits, exposed bedrock, and a low, vertical scarp at the bank toe on the inside of a channel bend.

A severe disturbance can initiate downcutting, transforming the system from one having a high-water table, appropriate floodplain, and high productivity to one of degraded water table, narrow (or no) active floodplain, and low productivity.

Incisement Class

None – The channel is vertically stable and not incised; 1-2 year high flows can access a floodplain appropriate to the stream type. Active downcutting is not evident. A mature, or nearly mature, vegetation community occupies much of the floodplain.

Slight – The extent of incisement is minimal. A 2-year flood event may still access some floodplain, partially or in spots. Any new floodplain is established with perennial vegetation. Downcutting may be progressing or there may be ongoing lateral erosion of high side walls.

Moderate – The extent of incisement is significant. Floods of 1-5 year magnitude cannot reach a floodplain. The channel may look like a gully. Active downcutting or high side walls are likely actively eroding. If new floodplains are establishing, they are only starting to become established with perennial vegetation.

Severe – The extent of incisement has caused a deep entrenchment. Small to moderate floods cannot reach the original floodplain. There is no floodplain at the bottom. Downcutting may still be occurring, or high side walls are eroding. Sediments are flowing downstream.

11.e) Natural surface or subsurface areas within the wetland are not altered by disturbance. If drainage tiles, ditches, dikes, or gullies are present, they are having a minor to non-existent impact to vegetation in the wetland.

Mark “Yes” if there are no active ditches or underground drainage tiles present that are moving water away from the site and affecting changes in vegetation. Ditches are sometimes created along the toe of a slope, and these may divert water from the valley bottom. Ditches may also traverse through a wetland. They are usually straight and don’t display the natural sinuosity of streams typical of gradual slopes on valley bottoms. Underground manmade drainage may include buried logs, rocks, plastic or kiln-dried clay structures used to convey water underground and effectively lower the water table. These drainages are not easily observed but are sometimes apparent where they empty into open channels downstream. Look in these areas for any structures that may be visible. Mark “Yes” if there are no nearby dikes or roads that are preventing floods of surface water from entering the wetland.

12. Is there an absence of significant threats to water levels in the wetland?

12.a) If present, is the outlet structure stable and allows water to pass securely?

The field crew should make an effort to visit wetland outlets to determine how water leaves the site and check the integrity of the outflow area. Both natural and constructed outlets play an important role in holding water within the wetland. If water exits the wetland through overland flow, it’s important to determine whether or not the channels are susceptible to erosion. Outlets should be composed of stable materials (e.g., embedded logs, rock, dense roots, and hardened surfaces). Dams are not automatically considered an impact. If dams are present, look for signs of instability. Animals burrowing into the dam, water flowing over the dam and causing erosion, or erosive forces on the upslope side of the dam can weaken its integrity. If the dam is leaking and water is seeping on the toe of the downstream slope, coloured water can indicate that sediment from the dam is leaving the site and weakening the dam. Human-made dams often include designated spillways that allow water safe passage that bypasses the dam and control maximum water levels. Check for blockages in the spillway or any erosion in the spillway. Mark “Yes” to this indicator question if the outlet structure lacks animal burrows, erosion or leakage.

12. b) If the wetland has a channel, is there no presence of active headcuts below or within the wetland (i.e., locations of active downcutting in channel)?

Check downstream for headcuts or the susceptibility for headcuts. Headcuts are “mini waterfalls” in water channels, where shear stress is high, and erosion occurs in soft materials. Headcuts will travel upstream until they reach a harden surface that prevents further erosion (e.g., bedrock). If a headcut reaches a wetland, it can continue through the wetland and cause the wetland to drain by acting like a ditch.

Mark “No” if any threats as indicated above are present.

12. c) Is less than 15 % of the shoreline of the wetland or any stream channels flowing through the wetland disturbed in any way?

For wetlands with defined streambanks: Structurally altered streambanks are those having impaired structural integrity (strength or stability). These banks are more susceptible to cracking and/or slumping. Count as streambank alteration such damage as livestock or wildlife hoof shear and concentrated trampling, vehicle or ATV tracks, and any other areas of human-caused disruption of bank integrity, including rip-rap or use of fill. The basic criterion is any disturbance to bank structure that increases

erosion potential or bank profile change. One exception is lateral bank cutting caused by stream flow, even if thought to result from upstream human manipulation of the flow. The intent of this indicator question is to assess only direct, on-site mechanical or structural damage to the banks. Each bank is considered separately, so total bank length for this indicator is approximately twice the reach length of the stream channel in the polygon (more if the stream is braided). **NOTE:** Constructed streambanks (especially those with rip-rap) may be stabilized at the immediate location but are likely to disrupt normal flow dynamics and cause erosion of banks downstream. The width of the bank to be assessed is proportional to the stream size as per the following conceptual guidelines.

Band width along the bank to be assessed for disturbance:

Rivers larger than 15 m channel width: assess first 15m from top of bank.

Small rivers and large streams 5-15 m channel width: assess first 5 m from top of bank.

Small streams up to 5 m channel width: assess 2m from top of bank.

For lentic wetlands with open water features: Record the percent of shoreline that is actively eroding or that shows any disturbances that increase the susceptibility of the shoreline to erosion. This may include loss of deep-rooted vegetation (e.g., woody rooted species or robust emergent vegetation), loss of rocks greater than 2.5cm in diameter, or loss of embedded coarse woody debris. Estimate the percentage of shoreline impacted compared to the shoreline length within the polygon assessed.

12. d) If the wetland has a stream channel, does the channel bank have vegetation with greater than 65 % deep binding rootmass?

All tree and shrub species are considered to have deep binding root masses. These species help stabilize the soil and filter sediments from overland flow. A rule of thumb in wetland herbaceous species is that perennial species have a range of root mass qualities, while annual plants lack deep binding roots. Some rhizomatous species such as the deep rooted *Carex* species (sedges) are excellent bank stabilizers. Others, such as *Poa pratensis* (Kentucky bluegrass), have only shallow roots and are poor bank stabilizers. Still others, such as *Juncus balticus* (wire rush), are intermediate in their ability to stabilize banks. The size and nature of the stream will determine which herbaceous species can be effective. The evaluator should try to determine if the types of root systems present in the polygon are in fact contributing to the stability of the streambanks. In situations where you are assessing a high cut bank (like the top of a valley wall), the top may be represented by an upland vegetation community, but the bottom may be a mix that is more representative of a riparian vegetation community. Do not assess the area that is only upland. In cases of tall, nearly vertical cut banks, assess the bottom portion that meets floodwaters. Omit from consideration those areas where the bank is comprised of bedrock, since these neither provide binding root mass, nor erode at a perceptible rate. **NOTE:** Rip-rap does not substitute for, act as, or preclude the need for deep, binding root mass.

Rivers larger than 15 m channel width: assess first 15m from top of bank.

Small rivers and large streams 5-15 m channel width: assess first 5 m from top of bank.

Small streams up to 5 m channel width: assess 2m from top of bank.

Water Quality

13. Does the water quality of the wetland appear to be within a reasonable range of natural variation?

Wetlands are well recognized as “sinks” for absorbing pollutants and sediment. Water leaving a wetland is often cleaner than water that enters a wetland. Streams entering wetlands often drop suspended sediments as water slows down and flow energy dissipates. The presence of extensive surface areas on emergent and submergent plants plus a mix of aerobic and anaerobic environments allow for a diversity of microorganisms and bacteria to break down and absorb pollutants. If clay soils are present in the basin, they contain zeolites for greater absorptive capacity. Many pollutants moving through wetlands at low levels can go undetected and are only observable in the field when the water quality is severely degraded. Mark “No” for Question 13 if either of Questions 13a or b is answered “No”.

13. a) Does the wetland lack any signs of excessive nutrient loading such as algae mats, blooms, fish kills?

If fish kills, algae blooms or mats are observed in the wetland, it typically indicates an overabundance of nutrients. High nutrients (e.g., phosphorous) often lead to increased microbial activity, which lowers the availability of oxygen in the water column and subsequently causes fish kills. High nutrients also increase algae growth.

13. b) Do basic water quality parameters (smell, colour, pH, turbidity, temperature) appear to be within a reasonable range of natural variation?

Where water is present at the surface, or in the test hole, use a calibrated water probe and record pH and temperature. Take the measurement three times and record the average. Consider the wetland class and determine if the pH is within the expected range for that particular wetland. Bogs = pH < 5.5, fens = pH > 5, swamps = pH < 7, and marshes = pH > 6.5. Shallow open water can have a broad range of pH values. If you have further classified your site to a wetland plant association, you can refer to *Wetlands of British Columbia: A Guide to Identification* (Mackenzie and Moran, 2004) to see a narrower range of expected pH values.

In organic soils that are fibric or mesic, bogs are more acidic than fens. In mineral soils or humic organic soils, swamps are more acidic than marshes. The ranges of pH as shown in Figure 7 below are: very acidic (VA) = pH < 4.5; moderately acidic (MA) = pH 4.5 to 5.5; somewhat acidic (SA) = pH 5.5 to 6.5; neutral (N) = 6.5 to 7.4; and alkaline (Ak) > 7.4.




Figure 7. Edatopic grid for abiotic factors and their relationship with different aquatic classes. Source: Wetlands of British Columbia, LMH 52.

Consider the smell, colour and turbidity of any open water. Record “No” if one or more of the water quality parameters appear abnormal. Keep in mind that some processes are normal. If there is an oily sheen, smell it to determine whether there is a petroleum odor. Natural oils can accumulate on the surface of water from the decomposition of dead plant material. If poked, plant oils will fracture like shattered glass, whereas petroleum will re-amalgamate relatively quickly. Wetlands can also naturally produce hydrogen sulfide, which releases a rotten egg smell due to anaerobic conditions.

Landscape

14. Is the riparian and upland habitat beside the wetland of adequate size and quality to mitigate impacts on critical activities (movements, feeding, breeding) by the area’s desired wildlife (e.g., grizzly, ungulates, martin, raptors, woodpeckers, songbirds, waterfowl, reptiles, amphibians, etc.)?

Successful use of an area by wildlife tends to be related to the size and quality of the habitat available. The relationship, however, is seldom simple because habitat quality can be exceedingly variable or difficult to summarize for large areas. In this assessment, wetlands are assumed to be important components of wildlife habitat, and their value is strongly related to how visible (or remote) the wetland is to predators and hunters, as well as how connected the wetland is to suitable amounts and types of riparian and upland habitat.

If the remote data available is sufficiently recent and accurate, most of the indicators can be answered in-office using GIS or visual quality assessment software. Field reviews, however, will probably still be needed to confirm the accuracy of the remote data. The visibility (or remoteness) of the wetland in particular (Indicator b) may be best assessed from various vantage points in the field.

14. a) Does 75% or more of the wetland have an upland area around it that has not been modified by human activities, 30 m wide in the case of small wetlands (0-5 ha), or 50 m wide for large (> 5 ha) wetlands?

For this question, consider harvesting, trails, roads, access by cattle. This calculation can be first estimated from remote imagery, and then further validated in the field.

14. b) Is 10% or less of the wetland's shoreline visible from any point on a road, pipeline or power line within 300 m of small wetlands (0-5ha), or 500 m of large wetlands (>5ha)?

The visibility (or remoteness) of the wetland is best assessed from various vantage points in the field. Prior to heading out to the field, study the area and identify right-of-ways of interest to investigate. If you encounter one point where you can see more than 10% of the wetland polygon, this is sufficient to record a “No” response. For trip planning, attempt to select areas where there appears to be narrow retention or heights of land that would provide a predator (including hunters) with an improved vantage point. In the example below, the purple lines indicate roads, the white dashed line is the distance measure, and the yellow lines are contours at 20 m intervals (for determining heights of land).




Figure 8. Map of right-of-ways within 500 m sight distance of a large (>5ha) wetland used to identify potential vantage points.

14. c) Do right-of-ways within 100 m of the wetland impinge on no more than 10% of the wetland's perimeter?

This can be visually estimated with air photos or satellite imagery. Calculations can also be made with mapping software or the FREP mapping tool for government employees. In the example below, the dashed white line is the 100 m buffer around the wetland. The perimeter of the wetland is 2377 m. Approximately 405 m of the wetland perimeter has a road within 100 m along its length. In this case, the road impinges on 17% of the wetland perimeter, thus the evaluator would record a “No” response.



Figure 9. Example of road impinging within 100 m of wetland area along the southwest (Q 14c).

14. d) *Is the percent of mature and old forest within two kilometers of the wetland perimeter greater than the minimal target for the area's biogeoclimatic zone and natural disturbance type?*

Landscape connectivity helps provide wildlife with sufficient and different habitat types to adequately meet their life needs. Gaps in connectivity can increase risk of predation, reduce suitable habitat, and ultimately limit chances of survival. The amount of intact habitat required can vary among species. The following indicator was adapted from a broadly utilized methodology in North America to estimate landscape connectivity around wetlands and other landscape features of interest. A similar methodology is utilized by the Conservation Data Centre of BC. This FREP wetland health evaluation applies a threshold of percent of mature and old forests that surround a wetland as a proxy of landscape connectivity. The targeted percent of mature and old forest varies by the site's natural disturbance regime and biogeoclimatic zone. Appendix 4 provides the minimum targets.

This indicator is best completed as a desktop exercise utilizing the best available wetland polygon that delineates the wetland boundaries.

Steps:

1. Place a 2000 m (2 km) spatial analysis buffer around the wetland.
2. Estimate the percent of old and mature forests within the buffer. Tip: If you have access to GIS software (e.g., QGIS), you can run a query on the Vegetation Resource Inventory spatial layer. Select or extract polygons within the fields: Forest Management Land Base "FOR_MGMT_LAND_BASE_IND" == Y and that have an age "PROJ_AGE_1" greater than the targeted age. For government employees, this can be achieved using the filter and VRI reporting

widgets in FREPmap. For targets, use the table in Appendix 4 based on the wetland’s dominant biogeoclimatic zone and natural disturbance type. If you don’t have access or the ability to conduct this type of spatial analyses, an estimate using aerial imagery and any other supporting information is sufficient.

3. Calculate the proportion of the polygon that meets the criteria.




Figure 10. Example of a spatial query for mature and old forests within a 2 km buffer of the wetland.

In the example shown in Figure 10, the wetland is within a natural disturbance type NDT3 and the biogeoclimatic zone and variant is SBSmc2. From the table in Appendix 4, the target mature and old forest is greater than 34%, and the minimal age is 100 years. The proportion of polygons meeting these criteria is 47.6% of the forest land base, therefore the wetland has sufficient mature and old forest.

15. Are surface and subsurface flows to the wetland intact?

Roads and other right-of-ways (ROWs) can have a significant impact on wetlands by diverting the flow of both surface and shallow ground water. The following questions are coarse-level indicators of potential impacts.

15.a) Do all mapped and unmapped streams at roads and ROWs appear to be in their original water courses?

Field surveyors should drive or otherwise traverse the roads and ROWs upslope of the wetland whenever possible. If available, this evaluation can be conducted using drone imagery. Look for mapped and unmapped streams and groundwater exposures where there have been cuts. Unmapped streams or groundwater exposures (i.e., new springs) are typically not part of the planning phase and may contribute significantly to changes in water inputs. If surface water is diverted upslope and away from its original trajectory (i.e., not provided with sufficient passage through a culvert or bridge), record “No”. Look for

drainage ditches alongside the road; look for flowing water, or evidence of flow, that suggests water is taken away from its natural course. If the wetland is adjacent to a lake or a river, only evaluate the face drainage that enters the wetland and not the drainage for the entire lake or river.

15.b) Is less than 25% of the contributing basin intercepted by roads or ROWs?

This is a map-based exercise that is best initiated before the field visit. It requires the use of topographic maps to estimate the hydrology and road layers. Record the amount of the basin that is separated from the wetland by a ROW.

If the surveyor observes additional roads or ROWs that were not on the map, the surveyor needs to draw in/digitize the location of this missing road on the map and re-calculate the percentage of the wetland drainage area potentially affected.

The contributing basin is the land where surface waters would flow into the wetland via streams or sheet flow on the surface or subsurface. First, estimate the area of the contributing basin on a map by interpreting surrounding streams and elevation contour lines or hill shade imagery. Next, estimate the amount of the contributing basin intercepted by ROWs.

Example: The white area in Figure 11a is a predefined sub-watershed area from the BC Freshwater Atlas. The light pink polygon is the approximate contributing basin for the wetland (light blue/green polygon). Both streams (blue lines) and contour lines (brown lines) were used to approximate the contributing basin. The areas within the contributing basin intercepted by roads (black lines) is displayed as dark pink in Figure 11b. In this example 501 out of 631 hectares are intercepted by right of ways (i.e., 79%).




Figure 11. A) Contributing basin shown as polygon in light pink; B) polygon with flow intercepted by ROW in dark pink.

Appendix 1 – Recommended Field Sampling Sequence

Introduction:

Steps 1 through 4 describe the process for completing your transect. However, there are other tables that have “continuous” information – you will fill these tables out AFTER you’ve completed all your transects. It’s helpful to mentally record, take photos, or start tallies (as appropriate), so that you can recollect what you’ve seen.

Continuous tables include:

- T.5.3. Bare Ground Hydrologically Connected to the wetland
- T. 6. Structural features missing or impacted within 10 m upland from wetland edge, and the condition of those features (the Form, Vigor and Recruitment)
- T.7.2. Distribution Code of Invasive Species
- T. 7.3 Observations about the Diversity of Decay and Size Classes of Snags and Coarse Woody Debris
- T.7.4. Impacts to vegetation (i.e., impacts that may prevent the vegetation community from achieving it’s climax community, pressures from overgrazing/browsing)
- T. 8. Structural features missing or impacted within the wetland, and the condition of those features (the Form, Vigor and Recruitment)
- T. 9. Physical and Hydrological Impacts to the Wetland. Take a moment to review questions within this table. Ensure you have an opportunity to assess the outlet, inlet, and any significant channel that is within the wetland.
- In addition, while approaching your sampling location, or walking around the wetland, observe any anomalies that are significant stressors that may be located outside of your transect (i.e., large patches of bare soil, invasive plants, disturbance/increaser species, low retention within RMZ or high disturbance in RMA, harvesting in wetland, dead or decadent trees, excessive windthrow or newly deposited coarse woody debris). Record anomalies that may affect your overall score in the second last column of Table 5.1 and Table 7.1, titled: “Large non-homogenous patches outside transect but within upland area/assessment polygon”.
- Step 1. At your sampling location, mark edge of wetland boundary with flagging tape. Record UTM Coordinates in T.4.1., you will fill out other columns in this table later.
- Step 2. Run a transect line upland from the wetland edge, and mark 10 m & 20m upland perpendicular from wetland edge with flagging tape. While you have your transect line laid out and you are exploring your upland location:
 - In Table 4.3., document invasive or disturbance species within the first 10 m upland from wetland edge. Make % estimates based on 10 m on either side of the transect line.
 - In Table T.5.1., there are 3 columns labeled T1-T3, one for each transect you will sample. Complete a transect column. Make % estimates based on 10 m on either side of the transect line for Question Indicators 3 and 5.
 - Table T.7.1. is completed based on the full transect length (i.e., 10 m upland plus wetland observations). Mentally take note of question indicators 1 to 9a in T.7.1.
 - Start recording old (O) and new (N) coarse woody debris that cross your transect line (i.e., Question Indicator 9c) in T.7.1. Note: if woody debris extends into the wetland you will need to continue recording CWD once you extend your transect into the wetland (Step 4). While observing coarse woody debris, also check off the decay class and diameter class in T.7.3. Once you’ve checked off a decay class or diameter range, you don’t need to check off the same class or diameter range again (this is a continuous observation).
 - If there is new windthrow (since disturbance), then also begin to make a tally for Question Indicators 10a,b. in T.5.2. Note: consider 20 m upland from wetland edge for wetlands without an RRZ, and 10 m upland from wetland edge for wetlands with an RRZ.

- In Table 7.3. also check off the decay and diameter classes of snags (dead standing trees) 10 m on either side of you.
- Reel in your rotary tape. You will now explore the wetland.
- Step 3. Run a transect line into the wetland that is perpendicular from the wetland perimeter. One of the crew members may stand on the edge of the wetland perimeter to guide the other into the wetland on a compass bearing and to help record notes.
 - In Table 4.1. document the transect bearing, end type (e.g., end of 50 m, channel, etc.), and Transect Length. Add 10 m to account for the upland portion. If you are entering a new wetland plant community at the end of your transect, then you may also add an additional 10 m to the transect length on the other end so that you can incorporate your observations of this new plant community (this may occur if you encounter, for instance, Shallow Open Water and are unable to walk further). Once you reach the extent of the transect, leave the rotary tape laid out. Remember that your transect represents one side of your wetland. Your transect length cannot be closer to any other part of the wetland perimeter than from where you placed your flagging tape for this transect at the wetland edge (Otherwise you are recording observations that are likely influenced from a different edge of the wetland).
 - Continue recording old (O) and new (N) coarse woody debris that cross your transect line (i.e., Question Indicator 9c) in T.7.1. While observing coarse woody debris, also check off the decay class and diameter class in T.7.3.
 - If there is new windthrow (since disturbance), then continue to make a tally for Question Indicators 10a,b,c. in T.7.1.
 - In Table 7.3., check off the decay and diameter classes of snags (dead standing trees) 10 m on either side of you.
 - Complete plots within each distinct vegetation zone. Fill out Table 4.2 (a row per plot) and Table 4.3 concurrently. While completing Table 4.3, only record: dominant native species in addition to invasive species. An exhaustive list is not necessary for this assessment. If you cannot identify a species, then make a quick note (e.g., Salix Sp.), record percent cover, and then move on! This portion of the assessment should be quick. Dominant species should help you classify your wetland. If you are not confident about the wetland class or plant association, then leave it blank in T.2. You will still be able to assess your site. Along the length of the transect make observations either 10 m from the plot centre, OR if the band of vegetation is narrower than 20 m, then to the extent of the vegetation zone where there is a distinct break. Make observations 10 m on either side of the transect line. The intent is only to document observations that represent the vegetation zone you are trying to describe.
 - After you've completed the plots on your transect, complete the rest of your transect information (i.e., column T1,T2, or T3) in T.7.1., taking into account observations you made from the 10 m upland portion plus what you observed along the transect in your wetland portion. Imagine the transect as one strip that is 20 m wide (10 m on either side)
- Step 4. Repeat steps 1 through 3 until you've completed all your transects.
- Step 5. Fill out the Continuous Tables as described at the introduction of this document.
- Step 6. Complete the remainder of T.4.1. (i.e., the fractional information in the last 3 columns)
- Step 7. Complete the weighted averages in T.5.1 by using weights recorded as Fb in Table 4.1. Then add percentages from transects and anomalies in the last column.
- Step 8. Complete the weighted averages in T.7.1. by using weights recorded as WF in Table 4.1 for question indicators 1 to 9a. Then add percentages from transects and anomalies in the last column.
- Step 9. Fill out pages 12 to 15 based on your records.
- Step 10. Summarize your findings, score your wetland, and document causal factors on page 16 and 17
- Step 11. Prepare a small map that shows your transects and key features within your wetland on page 18.

Appendix 2 – Common Invasive and Noxious Plants in BC

Invasive/Noxious Plant Species ¹	Scientific Name	Wetland Indicator Status ²	Page Number ³
Anchusa (Common Bugloss)	<i>Anchusa officinalis</i>		pNA, P50
Baby's Breath	<i>Gypsophila paniculata</i>		p320, P73
Black Knapweed	<i>Centaurea nigra</i>		pNA, PNA
Blueweed	<i>Echium vulgare</i>		p76, P47
Bohemian Knotweed	<i>Fallopia x bohemica</i>		P26
Brazilian Waterweed (syn. Brazilian Eloda)	<i>Egeria densa</i>	OBL	PNA
Brown Knapweed	<i>Centaurea jacea</i>	FACU	pNA, PNA
Buck's-horn Plantain	<i>Plantago coronopus</i>	FACW	PNA
Bull Thistle	<i>Cirsium vulgare</i>	FACU	pNA, P75
Canada Thistle	<i>Cirsium arvense</i>	FACU	p32, P9
Caulerpa	<i>Caulerpa taxifolia</i>		PNA
Common Burdock	<i>Arctium minus</i>	FACU	pNA, P48
Common Tansy	<i>Tanacetum vulgare</i>	FACU	p58, P51
Cordgrass	<i>Spartina spp.</i>	OBL/FAC	P11-14
Creeping Buttercup	<i>Ranunculus repens L.</i>	FAC	P79
Dalmation Toadflax	<i>Linaria dalmatica</i>		p136, P6
Dandelion	<i>Taraxacum</i>		PNA
Didymo	<i>Didymosphenia geminata</i>	OBL	PNA
Diffuse Knapweed	<i>Centaurea diffusa</i>		P29, P74
Dwarf Eelgrass	<i>Zostera japonica</i>	FACU	PNA
Eurasian Watermilfoil	<i>Myriophyllum spicatum</i>	OBL	PNA
European Common Reed	<i>Phragmites australis spp</i>	OBL	P10
Fanwort	<i>Cabomba aquatica</i>		PNA
Field Scabious	<i>Knautia arvense</i>		P382, P52
Flowering Rush	<i>Butomus umbellatus</i>	FACW/OBL	P17
Giant Hogweed	<i>Heracleum mantegazzianum</i>	FAC	P19
Giant Knotweed	<i>Fallopia sachalinensis</i>	FACU	PNA, P27
Giant Mannagrass/Reed Sweetgrass	<i>Glyceria maxima</i>	OBL	P20
Giant Reed	<i>Arundo donax</i>		PNA
Gorse	<i>Ulex europaeus</i>	FACU	pNA, P21

Hairy Willow Herb	<i>Epilobium hirsutum</i>	FACW	PNA
Hanging Sedge	<i>Carex pendula</i>		PNA
Himalayan Balsam (syn. Policeman's	<i>Impatiens glandulifera</i>	FACW	P84
Himalayan Knotweed	<i>Persicaria wallichii</i>	FAC	P28
Hoary Alyssum	<i>Berteroa incana</i>		pNA, P55
Hoary Cress	<i>Cardaria draba</i>		P272, P56
Hound's-tongue	<i>Cynoglossum officinale</i>	FACU	pNA, 22
Hydrilla	<i>Hydrilla verticillata</i>		PNA
Japanese Knotweed	<i>Fallopia japonica</i>	FACU	p91, P29
Japanese Wireweed	<i>Sargassum muticum</i>		P84
Leafy Spurge	<i>Euphorbia esula</i>		p368, P30
Marsh Plume Thistle	<i>Cirsium palustre</i>		pNA, P60
Meadow Hawkweed	<i>Hieracium caespitosum</i>		pNA, PNA
Meadow Knapweed	<i>Centaurea debeauxii</i>		pNA, P57
Nodding Thistle	<i>Carduus nutans</i>	FACU	p26, P87
Orange Hawkweed	<i>Hieracium aurantiacum</i>		p40, P54
Oxeye Daisy	<i>Leucanthemum vulgare</i>	FACU	p30, P62
Parrots Feather	<i>Myriophyllum aquaticum</i>	OBL	PNA
Perennial Pepperweed	<i>Lepidium latifolium</i>	FAC	pNA, P63
Plumeless Thistle	<i>Carduus acanthoides</i>		p27, P67
Poison Hemlock	<i>Conium maculatum</i>	FACW	PNA
Policeman's Helmet	<i>Impatiens glandulifera</i>	FACW	PNA
Puncture Vine	<i>Tribulus terrestris</i>		p96, P64
Purple Loosestrife	<i>Lythrum salicaria</i>	FACW	p224, P35
Reed Canary Grass	<i>Phalaris arundinacea</i>	FACU	PNA
Rush Skeletonweed	<i>Chondrilla juncea</i>		pNA, P36
Russian Knapweed	<i>Acroptilon repens</i>		pNA, P58
Russian Olive	<i>Elaeagnus angustifolia</i>	FAC	PNA
Scentless Chamomile	<i>Matricaria maritima</i>		p48, P37
Scotch Broom	<i>Cystisus scoparius</i>		pNA, P88
Scotch Thistle	<i>Onopordum acanthium</i>		pNA, P69
Spotted Knapweed	<i>Centaurea biebersteinii</i>		p28, P25
St. John's Wort	<i>Hypericum perforatum</i>	FACU	pNA, P89
Sulphur Cinquefoil	<i>Potentilla recta</i>		p357, P66
Swollen Bladderwort	<i>Utricularia inflata</i>	OBL	PNA

Tansy Ragwort	<i>Senecio jacobaea</i>	FACU	p59, P39
Teasel	<i>Dipsacus fullonum</i>	FAC	pNA, P90
Variable-leaf Milfoil	<i>Myriophyllum heterophyllum</i>		PNA
Wakame	<i>Undaria pinnatifida</i>		PNA
Water Primrose	<i>Ludwigia hexapetala</i>		PNA
Wild Chervil	<i>Anthriscus sylvestris</i>		P70
Wild Parsnip	<i>Pastinaca sativa</i>		PNA
Yellow Flag Iris	<i>Iris pseudacorus</i>	OBL	pNA, PNA
Yellow Floating Heart	<i>Nymphoides peltata</i>	OBL	PNA
Yellow Nut Sedge	<i>Cyperus esculentus</i>	FACW	P34
Yellow Starthistle	<i>Centaurea solstitialis</i>		p29, P45
Yellow Toadflax	<i>Linaria vulgaris</i>		p138, P40

¹Sources for selected plant list: BC Wildlife Federation. 2015. Priority Aquatic Invasive Plants in BC. Factsheet. Invasive Species Council of BC. Personal Communication (March 2018).

²Page numbers following a lower case ‘p’ are given for the descriptions found in Weeds of Canada. 1999. Royer and Dickerson. Lone Pine Publishing and The University of Alberta Press. Page numbers following an uppercase ‘P’ are from the Field Guide to Noxious Weeds and Other Selected Invasive Plants of British Columbia. 2014. Cranston, et al. British Columbia.

³Sources for Wetland Indicator Status: United States Department of Agriculture. 2018. Plants Database. Wetland Indicator Status. <https://plants.usda.gov/wetinfo.html>; and W.H. MacKenzie and J.R. Moran. 2004. Wetlands of British Columbia: A Guide to Identification. Res. Br., B.C. Min For., Victoria, B.C. Land Manage. Handb. No. 52.

Appendix 3 – Select List of Culturally Significant Plants to Indigenous Communities in BC

Species Name	Common Name	Class
<i>Alnus rubra</i> Bong.	red alder	FACU
<i>Angelica genuflexa</i> Nutt.	kneeling angelica	FACW
<i>Apocynum cannabinum</i> L.	Indian hemp	FAC
<i>Argentina anserina</i> (L.) Rydb. (syn. <i>Potentilla anserina</i> L.)	common silverweed	FACU
<i>Argentina egedii</i> (Wormsk.) Rydb. (syn. <i>Potentilla egedii</i> Wormsk.)	Pacific silverweed	FACW
<i>Athyrium filix-femina</i> (L.) Roth. ssp. <i>cyclosorum</i> (Rupr.) C. Chr.	common ladyfern	FACU
<i>Betula nana</i> L.	scrub birch (syn. <i>Betula glandulosa</i> Michx.)	FACU
<i>Betula occidentalis</i> Hook.	water birch	FACW
<i>Betula pumila</i> var. <i>glandulifera</i> Regel	bog birch	OBL
<i>C. virosa</i> L.	Mackenzie's water-hemlock	FACW
<i>Camassia quamash</i> (Pursh) Greene	common camas	FACW
<i>Carex obnupta</i> L.H. Bailey	slough sedge	FACW
<i>Chamaecyparis nootkatensis</i> (D. Don) Spach (syn. <i>Cupressus nootkatensis</i> D. Don)	Yellow cedar	FAC
<i>Cicuta douglasii</i> (DC.) J.M. Coult. & Rose	Douglas' water-hemlock	FACW
<i>Comarum palustre</i> L.	purple marshlocks	OBL
<i>Conocephalum conicum</i> (L.)	cone-headed liverwort	
<i>Cornus sericea</i> L. (syn. <i>Cornus stolonifera</i>)	red-osier dogwood, or "red willow"	FACU
<i>D. anglica</i> Huds.	English sundew	OBL
<i>Drosera rotundi folia</i> L.	Round leaved sundew	OBL
<i>Elaeagnus commutata</i> Bernh. ex Rydb	silverberry	FAC
<i>Empetrum nigrum</i> L.	black crowberry, "blackberry"	FACU
<i>Equisetum arvense</i> L.	common, or field horsetail	FACU
<i>Equisetum hyemale</i> L. ssp. <i>affine</i>	scouring rush, or branchless horsetail	FACW
<i>Equisetum telmateia</i> Ehrh.	giant horsetail	FACW
<i>Eriophorum angustifolium</i> Honck., and <i>E. chamissonis</i> C.A. May.	cottongrass	OBL

<i>Frangula purshiana</i> (DC.) Cooper (syn. <i>Rhamnus purshiana</i> DC.)	cascara	FAC
<i>Fritillaria camschatcensis</i> (L.) Ker Gawl.	northern rice-root	FACW
<i>Hierochloe hirta</i> (Schrank) Borbás	northern sweetgrass	FACW
<i>Juniperus communis</i> L.	common juniper	FACU
<i>Kalmia microphylla</i> (Hook.) A. Heller; syn. <i>Kalmia polifolia</i> <i>Wangenh.</i>	bog laurel, or swamp-laurel	OBL
Kamchatka fritillaria	sarana lily	FACW/OBL
Labrador-tea, and <i>R.</i> <i>neoglandulosum</i> Harmaja (syn. <i>L.</i> <i>glandulosum</i> Nutt.) (syn. <i>Rhododendron groenlandicum</i>)	trapper's tea, or western Labrador tea	FACU
<i>Ligusticum canbyi</i> (J.M. Coult. & Rose) J.M. Coult. & Rose	Canby's lovage or Canby's licorice-root	FAC
<i>Lonicera involucrata</i> (Richardson) Banks ex Spreng.	black twinberry or twinberry honeysuckle	FACU
<i>Lupinus nootkatensis</i> Donn ex Sims	Nootka lupine	FACU
<i>Lycopus uniflorus</i>	bugleweed	OBL
<i>Lysichiton americanus</i> Hultén & H. St. John	American skunk-cabbage	FACW
<i>Malus fusca</i> (Raf.) C.K. Schneid.	Pacific crabapple	FACU
<i>Mentha arvensis</i> L.	field mint or Canada mint	FACU
<i>Myrica gale</i> L.	sweetgale	OBL
<i>Nuphar lutea</i> (L.) Sm. ssp. <i>Polysepala</i> (Engelm.) E.O. Beal	yellow water-lily	OBL
<i>Oenanthe sarmentosa</i> C. Presl ex DC.	Water parsley	OBL
<i>Populus balsamifera</i> ssp. <i>trichocarpa</i> (Torr. & A. Gray ex Hook.) Brayshaw	black cottonwood	FAC
<i>Platanthera hyperborean</i> (L.) Lindl. (syn <i>Platanthera aquilonis</i>)	northern green orchid	FACW
<i>Physocarpus malvaceus</i> (Greene) Kuntze	mallow ninebark	Not listed
<i>Platanthera stricta</i> Lindl.	slender bog orchid	FACW
<i>Phalaris arundinacea</i> L.	reed canary grass	FACU

<i>Phragmites australis</i> (Cav.) Trin. Ex. Steud.	common reed grass and related spp.	OBL
<i>Physocarpus capitatus</i> (Pursh) Kuntze	Pacific ninebark	FACU
<i>Picea mariana</i> (Mill.) Britton, Sterns & Poggenb.	black spruce	FACU
<i>Pinus contorta</i> Douglas ex Louden	lodgepole pine	FACU
<i>Platanthera dilatata</i> (Pursh) Lindl. ex Beck	white bog orchid	FACW
<i>Populus balsamifera</i> L. ssp. <i>balsamifera</i>	balsam poplar	FAC
<i>Potamogeton natans</i> L.	floating pondweed	OBL
<i>Rhododendron groenlandicum</i> (Oeder) K.A. Kron & W.S. Judd (syn. <i>Ledum groenlandicum</i> Oeder)	Labrador-tea	FACW
<i>Ribes bracteosum</i> Douglas ex Hook.	gray currant or stink currant	FACU
<i>Rubus arcticus</i> L. ssp. <i>Acaulis</i> (Michx.) Focke	Nagoonberry or dwarf raspberry	FACW
<i>Rubus chamaemorus</i> L.	cloudberry or bakeapple	OBL
<i>Rubus spectabilis</i> Pursh	salmonberry	FACU
<i>Rumex aquaticus</i> L. var. <i>fenestratus</i> (Greene) Dorn (syn. <i>Rumex occidentalis</i> S. Watson)	western dock	FACW
<i>Salix bebbiana</i> Sarg.	Bebb's willow	FACU
<i>Sagittaria cuneata</i> Sheldon	arumleaf arrowhead	OBL
<i>Salix discolor</i> Muhl.	pussy willow	FACW
<i>Salix hookeriana</i> Barratt ex Hook.	Hooker's willow	FACW
<i>Schoenoplectus lacustris</i> (L.) Palla [syn. <i>S. tabernaemontani</i> (C.C. Gmel.) Palla]	lakeshore bulrush or softstem bulrush	OBL
<i>Salix scouleriana</i> Barratt ex Hook.	Scouler's willow	FAC
<i>Sorbus sitchensis</i> Sanson ex Bong.	Sitka mountain ash	FAC
<i>Sagittaria latifolia</i> Willd.	wapato, or broadleaf arrowhead	OBL
<i>Salix barclayi</i> Andersson	Barclay's willow and other mountain willows	FACU
<i>Salix exigua</i> Nutt.	sandbar willow, or rope willow	FACW
<i>Salix lucida</i> Muhl. ssp. <i>lasiandra</i> (Benth.) E. Murray	Pacific willow	FACW
<i>Sambucus racemosa</i> L.	red elderberry	FACU

<i>Schoenoplectus acutus</i> (Muhl. ex Bigelow) A. Löve & D. Löve	hardstem bulrush	OBL
<i>Schoenoplectus americanus</i> (Pers.) Volkart ex Schinz & R. Keller (syn. <i>Scirpusolneyi</i> A. Gray)	Olney's three-square bulrush	FACW
<i>Scirpus microcarpus</i> J. Presl & C. Presl	small-flowered bulrush or "cut-grass"	OBL
<i>Sium suave</i> Walter	Water parsnip	OBL
<i>Sphagnum</i> spp.	Sphagnum moss or peat moss (several species)	OBL
<i>Spiraea douglasii</i> Hook.	hardhack	FACW
<i>Stachys chamissonis</i> Benth. var. <i>cooleyae</i> (A. Heller) G. Mulligan & D. Munro	coastal hedge-nettle	OBL
<i>Taxus brevifolia</i> Nutt.	Pacific yew or western yew	FACU
<i>Thuja plicata</i> Donn ex D. Don	western redcedar	FACU
<i>Tricholoma populinum</i> Lange	cottonwood mushroom	Not listed
<i>Trifolium wormskioldii</i> Lehm.	springbank clover	FACW
<i>Triglochin maritima</i> L.	seaside arrowgrass	OBL
<i>Typha latifolia</i> L.	cattail	OBL
<i>Viburnum. opulus</i> L.	American highbush cranberry or bush cranberry	FACW
<i>Vaccinium oxycoccus</i> L.	bog cranberry or small cranberry	OBL
<i>Vaccinium uliginosum</i> L.	bog bilberry	FACU
<i>Vaccinium vitis-idaea</i> L.	lingonberry or lowbush cranberry	FACU
<i>Veratrum viride</i> Aiton	false hellebore, or green false hellebore	FACU
<i>Viburnum edule</i> (Michx.) Raf.	highbush cranberry or squashberry	FACU

Sources: Turner, Nancy J. Food plants of coastal first peoples. Victoria: Royal BC Museum, 2010. Print.
Turner, Nancy J. Food plants of interior first peoples. Victoria: Royal BC Museum, 2007. Print.

Appendix 4 – Wetland Indicator Status Categories

Indicator Code	Indicator Status	Designation	Comment
OBL	Obligate Wetland	Hydrophyte	Almost always occur in wetlands.
FACW	Facultative Wetland	Hydrophyte	Usually occur in wetlands but may occur in non-wetlands.
FAC	Facultative	Hydrophyte	Occur in wetlands and non-wetlands.
FACU	Facultative Upland	Non-hydrophyte	Usually occur in non-wetlands but may occur in wetlands.
UPL	Obligate Upland	Non-hydrophyte	Almost never occur in wetlands.

Source: United States Department of Agriculture. 2018. Plants Database. Wetland Indicator Status.

<https://plants.usda.gov/wetinfo.html>

Appendix 5 Minimal Targets for Mature and Old Forest Coverage

Natural Disturbance Type	BEC & Variant	Upland-Wetland Objective	Minimal % Target for Mature + Old	Minimal Age
NDT1				
	CWH	High	54	80
	ICH	High	51	100
	ESSF	High	54	120
	MH	High	54	120
NDT2				
	CWH	Mod	34	80
	CDF	Mod	34	80
	ICH	Mod	31	100
	SBS	Mod	31	100
	ESSF	Mod	28	120
	SWB	Mod	28	120
NDT3				
	SBPS	Low	8	100
	SBSdk	Low	11	100
	SBSdw	Low	11	100
	SBSmk	Low	11	100
	SBSmc3	Low	11	100
	SBSwk1	Low	11	100
	SBS - other	High	34	100
	BWBSmw	Low	11	80/100*
	BWBSdk	Low	11	80/100*
	BWBSwk	Low	11	80/100*
	BWBS - other	High	34	80/100*
	MS - other	High	39	100
	MSxv	moderate-high	26	100
	ESSF	High	34	120
	ICH	High	34	100
	CWH	High	34	80
NDT4				
	ICH	high	51	100
	IDFdk	moderate-high	34	100
	IDF - other	high	51	100
	PP	high	51	100

*For BWBS, the minimal target age is 80 years for deciduous prominent & 100 years for coniferous prominent.

Source: Parminter, J. 1995. Biodiversity Guidebook. Forest Practices Code of British Columbia. B.C. Min. For. and B.C. Environ., Victoria, B.C.